
## Krishna K Mishra

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10640817/publications.pdf Version: 2024-02-01



Кріснил К Міснрл

#	Article	IF	CITATIONS
1	Virulence Gene-Associated Mutant Bacterial Colonies Generate Differentiating Two-Dimensional Laser Scatter Fingerprints. Applied and Environmental Microbiology, 2016, 82, 3256-3268.	3.1	17
2	Genetic organization and molecular characterization of secA2 locus in Listeria species. Gene, 2011, 489, 76-85.	2.2	22
3	N-Terminal Gly224–Gly411 Domain in Listeria Adhesion Protein Interacts with Host Receptor Hsp60. PLoS ONE, 2011, 6, e20694.	2.5	36
4	LAP, an alcohol acetaldehyde dehydrogenase enzyme in Listeria, promotes bacterial adhesion to enterocyte-like Caco-2 cells only in pathogenic species. Microbiology (United Kingdom), 2010, 156, 2782-2795.	1.8	100
5	Expression of LAP, a SecA2-dependent secretory protein, is induced under anaerobic environment. Microbes and Infection, 2009, 11, 859-867.	1.9	86
6	Coordinate regulation of a family of promastigote-enriched mRNAs by the 3′UTR PRE element in Leishmania mexicana. Molecular and Biochemical Parasitology, 2008, 157, 54-64.	1.1	20
7	Meiotic Reestablishment of Post-Transcriptional Gene Silencing is Regulated by Aberrant RNA Formation in Tomato (Lycopersicon esculentum cv. Mill.). Molecular Breeding, 2005, 16, 139-149.	2.1	3
8	A Negative Regulatory Element Controls mRNA Abundance of the Leishmania mexicana Paraflagellar Rod Gene PFR2. Eukaryotic Cell, 2003, 2, 1009-1017.	3.4	46
9	Post-transcriptional silencing of pectin methylesterase gene in transgenic tomato fruits results from impaired pre-mRNA processing. Plant Journal, 1998, 14, 583-592.	5.7	28
10	Modulation and recovery of immune response of BALB/c mice to <i>Shigella dysenteriae</i> antigens after cyclophosphamide treatment. Immunology and Cell Biology, 1994, 72, 419-426.	2.3	1