Luis Cartuche

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1065910/publications.pdf

Version: 2024-02-01

759233 794594 30 418 12 19 h-index citations g-index papers 30 30 30 609 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Enantiomeric Composition, Antioxidant Capacity and Anticholinesterase Activity of Essential Oil from Leaves of Chirimoya (Annona cherimola Mill.). Plants, 2022, 11, 367.	3.5	6
2	Chemical composition, enantiomeric analysis and anticholinesterase activity of <i>Lepechinia betonicifolia</i> essential oil from Ecuador. Pharmaceutical Biology, 2022, 60, 206-211.	2.9	6
3	Chemical Composition, Enantiomeric Distribution and Anticholinesterase and Antioxidant Activity of the Essential Oil of Diplosthephium juniperinum. Plants, 2022, 11, 1188.	3.5	6
4	Suillin: A mixed-type acetylcholinesterase inhibitor from Suillus luteus which is used by Saraguros indigenous, southern Ecuador. PLoS ONE, 2022, 17, e0268292.	2.5	8
5	Phytochemical profile, antimicrobial and antioxidant activities of essential oil extracted from Ecuadorian species <i>Piper ecuadorense</i> sodiro. Natural Product Research, 2021, 35, 6014-6019.	1.8	10
6	Study of Volatile Secondary Metabolites Present in Piper carpunya Leaves and in the Traditional Ecuadorian Beverage Guaviduca. Plants, 2021, 10, 338.	3.5	13
7	Chemical Constituents of the Essential Oil from Ecuadorian Endemic Species Croton ferrugineus and Its Antimicrobial, Antioxidant and α-Glucosidase Inhibitory Activity. Molecules, 2021, 26, 4608.	3.8	12
8	Variability of the Chemical Composition and Bioactivity between the Essential Oils Isolated from Male and Female Specimens of Hedyosmum racemosum (Ruiz & Pav.) G. Don. Molecules, 2021, 26, 4613.	3.8	5
9	A Rare Dirhamnosyl Flavonoid and Other Radicalâ€Scavenging Metabolites from <i>Cynophalla mollis</i> (Kunth) J.â€Presl and <i>Colicodendron scabridum</i> (Kunt) Seem. (Capparaceae) of Ecuador. Chemistry and Biodiversity, 2021, 18, e2100260.	2.1	6
10	Extraction and Study of the Essential Oil of Copal (Dacryodes peruviana), an Amazonian Fruit with the Highest Yield Worldwide. Plants, 2020, 9, 1658.	3.5	20
11	Evaluation of Indolocarbazoles from Streptomyces sanyensis as a Novel Source of Therapeutic Agents against the Brain-Eating Amoeba Naegleria fowleri. Microorganisms, 2020, 8, 789.	3.6	13
12	Antikinetoplastid Activity of Indolocarbazoles from Streptomyces sanyensis. Biomolecules, 2020, 10, 657.	4.0	24
13	Antiamoebic Activities of Indolocarbazole Metabolites Isolated from Streptomyces sanyensis Cultures. Marine Drugs, 2019, 17, 588.	4.6	11
14	Staurosporine from Streptomyces sanyensis activates Programmed Cell Death in Acanthamoeba via the mitochondrial pathway and presents low in vitro cytotoxicity levels in a macrophage cell line. Scientific Reports, 2019, 9, 11651.	3.3	27
15	Novel Flavonoid Glycosides of Quercetin from Leaves and Flowers of Gaiadendron punctatum G.Don. (Violeta de Campo), used by the Saraguro Community in Southern Ecuador, Inhibit α-Glucosidase Enzyme. Molecules, 2019, 24, 4267.	3.8	13
16	Chemical Constituents of Croton thurifer Kunth as \hat{l} ±-Glucosidase Inhibitors. Records of Natural Products, 2019, 14, 31-41.	1.3	5
17	Inactivation of a wild isolated Klebsiella pneumoniae by photo-chemical processes: UV-C, UV-C/H2O2 and UV-C/H2O2/Fe3+. Catalysis Today, 2018, 313, 94-99.	4.4	22
18	PRELIMINARY PHYTOCHEMICAL STUDY OF THE ECUADORIAN PLANT CROTON ELEGANS KUNTH (EUPHORBIACEAE). Journal of the Chilean Chemical Society, 2018, 63, 3875-3877.	1.2	11

#	Article	lF	CITATIONS
19	α-Glucosidase Inhibition and Antibacterial Activity of Secondary Metabolites from the Ecuadorian Species Clinopodium taxifolium (Kunth) Govaerts. Molecules, 2018, 23, 146.	3.8	16
20	Chemical composition and antimicrobial activity of Myrcianthes fragrans essential oil, a natural aromatizer of the traditional Ecuadorian beverage colada morada. Journal of Ethnopharmacology, 2018, 225, 319-326.	4.1	12
21	Acorenone B: AChE and BChE Inhibitor as a Major Compound of the Essential Oil Distilled from the Ecuadorian Species Niphogeton dissecta (Benth.) J.F. Macbr. Pharmaceuticals, 2017, 10, 84.	3.8	25
22	Main Constituents and Antidiabetic Properties of <i>Otholobium mexicanum</i> . Natural Product Communications, 2017, 12, 1934578X1701200.	0.5	3
23	Chemical Constituents of Muehlenbeckia tamnifolia (Kunth) Meisn (Polygonaceae) and Its In Vitro α-Amilase and α-Glucosidase Inhibitory Activities. Molecules, 2016, 21, 1461.	3.8	65
24	Biological Activity and Chemical Composition of the Essential Oil fromChromolaena laevigata(Lam.) R.M. King & H. Rob. (Asteraceae) from Loja, Ecuador. Journal of Essential Oil-bearing Plants: JEOP, 2016, 19, 384-390.	1.9	4
25	Antibacterial and cytotoxic activity from the extract and fractions of a marine derived bacterium from the Streptomycesgenus. Pharmaceutical Biology, 2015, 53, 1826-1830.	2.9	4
26	Chemical composition and biological activity of the essential oil of <i>Baccharis obtusifolia</i> Kunth from Loja, Ecuador. Journal of Essential Oil Research, 2015, 27, 212-216.	2.7	17
27	Antifungal activity of raw extract and flavanons isolated from Piper ecuadorense from Ecuador. Revista Brasileira De Farmacognosia, 2013, 23, 370-373.	1.4	17
28	Chemical composition, antifungal and antibacterial activity of the essential oil from <i>Baccharis latifolia </i> i (Ruiz & amp; Pav.) Pers. (Asteraceae) from Loja, Ecuador. Journal of Essential Oil Research, 2013, 25, 233-238.	2.7	15
29	Preparation, Characterization and Antibacterial Activity of Poly(<l>ε</l> -caprolactone) Electrospun Fibers Loaded with Amoxicillin for Controlled Release in Biomedical Applications. Journal of Nanoscience and Nanotechnology, 2013, 13, 1717-1726.	0.9	16
30	Study of the essential oil from native amazonian species of Ecuador Piper lineatum, presence of apiole and safrole. Natural Volatiles and Essential Oils (discontinued), 0, , .	1.1	6