Patrick Rodriguez

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10664793/publications.pdf

Version: 2024-02-01

840776 1281871 14 1,668 11 11 citations h-index g-index papers 14 14 14 2730 docs citations times ranked citing authors all docs

#	Article	IF	Citations
1	A phospho-dependent mechanism involving NCoR and KMT2D controls a permissive chromatin state at Notch target genes. Nucleic Acids Research, 2016, 44, 4703-4720.	14.5	77
2	BCL9/9L-Î ² -catenin Signaling is Associated With Poor Outcome in Colorectal Cancer. EBioMedicine, 2015, 2, 1932-1943.	6.1	58
3	Histone demethylase KDM5A is an integral part of the core Notch–RBP-J repressor complex. Genes and Development, 2010, 24, 590-601.	5.9	162
4	Proteomics Analysis of Ring1B/Rnf2 Interactors Identifies a Novel Complex with the Fbxl10/Jhdm1B Histone Demethylase and the Bcl6 Interacting Corepressor. Molecular and Cellular Proteomics, 2007, 6, 820-834.	3.8	202
5	Isolation of Transcription Factor Complexes by In Vivo Biotinylation Tagging and Direct Binding to Streptavidin Beads., 2006, 338, 305-323.		25
6	Novel binding partners of Ldb1 are required for haematopoietic development. Development (Cambridge), 2006, 133, 4913-4923.	2.5	115
7	ETO2 coordinates cellular proliferation and differentiation during erythropoiesis. EMBO Journal, 2006, 25, 357-366.	7.8	126
8	The mediator complex functions as a coactivator for GATA-1 in erythropoiesis via subunit Med1/TRAP220. Proceedings of the National Academy of Sciences of the United States of America, 2006, 103, 18504-18509.	7.1	89
9	Dynamic Change in the Stochiometry of ETO2 and SCL Governs the Transition to Terminal Differentiation in Erythroid Progenitors Blood, 2006, 108, 1169-1169.	1.4	O
10	Isolation and Characterization of Hematopoietic Transcription Factor Complexes byin VivoBiotinylation Tagging and Mass Spectrometry. Annals of the New York Academy of Sciences, 2005, 1054, 55-67.	3.8	29
11	GATA-1 forms distinct activating and repressive complexes in erythroid cells. EMBO Journal, 2005, 24, 2354-2366.	7.8	255
12	ETO-2 Associates with SCL in Erythroid Cells and Megakaryocytes and Provides Repressor Functions in Erythropoiesis. Molecular and Cellular Biology, 2005, 25, 10235-10250.	2.3	130
13	GATA-1 Forms Distinct Activating and Repressive Complexes in Erythroid Cells Blood, 2004, 104, 356-356.	1.4	0
14	Efficient biotinylation and single-step purification of tagged transcription factors in mammalian cells and transgenic mice. Proceedings of the National Academy of Sciences of the United States of America, 2003, 100, 7480-7485.	7.1	400