
Réjane Streiff

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10669117/publications.pdf Version: 2024-02-01



RÃOIANE STREIFE

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Within-population genetic structure in Quercus robur L. and Quercus petraea (Matt.) Liebl. assessed with isozymes and microsatellites. Molecular Ecology, 1998, 7, 317-328. | 3.9 | 299 |
| 2 | Genomics of adaptation to host-plants in herbivorous insects. Briefings in Functional Genomics, 2015, 14, 413-423. | 2.7 | 135 |
| 3 | Comparative study of genetic variation and differentiation of two pedunculate oak (Quercus robur) stands using microsatellite and allozyme loci. Heredity, 1999, 83, 597-603. | 2.6 | 48 |
| 4 | â€~Becoming a species by becoming a pest' or how two maize pests of the genus <i>Ostrinia</i> possibly evolved through parallel ecological speciation events. Molecular Ecology, 2014, 23, 325-342. | 3.9 | 46 |
| 5 | Genetic Architecture of Sexual Selection: QTL Mapping of Male Song and Female Receiver Traits in an Acoustic Moth. PLoS ONE, 2012, 7, e44554. | 2.5 | 30 |
| 6 | Scanning the European corn borer (<i>Ostrinia</i> spp.) genome for adaptive divergence between hostâ€affiliated sibling species. Molecular Ecology, 2011, 20, 1414-1430. | 3.9 | 29 |
| 7 | Animal choruses emerge from receiver psychology. Scientific Reports, 2016, 6, 34369. | 3.3 | 20 |
| 8 | Group synchrony and alternation as an emergent property: elaborate chorus structure in a bushcricket is an incidental by-product of female preference for leading calls. Behavioral Ecology and Sociobiology, 2015, 69, 1957-1973. | 1.4 | 19 |
| 9 | When History Repeats Itself: Exploring the Genetic Architecture of Host-Plant Adaptation in Two Closely Related Lepidopteran Species. PLoS ONE, 2013, 8, e69211. | 2.5 | 13 |
| 10 | MODELING SURVIVAL AND MARK LOSS IN MOLTING ANIMALS: RECAPTURE, DEAD RECOVERIES, AND EXUVIA RECOVERIES. Ecology, 2007, 88, 289-295. | 3.2 | 11 |
| 11 | De novo transcriptomic resources for two sibling species of moths: Ostrinia nubilalis and O. scapulalis. BMC Research Notes, 2013, 6, 73. | 1.4 | 9 |
| 12 | Organisation spatiale de la diversité génétique et flux polliniques dans une chênaie mixte. Genetics Selection Evolution, 1998, 30, 1. | 3.0 | 6 |
| 13 | Genetic mapping of two components of reproductive isolation between two sibling species of moths, Ostrinia nubilalis and O. scapulalis. Heredity, 2014, 112, 370-381. | 2.6 | 6 |
| 14 | Fine-scale interactions between habitat quality and genetic variation suggest an impact of grazing on the critically endangered Crau Plain grasshopper (Pamphagidae: Prionotropis rhodanica). Journal of Orthoptera Research, 2018, 27, 61-73. | 1.0 | 4 |
| 15 | Characterization of 16 novel microsatellite loci for Ephippiger diurnus (Orthoptera: Tettigoniidae) using pyrosequencing technology and cross-species amplification. European Journal of Entomology, 0, 113, 302-306. | 1.2 | 3 |
| | | | |

Pleistocene origins of chorusing diversity in Mediterranean bush-cricket populations (Ephippiger) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 1