Réjane Streiff

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10669117/publications.pdf

Version: 2024-02-01

		1040056 996975	
16	679	9	15
papers	citations	h-index	g-index
16	16	16	942
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	Pleistocene origins of chorusing diversity in Mediterranean bush-cricket populations (Ephippiger) Tj ETQq1 1 0.78	14314 rgBT	- Overlock 1
2	Fine-scale interactions between habitat quality and genetic variation suggest an impact of grazing on the critically endangered Crau Plain grasshopper (Pamphagidae: Prionotropis rhodanica). Journal of Orthoptera Research, 2018, 27, 61-73.	1.0	4
3	Animal choruses emerge from receiver psychology. Scientific Reports, 2016, 6, 34369.	3.3	20
4	Group synchrony and alternation as an emergent property: elaborate chorus structure in a bushcricket is an incidental by-product of female preference for leading calls. Behavioral Ecology and Sociobiology, 2015, 69, 1957-1973.	1.4	19
5	Genomics of adaptation to host-plants in herbivorous insects. Briefings in Functional Genomics, 2015, 14, 413-423.	2.7	135
6	â€Becoming a species by becoming a pest' or how two maize pests of the genus <i>Ostrinia</i> possibly evolved through parallel ecological speciation events. Molecular Ecology, 2014, 23, 325-342.	3.9	46
7	Genetic mapping of two components of reproductive isolation between two sibling species of moths, Ostrinia nubilalis and O. scapulalis. Heredity, 2014, 112, 370-381.	2.6	6
8	De novo transcriptomic resources for two sibling species of moths: Ostrinia nubilalis and O. scapulalis. BMC Research Notes, 2013, 6, 73.	1.4	9
9	When History Repeats Itself: Exploring the Genetic Architecture of Host-Plant Adaptation in Two Closely Related Lepidopteran Species. PLoS ONE, 2013, 8, e69211.	2.5	13
10	Genetic Architecture of Sexual Selection: QTL Mapping of Male Song and Female Receiver Traits in an Acoustic Moth. PLoS ONE, 2012, 7, e44554.	2.5	30
11	Scanning the European corn borer (<i>Ostrinia</i> spp.) genome for adaptive divergence between hostâ€affiliated sibling species. Molecular Ecology, 2011, 20, 1414-1430.	3.9	29
12	MODELING SURVIVAL AND MARK LOSS IN MOLTING ANIMALS: RECAPTURE, DEAD RECOVERIES, AND EXUVIA RECOVERIES. Ecology, 2007, 88, 289-295.	3.2	11
13	Comparative study of genetic variation and differentiation of two pedunculate oak (Quercus robur) stands using microsatellite and allozyme loci. Heredity, 1999, 83, 597-603.	2.6	48
14	Organisation spatiale de la diversité génétique et flux polliniques dans une chênaie mixte. Genetics Selection Evolution, 1998, 30, 1.	3.0	6
15	Within-population genetic structure in Quercus robur L. and Quercus petraea (Matt.) Liebl. assessed with isozymes and microsatellites. Molecular Ecology, 1998, 7, 317-328.	3.9	299
16	Characterization of 16 novel microsatellite loci for Ephippiger diurnus (Orthoptera: Tettigoniidae) using pyrosequencing technology and cross-species amplification. European Journal of Entomology, 0, 113, 302-306.	1.2	3