

David Cox

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10670214/publications.pdf>

Version: 2024-02-01

6
papers

448
citations

1477746
6
h-index

1872312
6
g-index

6
all docs

6
docs citations

6
times ranked

421
citing authors

#	ARTICLE	IF	CITATIONS
1	Monotherapy with Tadalafil or Tamsulosin Similarly Improved Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia in an International, Randomised, Parallel, Placebo-Controlled Clinical Trial. <i>European Urology</i> , 2012, 61, 917-925.	0.9	253
2	Efficacy and Safety of Tadalafil 5 mg Once Daily for Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia: Subgroup Analyses of Pooled Data From 4 Multinational, Randomized, Placebo-controlled Clinical Studies. <i>Urology</i> , 2013, 82, 667-673.	0.5	59
3	Tadalafil Once Daily Improves Ejaculatory Function, Erectile Function, and Sexual Satisfaction in Men with Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia and Erectile Dysfunction: Results from a Randomized, Placebo- and Tamsulosin-Controlled, 12-Week Double-Blind Study. <i>Journal of Sexual Medicine</i> , 2013, 10, 857-865.	0.3	50
4	Effects of Tadalafil Once Daily on Maximum Urinary Flow Rate in Men with Lower Urinary Tract Symptoms Suggestive of Benign Prostatic Hyperplasia. <i>Journal of Urology</i> , 2014, 191, 1045-1050.	0.2	35
5	Effects of tadalafil on nighttime voiding (nocturia) in men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia: a post hoc analysis of pooled data from four randomized, placebo-controlled clinical studies. <i>World Journal of Urology</i> , 2014, 32, 1127-1132.	1.2	30
6	Effect of Tadalafil Once Daily on Prostate Blood Flow and Perfusion in Men With Lower Urinary Tract Symptoms Secondary to Benign Prostatic Hyperplasia: A Randomized, Double-blind, Multicenter, Placebo-controlled Trial. <i>Urology</i> , 2014, 84, 412-420.	0.5	21