

# David Cox

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/10670214/david-cox-publications-by-year.pdf>

**Version:** 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

6

papers

359

citations

6

h-index

6

g-index

6

ext. papers

411

ext. citations

3.2

avg, IF

2.8

L-index

#	Paper	IF	Citations
6	Effects of tadalafil on nighttime voiding (nocturia) in men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia: a post hoc analysis of pooled data from four randomized, placebo-controlled clinical studies. <i>World Journal of Urology</i> , <b>2014</b> , 32, 1127-32	4	26
5	Effect of tadalafil once daily on prostate blood flow and perfusion in men with lower urinary tract symptoms secondary to benign prostatic hyperplasia: a randomized, double-blind, multicenter, placebo-controlled trial. <i>Urology</i> , <b>2014</b> , 84, 412-9	1.6	19
4	Effects of tadalafil once daily on maximum urinary flow rate in men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia. <i>Journal of Urology</i> , <b>2014</b> , 191, 1045-50	2.5	26
3	Efficacy and safety of tadalafil 5 mg once daily for lower urinary tract symptoms suggestive of benign prostatic hyperplasia: subgroup analyses of pooled data from 4 multinational, randomized, placebo-controlled clinical studies. <i>Urology</i> , <b>2013</b> , 82, 667-73	1.6	47
2	Tadalafil once daily improves ejaculatory function, erectile function, and sexual satisfaction in men with lower urinary tract symptoms suggestive of benign prostatic hyperplasia and erectile dysfunction: results from a randomized, placebo- and tamsulosin-controlled, 12-week double-blind trial. <i>Urology</i> , <b>2013</b> , 82, 677-85	1.1	41
1	Monotherapy with tadalafil or tamsulosin similarly improved lower urinary tract symptoms suggestive of benign prostatic hyperplasia in an international, randomised, parallel, placebo-controlled clinical trial. <i>European Urology</i> , <b>2012</b> , 61, 917-25	10.2	200