

# Jaime Molina-Ochoa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10679835/publications.pdf>

Version: 2024-02-01

13  
papers

470  
citations

933447

10  
h-index

1125743

13  
g-index

13  
all docs

13  
docs citations

13  
times ranked

512  
citing authors

#	ARTICLE	IF	CITATIONS
1	Genetic Variability of Spined Soldier Bugs (Hemiptera: Pentatomidae) Sampled from Distinct Field Sites and Laboratory Colonies in the United States. Florida Entomologist, 2014, 97, 1631-1639.	0.5	2
2	Survey of entomopathogenic nematodes from the families Steinernematidae and Heterorhabditidae (Nematoda: Rhabditida) in Colima, MÃ©xico. International Journal of Tropical Insect Science, 2014, 34, 53-57.	1.0	5
3	Occurrence of Entomopathogenic Fungi and Parasitic Nematodes on <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) Larvae Collected in Central Chiapas, MÃ©xico. Florida Entomologist, 2013, 96, 498-503.	0.5	24
4	Use of the entomopathogenic fungi <i>Metarhizium anisopliae</i> , <i>Cordyceps bassiana</i> and <i>Isaria fumosorosea</i> to control <i>Diaphorina citri</i> (Hemiptera: Psyllidae) in Persian lime under field conditions. International Journal of Tropical Insect Science, 2012, 32, 39-44.	1.0	43
5	Spatial Genetic Variation among <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) Sampled from the United States, Puerto Rico, Panama, and Argentina. Annals of the Entomological Society of America, 2012, 105, 359-367.	2.5	12
6	Population genetic structure of two primary parasitoids of <i>Spodoptera frugiperda</i> (Lepidoptera), <i>Chelonus insularis</i> and <i>Campoletis sonorensis</i> (Hymenoptera): to what extent is the host plant important?. Molecular Ecology, 2010, 19, 2168-2179.	3.9	12
7	<i>Steinernema diaprepesi</i> (Nematoda: Steinernematidae): Its Occurrence in Western Mexico and Susceptibility of Engorged Cattle Ticks <i>Boophilus microplus</i> (Acari: Ixodidae). Florida Entomologist, 2009, 92, 661.	0.5	12
8	Lyophilization of <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) larvae yields high-quality DNA for use in AFLP genetic studies. International Journal of Tropical Insect Science, 2009, 29, 79.	1.0	2
9	Population Variation of the Fall Armyworm, <i>Spodoptera frugiperda</i> , in the Western Hemisphere. Journal of Insect Science, 2007, 7, 1-10.	1.5	116
10	Survey for Hymenopteran and Dipteran Parasitoids of the Fall Armyworm (Lepidoptera: Noctuidae) in Chiapas, Mexico. Journal of Agricultural and Urban Entomology, 2007, 24, 35-42.	0.6	36
11	PATHOGENS AND PARASITIC NEMATODES ASSOCIATED WITH POPULATIONS OF FALL ARMYWORM (LEPIDOPTERA: NOCTUIDAE) LARVAE IN MEXICO. Florida Entomologist, 2003, 86, 244-253.	0.5	36
12	PARASITIDS AND PARASITES OF SPODOPTERA FRUGIPERDA (LEPIDOPTERA: NOCTUIDAE) IN THE AMERICAS AND CARIBBEAN BASIN: AN INVENTORY. Florida Entomologist, 2003, 86, 254-289.	0.5	140
13	Occurrence of Entomopathogens of <i>Spodoptera frugiperda</i> (Lepidoptera: Noctuidae) in the Mexican States of MichoacÃ¡n, Colima, Jalisco and Tamaulipas. Florida Entomologist, 2001, 84, 23.	0.5	30