

Jiangye Xu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1072794/publications.pdf>

Version: 2024-02-01

21
papers

1,113
citations

567281

15
h-index

713466

21
g-index

24
all docs

24
docs citations

24
times ranked

1132
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Hypoxic glioma-derived exosomes deliver microRNA-1246 to induce M2 macrophage polarization by targeting TERF2IP via the STAT3 and NF- κ B pathways. <i>Oncogene</i> , 2020, 39, 428-442. | 5.9 | 223 |
| 2 | Exosomes derived from mesenchymal stem cells attenuate the progression of atherosclerosis in ApoE ^{-/-} mice via miR-let7 mediated infiltration and polarization of M2 macrophage. <i>Biochemical and Biophysical Research Communications</i> , 2019, 510, 565-572. | 2.1 | 136 |
| 3 | EWSR1-induced circNEIL3 promotes glioma progression and exosome-mediated macrophage immunosuppressive polarization via stabilizing IGF2BP3. <i>Molecular Cancer</i> , 2022, 21, 16. | 19.2 | 115 |
| 4 | Glioma exosomes mediate the expansion and function of myeloid-derived suppressor cells through microRNA-29a and Hbp1 and microRNA-92a and Prkar1a pathways. <i>International Journal of Cancer</i> , 2019, 144, 3111-3126. | 5.1 | 107 |
| 5 | Hypoxic glioma-derived exosomes promote M2-like macrophage polarization by enhancing autophagy induction. <i>Cell Death and Disease</i> , 2021, 12, 373. | 6.3 | 93 |
| 6 | Transfer of MicroRNA via Macrophage-Derived Extracellular Vesicles Promotes Proneural-to-Mesenchymal Transition in Glioma Stem Cells. <i>Cancer Immunology Research</i> , 2020, 8, 966-981. | 3.4 | 55 |
| 7 | Allosteric mechanisms underlie GPCR signaling to SH3-domain proteins through arrestin. <i>Nature Chemical Biology</i> , 2018, 14, 876-886. | 8.0 | 50 |
| 8 | Cullin-7 (CUL7) is overexpressed in glioma cells and promotes tumorigenesis via NF- κ B activation. <i>Journal of Experimental and Clinical Cancer Research</i> , 2020, 39, 59. | 8.6 | 41 |
| 9 | Hypoxia-induced lncRNA PDIA3P1 promotes mesenchymal transition via sponging of miR-124-3p in glioma. <i>Cell Death and Disease</i> , 2020, 11, 168. | 6.3 | 40 |
| 10 | Mucin O-glycosylating enzyme GALNT2 facilitates the malignant character of glioma by activating the EGFR/PI3K/Akt/mTOR axis. <i>Clinical Science</i> , 2019, 133, 1167-1184. | 4.3 | 37 |
| 11 | MicroRNA-29a-3p delivery via exosomes derived from engineered human mesenchymal stem cells exerts tumour suppressive effects by inhibiting migration and vasculogenic mimicry in glioma. <i>Aging</i> , 2021, 13, 5055-5068. | 3.1 | 37 |
| 12 | The dual role of glioma exosomal microRNAs: glioma eliminates tumor suppressor miR-1298-5p via exosomes to promote immunosuppressive effects of MDSCs. <i>Cell Death and Disease</i> , 2022, 13, 426. | 6.3 | 32 |
| 13 | Exosomes derived from hypoxic glioma deliver miR-1246 and miR-10b-5p to normoxic glioma cells to promote migration and invasion. <i>Laboratory Investigation</i> , 2021, 101, 612-624. | 3.7 | 28 |
| 14 | SPI1-induced downregulation of FTO promotes GBM progression by regulating pri-miR-10a processing in an m6A-dependent manner. <i>Molecular Therapy - Nucleic Acids</i> , 2022, 27, 699-717. | 5.1 | 23 |
| 15 | PDIA3P1 promotes Temozolomide resistance in glioblastoma by inhibiting C/EBP β degradation to facilitate proneural-to-mesenchymal transition. <i>Journal of Experimental and Clinical Cancer Research</i> , 2022, 41, . | 8.6 | 20 |
| 16 | Cell surface GRP78 regulates BACE2 via lysosome-dependent manner to maintain mesenchymal phenotype of glioma stem cells. <i>Journal of Experimental and Clinical Cancer Research</i> , 2021, 40, 20. | 8.6 | 17 |
| 17 | <p>GALE Promotes the Proliferation and Migration of Glioblastoma Cells and Is Regulated by miR-let-7i-5p</p>. <i>Cancer Management and Research</i> , 2019, Volume 11, 10539-10554. | 1.9 | 15 |
| 18 | A Comprehensive Analysis of METTL1 to Immunity and Stemness in Pan-Cancer. <i>Frontiers in Immunology</i> , 2022, 13, 795240. | 4.8 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | PLEKHG5 is a novel prognostic biomarker in glioma patients. International Journal of Clinical Oncology, 2019, 24, 1350-1358. | 2.2 | 14 |
| 20 | The Non-N6-Methyladenosine Epitranscriptome Patterns and Characteristics of Tumor Microenvironment Infiltration and Mesenchymal Transition in Glioblastoma. Frontiers in Immunology, 2021, 12, 809808. | 4.8 | 11 |
| 21 | Nitidine Chloride Is a Potential Alternative Therapy for Glioma Through Inducing Endoplasmic Reticulum Stress and Alleviating Epithelial-Mesenchymal Transition. Integrative Cancer Therapies, 2020, 19, 153473541990092. | 2.0 | 4 |