

Scott G Clark

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10731827/publications.pdf>

Version: 2024-02-01

15
papers

5,376
citations

623734

14
h-index

1058476

14
g-index

15
all docs

15
docs citations

15
times ranked

6123
citing authors

#	ARTICLE	IF	CITATIONS
1	The conserved transmembrane RING finger protein PLR-1 downregulates Wnt signaling by reducing Frizzled, Ror and Ryk cell-surface levels in <i>C. elegans</i> . <i>Development (Cambridge)</i> , 2014, 141, 617-628.	2.5	36
2	Axons Degenerate in the Absence of Mitochondria in <i>C. elegans</i> . <i>Current Biology</i> , 2014, 24, 760-765.	3.9	86
3	<i>C. elegans</i> AP-2 and Retromer Control Wnt Signaling by Regulating MIG-14/Wntless. <i>Developmental Cell</i> , 2008, 14, 132-139.	7.0	189
4	Multiple Wnts and Frizzled Receptors Regulate Anteriorly Directed Cell and Growth Cone Migrations in <i>Caenorhabditis elegans</i> . <i>Developmental Cell</i> , 2006, 10, 367-377.	7.0	151
5	Wnt signaling establishes anteroposterior neuronal polarity and requires retromer in <i>C. elegans</i> . <i>Development (Cambridge)</i> , 2006, 133, 1757-1766.	2.5	199
6	Compartment-specific perturbation of protein handling activates genes encoding mitochondrial chaperones. <i>Journal of Cell Science</i> , 2004, 117, 4055-4066.	2.0	522
7	<i>C. elegans</i> ZAG-1, a Zn-finger-homeodomain protein, regulates axonal development and neuronal differentiation. <i>Development (Cambridge)</i> , 2003, 130, 3781-3794.	2.5	118
8	A survival pathway for <i>Caenorhabditis elegans</i> with a blocked unfolded protein response. <i>Journal of Cell Biology</i> , 2002, 158, 639-646.	5.2	181
9	IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. <i>Nature</i> , 2002, 415, 92-96.	27.8	2,452
10	PAG-3, a Zn-finger transcription factor, determines neuroblast fate in <i>C. elegans</i> . <i>Development (Cambridge)</i> , 2002, 129, 1763-1774.	2.5	34
11	Gain-of-Function Mutations in the <i>Caenorhabditis elegans</i> lin-1 ETS Gene Identify a C-Terminal Regulatory Domain Phosphorylated by ERK MAP Kinase. <i>Genetics</i> , 1998, 149, 1809-1822.	2.9	106
12	Control of cell fates in the central body region of <i>C. elegans</i> by the homeobox gene lin-39. <i>Cell</i> , 1993, 74, 43-55.	28.9	261
13	<i>C. elegans</i> cell-signalling gene sem-5 encodes a protein with SH2 and SH3 domains. <i>Nature</i> , 1992, 356, 340-344.	27.8	632
14	<i>Caenorhabditis elegans</i> ras gene let-60 acts as a switch in the pathway of vulval induction. <i>Nature</i> , 1990, 348, 503-509.	27.8	408
15	IRE1 couples endoplasmic reticulum load to secretory capacity by processing the XBP-1 mRNA. , 0, .		1