

Bernardo Trigatti

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10740778/publications.pdf>

Version: 2024-02-01

13
papers

1,298
citations

932766

10
h-index

1281420

11
g-index

13
all docs

13
docs citations

13
times ranked

1174
citing authors

#	ARTICLE	IF	CITATIONS
1	Loss-of-function PCSK9 mutants evade the unfolded protein response sensor GRP78 and fail to induce endoplasmic reticulum stress when retained. <i>Journal of Biological Chemistry</i> , 2018, 293, 7329-7343.	1.6	29
2	Intestinal scavenger receptor class B type I as a novel regulator of chylomicron production in healthy and diet-induced obese states. <i>American Journal of Physiology - Renal Physiology</i> , 2015, 309, G350-G359.	1.6	21
3	The Scavenger Receptor Class B Type I. , 2010, , 153-178.		0
4	Hepatic uptake and metabolism of phosphatidylcholine associated with high density lipoproteins. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 2009, 1790, 538-551.	1.1	28
5	Conversion of Low Density Lipoprotein-associated Phosphatidylcholine to Triacylglycerol by Primary Hepatocytes. <i>Journal of Biological Chemistry</i> , 2008, 283, 6449-6458.	1.6	33
6	Myocardial Infarction Following Atherosclerosis in Murine Models. <i>Current Drug Targets</i> , 2008, 9, 217-223.	1.0	9
7	Hepatic steatosis and inflammation: Role of western diet and lipoproteins. <i>FASEB Journal</i> , 2007, 21, A846.	0.2	0
8	Hepatic Lipase Deficiency Delays Atherosclerosis, Myocardial Infarction, and Cardiac Dysfunction and Extends Lifespan in SR-BI/Apolipoprotein E Double Knockout Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2006, 26, 548-554.	1.1	37
9	Hepatic cholesterol and bile acid metabolism and intestinal cholesterol absorption in scavenger receptor class B type I-deficient mice. <i>Journal of Lipid Research</i> , 2001, 42, 170-180.	2.0	256
10	The role of the high-density lipoprotein receptor SR-BI in cholesterol metabolism. <i>Current Opinion in Lipidology</i> , 2000, 11, 123-131.	1.2	172
11	The Efficient Cellular Uptake of High Density Lipoprotein Lipids via Scavenger Receptor Class B Type I Requires Not Only Receptor-mediated Surface Binding but Also Receptor-specific Lipid Transfer Mediated by Its Extracellular Domain. <i>Journal of Biological Chemistry</i> , 1998, 273, 26338-26348.	1.6	198
12	Murine SR-BI, a High Density Lipoprotein Receptor That Mediates Selective Lipid Uptake, Is N-Glycosylated and Fatty Acylated and Colocalizes with Plasma Membrane Caveolae. <i>Journal of Biological Chemistry</i> , 1997, 272, 13242-13249.	1.6	330
13	Scavenger receptor BI - a cell surface receptor for high density lipoprotein. <i>Current Opinion in Lipidology</i> , 1997, 8, 181-188.	1.2	185