

# Jyrki Tenhunen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10743571/publications.pdf>

Version: 2024-02-01

46  
papers

3,576  
citations

361296  
20  
h-index

265120  
42  
g-index

47  
all docs

47  
docs citations

47  
times ranked

4584  
citing authors

#	ARTICLE	IF	CITATIONS
1	Mono-(2-ethylhexyl) phthalate Promotes Dengue Virus Infection by Decreasing IL-23-Mediated Antiviral Responses. <i>Frontiers in Immunology</i> , 2021, 12, 599345.	2.2	5
2	Never quite there? â€” Hyperventilation in cardiopulmonary resuscitation. <i>Resuscitation</i> , 2021, 165, 138-139.	1.3	2
3	Plasma hyaluronan, hyaluronidase activity and endogenous hyaluronidase inhibition in sepsis: an experimental and clinical cohort study. <i>Intensive Care Medicine Experimental</i> , 2021, 9, 53.	0.9	3
4	Endotoxin Removal in Septic Shock with the Alteco LPS Adsorber Was Safe But Showed no Benefit Compared to Placebo in the Double-Blind Randomized Controlled Trialâ€™the Asset Study. <i>Shock</i> , 2020, 54, 224-231.	1.0	11
5	The antisecretory peptide AF-16 may modulate tissue edema but not inflammation in experimental peritonitis induced sepsis. <i>PLoS ONE</i> , 2020, 15, e0232302.	1.1	1
6	Title is missing!. , 2020, 15, e0232302.		0
7	Title is missing!. , 2020, 15, e0232302.		0
8	Title is missing!. , 2020, 15, e0232302.		0
9	Title is missing!. , 2020, 15, e0232302.		0
10	High levels of serum hyaluronan is an early predictor of dengue warning signs and perturbs vascular integrity. <i>EBioMedicine</i> , 2019, 48, 425-441.	2.7	29
11	S100B, NSE and MMP-9 fail to predict neurologic outcome while elevated S100B associates with milder initial clinical presentation after aneurysmal subarachnoid hemorrhage. <i>Journal of the Neurological Sciences</i> , 2018, 390, 129-134.	0.3	9
12	Time-courses of plasma IL-6 and HMGB-1 reflect initial severity of clinical presentation but do not predict poor neurologic outcome following subarachnoid hemorrhage. <i>ENeurologicalSci</i> , 2017, 6, 55-62.	0.5	14
13	Successful management of superâ€™refractory status epilepticus with thalamic deep brain stimulation. <i>Annals of Neurology</i> , 2017, 81, 142-146.	2.8	36
14	Plasma Soluble Urokinase-Type Plasminogen Activator Receptor Is Not Associated with Neurological Outcome in Patients with Aneurysmal Subarachnoid Hemorrhage. <i>Frontiers in Neurology</i> , 2017, 8, 144.	1.1	5
15	HMGB1 and Extracellular Histones Significantly Contribute to Systemic Inflammation and Multiple Organ Failure in Acute Liver Failure. <i>Mediators of Inflammation</i> , 2017, 2017, 1-6.	1.4	56
16	HMGB1 and Histones Play a Significant Role in Inducing Systemic Inflammation and Multiple Organ Dysfunctions in Severe Acute Pancreatitis. <i>International Journal of Inflammation</i> , 2017, 2017, 1-6.	0.9	46
17	Increased plasma UCH-L1 after aneurysmal subarachnoid hemorrhage is associated with unfavorable neurologic outcome. <i>Journal of the Neurological Sciences</i> , 2016, 361, 144-149.	0.3	15
18	Abdominal Septic Shock â€™ Endotoxin Adsorption Treatment (ASSET) â€™ endotoxin removal in abdominal and urogenital septic shock with the Altecoâ€™ LPS Adsorber: study protocol for a double-blinded, randomized placebo-controlled trial. <i>Trials</i> , 2016, 17, 587.	0.7	14

#	ARTICLE	IF	CITATIONS
19	Simultaneous beat-to-beat assessment of arterial blood pressure and quality of cardiopulmonary resuscitation in out-of-hospital and in-hospital settings. <i>Resuscitation</i> , 2015, 96, 163-169.	1.3	17
20	Predictive value of urine interleukin-18 in the evolution and outcome of acute kidney injury in critically ill adult patients. <i>British Journal of Anaesthesia</i> , 2015, 114, 460-468.	1.5	47
21	Plasma hyaluronan and hemorheology in patients with septic shock: A clinical and experimental study. <i>Clinical Hemorheology and Microcirculation</i> , 2014, 56, 133-144.	0.9	15
22	Moderate Intra-Abdominal Hypertension Leads to Anaerobic Metabolism in the Rectus Abdominis Muscle Tissue of Critically Ill Patients: A Prospective Observational Study. <i>BioMed Research International</i> , 2014, 2014, 1-8.	0.9	3
23	Confined ischemia may improve remote myocardial outcome after rat cardiac arrest. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014, 74, 27-36.	0.6	2
24	Medical emergency team activation: performance of conventional dichotomised criteria versus national early warning score. <i>Acta Anaesthesiologica Scandinavica</i> , 2014, 58, 411-419.	0.7	41
25	The Urine Protein NGAL Predicts Renal Replacement Therapy, but Not Acute Kidney Injury or 90-Day Mortality in Critically Ill Adult Patients. <i>Anesthesia and Analgesia</i> , 2014, 119, 95-102.	1.1	21
26	Effect of mattress and bed frame deflection on real chest compression depth measured with two CPR sensors. <i>Resuscitation</i> , 2014, 85, 840-843.	1.3	19
27	Long-term outcomes in patients with severe sepsis randomised to resuscitation with hydroxyethyl starch 130/0.42 or Ringer's acetate. <i>Intensive Care Medicine</i> , 2014, 40, 927-934.	3.9	35
28	The predictive value of soluble urokinase plasminogen activator receptor (SuPAR) regarding 90-day mortality and 12-month neurological outcome in critically ill patients after out-of-hospital cardiac arrest. Data from the prospective FINNRESUSCI study. <i>Resuscitation</i> , 2014, 85, 1562-1567.	1.3	15
29	Therapeutic hypothermia after out-of-hospital cardiac arrest in Finnish intensive care units: the FINNRESUSCI study. <i>Intensive Care Medicine</i> , 2013, 39, 826-837.	3.9	133
30	Incidence, risk factors and 90-day mortality of patients with acute kidney injury in Finnish intensive care units: the FINNAKI study. <i>Intensive Care Medicine</i> , 2013, 39, 420-428.	3.9	348
31	SuPAR and PAI-1 in critically ill, mechanically ventilated patients. <i>Intensive Care Medicine</i> , 2013, 39, 489-496.	3.9	296
32	Real-time audiovisual feedback system in a physician-staffed helicopter emergency medical service in Finland: the quality results and barriers to implementation. <i>Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine</i> , 2013, 21, 50.	1.1	23
33	Validation of Intraluminal and Intraperitoneal microdialysis in ischemic small intestine. <i>BMC Gastroenterology</i> , 2013, 13, 170.	0.8	6
34	Deeper chest compression – More complications for cardiac arrest patients?. <i>Resuscitation</i> , 2013, 84, 760-765.	1.3	164
35	Factors associated with delayed activation of medical emergency team and excess mortality: An Utstein-style analysis. <i>Resuscitation</i> , 2013, 84, 173-178.	1.3	72
36	Hydroxyethyl Starch 130/0.42 versus Ringer's Acetate in Severe Sepsis. <i>New England Journal of Medicine</i> , 2012, 367, 124-134.	13.9	1,594

#	ARTICLE	IF	CITATIONS
37	Therapeutic hypothermia after cardiac arrest â€“ cerebral perfusion and metabolism during upper and lower threshold normocapnia. Resuscitation, 2011, 82, 1174-1179.	1.3	55
38	Comparing the effect of hydroxyethyl starch 130/0.4 with balanced crystalloid solution on mortality and kidney failure in patients with severe sepsis (6S - Scandinavian Starch for Severe Sepsis/Septic) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50 Trials, 2011, 12, 24.	0.7	52
39	Glutamate release predicts ongoing myocardial ischemia of rat hearts. Scandinavian Journal of Clinical and Laboratory Investigation, 2010, 70, 217-224.	0.6	19
40	Acute respiratory failure in intensive care units. FINNALI: a prospective cohort study. Intensive Care Medicine, 2009, 35, 1352-1361.	3.9	112
41	Incidence of iatrogenic dyscarbia during mild therapeutic hypothermia after successful resuscitation from out-of-hospital cardiac arrest. Resuscitation, 2009, 80, 990-993.	1.3	64
42	Hypothermic preconditioning of donor organs prior to harvesting and ischaemia using ice-cold intravenous fluids. Medical Hypotheses, 2009, 73, 65-66.	0.8	9
43	Prehospital induction of therapeutic hypothermia during CPR: A pilot study. Resuscitation, 2008, 76, 360-363.	1.3	88
44	Induction of therapeutic hypothermia during prehospital CPR using ice-cold intravenous fluid. Resuscitation, 2008, 79, 205-211.	1.3	63
45	CAN WE DISTINGUISH BETWEEN DIFFERENT TYPES OF LOCAL PERFUSION/METABOLIC DERANGEMENT BY REGIONAL VENOUS CONCENTRATIONS OF INTERMEDIARY ENERGY SUBSTRATES?. Shock, 2004, 22, 191-192.	1.0	1
46	Jejunal luminal microdialysate lactate in cardiac tamponade â€“ effect of low systemic blood flow on gut mucosa. Intensive Care Medicine, 2002, 28, 953-962.	3.9	15