## Sharon Kean

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10753034/publications.pdf

Version: 2024-02-01

1039406 1372195 2,554 9 9 10 citations h-index g-index papers 10 10 10 3271 citing authors docs citations times ranked all docs

| # | ARTICLE  | IF           | CITATION |
|---|--|--------------|----------|
| 1 | Primary care-led weight management for remission of type 2 diabetes (DiRECT): an open-label, cluster-randomised trial. Lancet, The, 2018, 391, 541-551.  | 6.3          | 1,282    |
| 2 | Durability of a primary care-led weight-management intervention for remission of type 2 diabetes: 2-year results of the DiRECT open-label, cluster-randomised trial. Lancet Diabetes and Endocrinology,the, 2019, 7, 344-355.                                      | 5 <b>.</b> 5 | 569      |
| 3 | Thyroid Hormone Therapy for Older Adults with Subclinical Hypothyroidism. New England Journal of Medicine, 2017, 376, 2534-2544.   | 13.9         | 366      |
| 4 | The Diabetes Remission Clinical Trial (DiRECT): protocol for a cluster randomised trial. BMC Family Practice, 2016, 17, 20.  | 2.9          | 86       |
| 5 | Association Between Levothyroxine Treatment and Thyroid-Related Symptoms Among Adults Aged 80 Years and Older With Subclinical Hypothyroidism. JAMA - Journal of the American Medical Association, 2019, 322, 1977.  | 3.8          | 78       |
| 6 | The effect of a programme to improve men's sedentary time and physical activity: The European Fans in Training (EuroFIT) randomised controlled trial. PLoS Medicine, 2019, 16, e1002736.   | 3.9          | 61       |
| 7 | Clinical and metabolic features of the randomised controlled Diabetes Remission Clinical Trial (DiRECT) cohort. Diabetologia, 2018, 61, 589-598.   | 2.9          | 46       |
| 8 | Study protocol; Thyroid hormone Replacement for Untreated older adults with Subclinical hypothyroidism - a randomised placebo controlled Trial (TRUST). BMC Endocrine Disorders, 2017, 17, 6.  | 0.9          | 36       |
| 9 | Xanthine oxidase inhibition for the improvement of long-term outcomes following ischaemic stroke and transient ischaemic attack (XILO-FIST) – Protocol for a randomised double blind placebo-controlled clinical trial. European Stroke Journal, 2018, 3, 281-290. | 2.7          | 26       |