
David J Keene

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/1076627/publications.pdf Version: 2024-02-01



DAVID I KEENE

#	Article	IF	CITATIONS
1	Changing the System - Major Trauma Patients and Their Outcomes in the NHS (England) 2008–17. EClinicalMedicine, 2018, 2-3, 13-21.	7.1	208
2	Ageing, Muscle Power and Physical Function: A Systematic Review and Implications for Pragmatic Training Interventions. Sports Medicine, 2016, 46, 1311-1332.	6.5	157
3	Prognostic models for identifying risk of poor outcome in people with acute ankle sprains: the SPRAINED development and external validation study. Health Technology Assessment, 2018, 22, 1-112.	2.8	124
4	Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults. JAMA - Journal of the American Medical Association, 2016, 316, 1455.	7.4	94
5	The Ankle Injury Management (AIM) trial: a pragmatic, multicentre, equivalence randomised controlled trial and economic evaluation comparing close contact casting with open surgical reduction and internal fixation in the treatment of unstable ankle fractures in patients aged over 60 years. Health Technology Assessment, 2016, 20, 1-158.	2.8	73
6	Effect of platelet-rich plasma on healing tissues in acute ruptured Achilles tendon: a human immunohistochemistry study. Lancet, The, 2015, 385, S19.	13.7	59
7	Is cervical spine rotation, as used in the standard vertebrobasilar insufficiency test, associated with a measureable change in intracranial vertebral artery blood flow?. Manual Therapy, 2004, 9, 220-227.	1.6	54
8	Progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of patients with rotator cuff disorders (GRASP): a multicentre, pragmatic, 2 × 2 factorial, randomised controlled trial. Lancet, The, 2021, 398, 416-428.	13.7	53
9	Platelet rich plasma injection for acute Achilles tendon rupture: PATH-2 randomised, placebo controlled, superiority trial. BMJ: British Medical Journal, 2019, 367, l6132.	2.3	51
10	Negative-pressure wound therapy versus standard dressings for adults with an open lower limb fracture: the WOLLF RCT. Health Technology Assessment, 2018, 22, 1-162.	2.8	42
11	Early Ankle Movement Versus Immobilization in the Postoperative Management of Ankle Fracture in Adults: A Systematic Review and Meta-analysis. Journal of Orthopaedic and Sports Physical Therapy, 2014, 44, 690-C7.	3.5	41
12	Prognostic factors for recovery following acute lateral ankle ligament sprain: a systematic review. BMC Musculoskeletal Disorders, 2017, 18, 421.	1.9	34
13	Risk of Mortality: The Relationship With Associated Injuries and Fracture Treatment Methods in Patients With Unilateral or Bilateral Femoral Shaft Fractures. Journal of Trauma, 2010, 69, 405-410.	2.3	33
14	Research priorities in fragility fractures of the lower limb and pelvis: a UK priority setting partnership with the James Lind Alliance. BMJ Open, 2018, 8, e023301.	1.9	31
15	Protocol for a randomised controlled trial of standard wound management versus negative pressure wound therapy in the treatment of adult patients with an open fracture of the lower limb: UK Wound management of Open Lower Limb Fractures (UK WOLFF). BMJ Open, 2015, 5, e009087.	1.9	23
16	Measuring physical capacity and performance in older people. Best Practice and Research in Clinical Rheumatology, 2017, 31, 243-254.	3.3	22
17	Clinical and cost-effectiveness of progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of rotator cuff disorders: protocol for a 2x2 factorial randomised controlled trial (the GRASP trial). BMJ Open, 2017, 7, e018004.	1.9	20
18	Three-Year Follow-up of a Trial of Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults. JAMA - Journal of the American Medical Association, 2018, 319, 1274.	7.4	19

David J Keene

#	Article	IF	CITATIONS
19	Intrarater reliability and agreement of linear encoder derived heel-rise endurance test outcome measures in healthy adults. Journal of Electromyography and Kinesiology, 2017, 36, 34-39.	1.7	17
20	Platelet rich Plasma in Achilles Tendon Healing 2 (PATH-2) trial: protocol for a multicentre, participant and assessor-blinded, parallel-group randomised clinical trial comparing platelet-rich plasma (PRP) injection versus placebo injection for Achilles tendon rupture. BMJ Open, 2017, 7, e018135.	1.9	16
21	Chronic pain with neuropathic characteristics after surgery for major trauma to the lower limb: prevalence, predictors, and association with pain severity, disability, and quality of life in the UK WHiST trial. Bone and Joint Journal, 2021, 103-B, 1047-1054.	4.4	16
22	Ankle Injury Management (AIM): design of a pragmatic multi-centre equivalence randomised controlled trial comparing Close Contact Casting (CCC) to Open surgical Reduction and Internal Fixation (ORIF) in the treatment of unstable ankle fractures in patients over 60Âyears. BMC Musculoskeletal Disorders, 2014, 15, 79.	1.9	14
23	The application of multilevel modelling to account for the influence of walking speed in gait analysis. Gait and Posture, 2016, 43, 216-219.	1.4	12
24	Fall prevention interventions in primary care to reduce fractures and falls in people aged 70 years and over: the PreFIT three-arm cluster RCT. Health Technology Assessment, 2021, 25, 1-114.	2.8	12
25	How effective are platelet rich plasma injections in treating musculoskeletal soft tissue injuries?. BMJ, The, 2016, 352, i517.	6.0	11
26	Helicopter and ground emergency medical services transportation to hospital after major trauma in England: a comparative cohort study. Trauma Surgery and Acute Care Open, 2020, 5, e000508.	1.6	11
27	Upper extremity deep vein thrombosis (Paget-Schroetter syndrome) after surfing: A case report. Manual Therapy, 2015, 20, 358-360.	1.6	10
28	Participation in a trial in the emergency situation: a qualitative study of patient experience in the UK WOLLF trial. Trials, 2018, 19, 328.	1.6	10
29	Progressive exercise compared with best-practice advice, with or without corticosteroid injection, for rotator cuff disorders: the GRASP factorial RCT. Health Technology Assessment, 2021, 25, 1-158.	2.8	10
30	Progressive functional exercise versus best practice advice for adults aged 50 years or over after ankle fracture: protocol for a pilot randomised controlled trial in the UK - the Ankle Fracture Treatment: Enhancing Rehabilitation (AFTER) study. BMJ Open, 2019, 9, e030877.	1.9	8
31	Development and implementation of the physiotherapy-led exercise interventions for the treatment of rotator cuff disorders for the †Getting it Right: Addressing Shoulder Pain' (GRASP) trial. Physiotherapy, 2020, 107, 252-266.	0.4	8
32	Characteristics of L-PRP preparations for treating Achilles tendon rupture within the PATH-2 study. Platelets, 2021, 32, 273-279.	2.3	8
33	Platelet-rich plasma injection for adults with acute Achilles tendon rupture: the PATH-2 RCT. Efficacy and Mechanism Evaluation, 2019, 6, 1-98.	0.7	8
34	Implications of the Ankle Injury Management (AIM) trial. Bone and Joint Journal, 2019, 101-B, 1472-1475.	4.4	7
35	Objectively quantified lower limb strength recovery in people treated surgically or non-surgically after patellar dislocation: A systematic review. Physical Therapy in Sport, 2021, 51, 110-138.	1.9	7
36	Factors associated with mobility outcomes in older people post-ankle fracture: An observational cohort study focussing on peripheral vessel function. Injury, 2013, 44, 987-993.	1.7	6

David J Keene

#	Article	IF	CITATIONS
37	Exercise for people with a fragility fracture of the pelvis or lower limb: a systematic review of interventions evaluated in clinical trials and reporting quality. BMC Musculoskeletal Disorders, 2020, 21, 435.	1.9	6
38	Predicting patient-reported and objectively measured functional outcome 6 months after ankle fracture in people aged 60 years or over in the UK: prognostic model development and internal validation. BMJ Open, 2019, 9, e029813.	1.9	6
39	Development and prospective external validation of a tool to predict poor recovery at 9 months after acute ankle sprain in UK emergency departments: the SPRAINED prognostic model. BMJ Open, 2018, 8, e022802.	1.9	5
40	Patient and informal carer experience of hip fracture: a qualitative study using interviews and observation in acute orthopaedic trauma. BMJ Open, 2021, 11, e042040.	1.9	5
41	Functional Outcomes Following Plating or Tension Band Wiring of Olecranon Fractures. European Journal of Trauma and Emergency Surgery, 2003, 29, 273-277.	0.3	4
42	Platelet-rich plasma in Achilles tendon healing 2 (PATH-2) trial: statistical analysis plan for a multicentre, double-blinded, parallel-group, placebo-controlled randomised clinical trial. Trials, 2018, 19, 464.	1.6	4
43	Understanding patient and relative/carer experience of hip fracture in acute care: A qualitative study protocol. International Journal of Orthopaedic and Trauma Nursing, 2017, 25, 36-41.	0.9	3
44	The findings of a surgical hip fracture trial were generalizable to the UK national hip fracture database. Journal of Clinical Epidemiology, 2021, 131, 141-151.	5.0	3
45	Co-producing a multi-stakeholder Core Outcome Set for distal Tibia and Ankle fractures (COSTA): a study protocol. Trials, 2021, 22, 443.	1.6	3
46	Quantitative biomechanical comparison of ankle fracture casting methods. Biomedizinische Technik, 2015, 60, 263-7.	0.8	2
47	The effects of ankle supports on gait in adults: A randomized cross-over study. Journal of Electromyography and Kinesiology, 2015, 25, 973-981.	1.7	2
48	The Immediate Effects of Different Types of Ankle Support Introduced 6 Weeks After Surgical Internal Fixation for Ankle Fracture on Gait and Pain: A Randomized Crossover Trial. Journal of Orthopaedic and Sports Physical Therapy, 2016, 46, 157-167.	3.5	2
49	Progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of rotator cuff disorders: statistical analysis plan for the Getting it Right: Addressing Shoulder Pain (GRASP) 2 × 2 factorial multicentre randomised controlled trial. Trials, 2020. 21. 767.	1.6	2
50	Is an individually tailored programme of intense leg resistance and dynamic exercise acceptable to adults with an acute lateral patellar dislocation? A feasibility study. Pilot and Feasibility Studies, 2021, 7, 197.	1.2	2
51	Flexibility and resistance exercises versus usual care for improving pain and function after distal radius fracture in adults aged 50 years or over: protocol for the WISE randomised multicentre feasibility trial. Pilot and Feasibility Studies, 2022, 8, 55.	1.2	2
52	48â€The Effects Of Different Types Of Ankle Support On Gait: A Randomised Cross-over Study. British Journal of Sports Medicine, 2014, 48, A31-A31.	6.7	0
53	Immobilisation versus early ankle movement for treating acute lateral ankle ligament injuries in adults. The Cochrane Library, 0, , .	2.8	0
54	P135 Lower limb muscle strength and balance in older adults with distal radius fracture: a systematic review. Rheumatology, 2022, 61, .	1.9	0

#	Article	IF	CITATIONS
55	Prolonged ischemia of the ileum and colon after surgical mucosectomy explains contraction and failure of "mucus free―bladder augmentation. Journal of Pediatric Urology, 2022, , .	1.1	0