

David J Keene

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1076627/publications.pdf>

Version: 2024-02-01

55
papers

1,380
citations

471509

17
h-index

361022

35
g-index

58
all docs

58
docs citations

58
times ranked

2020
citing authors

#	ARTICLE	IF	CITATIONS
1	Flexibility and resistance exercises versus usual care for improving pain and function after distal radius fracture in adults aged 50 years or over: protocol for the WISE randomised multicentre feasibility trial. Pilot and Feasibility Studies, 2022, 8, 55.	1.2	2
2	P135 Lower limb muscle strength and balance in older adults with distal radius fracture: a systematic review. Rheumatology, 2022, 61, .	1.9	0
3	Prolonged ischemia of the ileum and colon after surgical mucosectomy explains contraction and failure of mucus free bladder augmentation. Journal of Pediatric Urology, 2022, , .	1.1	0
4	Characteristics of L-PRP preparations for treating Achilles tendon rupture within the PATH-2 study. Platelets, 2021, 32, 273-279.	2.3	8
5	The findings of a surgical hip fracture trial were generalizable to the UK national hip fracture database. Journal of Clinical Epidemiology, 2021, 131, 141-151.	5.0	3
6	Patient and informal carer experience of hip fracture: a qualitative study using interviews and observation in acute orthopaedic trauma. BMJ Open, 2021, 11, e042040.	1.9	5
7	Fall prevention interventions in primary care to reduce fractures and falls in people aged 70 years and over: the PreFIT three-arm cluster RCT. Health Technology Assessment, 2021, 25, 1-114.	2.8	12
8	Chronic pain with neuropathic characteristics after surgery for major trauma to the lower limb: prevalence, predictors, and association with pain severity, disability, and quality of life in the UK WHIST trial. Bone and Joint Journal, 2021, 103-B, 1047-1054.	4.4	16
9	Progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of patients with rotator cuff disorders (GRASP): a multicentre, pragmatic, 2-factorial, randomised controlled trial. Lancet, The, 2021, 398, 416-428.	13.7	53
10	Co-producing a multi-stakeholder Core Outcome Set for distal Tibia and Ankle fractures (COSTA): a study protocol. Trials, 2021, 22, 443.	1.6	3
11	Progressive exercise compared with best-practice advice, with or without corticosteroid injection, for rotator cuff disorders: the GRASP factorial RCT. Health Technology Assessment, 2021, 25, 1-158.	2.8	10
12	Objectively quantified lower limb strength recovery in people treated surgically or non-surgically after patellar dislocation: A systematic review. Physical Therapy in Sport, 2021, 51, 110-138.	1.9	7
13	Is an individually tailored programme of intense leg resistance and dynamic exercise acceptable to adults with an acute lateral patellar dislocation? A feasibility study. Pilot and Feasibility Studies, 2021, 7, 197.	1.2	2
14	Development and implementation of the physiotherapy-led exercise interventions for the treatment of rotator cuff disorders for the "Getting it Right: Addressing Shoulder Pain" (GRASP) trial. Physiotherapy, 2020, 107, 252-266.	0.4	8
15	Helicopter and ground emergency medical services transportation to hospital after major trauma in England: a comparative cohort study. Trauma Surgery and Acute Care Open, 2020, 5, e000508.	1.6	11
16	Progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of rotator cuff disorders: statistical analysis plan for the Getting it Right: Addressing Shoulder Pain (GRASP) 2-factorial multicentre randomised controlled trial. Trials, 2020, 21, 767.	1.6	2
17	Exercise for people with a fragility fracture of the pelvis or lower limb: a systematic review of interventions evaluated in clinical trials and reporting quality. BMC Musculoskeletal Disorders, 2020, 21, 435.	1.9	6
18	Implications of the Ankle Injury Management (AIM) trial. Bone and Joint Journal, 2019, 101-B, 1472-1475.	4.4	7

#	ARTICLE	IF	CITATIONS
19	Platelet rich plasma injection for acute Achilles tendon rupture: PATH-2 randomised, placebo controlled, superiority trial. <i>BMJ: British Medical Journal</i> , 2019, 367, l6132.	2.3	51
20	Progressive functional exercise versus best practice advice for adults aged 50 years or over after ankle fracture: protocol for a pilot randomised controlled trial in the UK - the Ankle Fracture Treatment: Enhancing Rehabilitation (AFTER) study. <i>BMJ Open</i> , 2019, 9, e030877.	1.9	8
21	Predicting patient-reported and objectively measured functional outcome 6 months after ankle fracture in people aged 60 years or over in the UK: prognostic model development and internal validation. <i>BMJ Open</i> , 2019, 9, e029813.	1.9	6
22	Platelet-rich plasma injection for adults with acute Achilles tendon rupture: the PATH-2 RCT. Efficacy and Mechanism Evaluation, 2019, 6, 1-98.	0.7	8
23	Three-Year Follow-up of a Trial of Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults. <i>JAMA - Journal of the American Medical Association</i> , 2018, 319, 1274.	7.4	19
24	Development and prospective external validation of a tool to predict poor recovery at 9 months after acute ankle sprain in UK emergency departments: the SPRAINED prognostic model. <i>BMJ Open</i> , 2018, 8, e022802.	1.9	5
25	Research priorities in fragility fractures of the lower limb and pelvis: a UK priority setting partnership with the James Lind Alliance. <i>BMJ Open</i> , 2018, 8, e023301.	1.9	31
26	Platelet-rich plasma in Achilles tendon healing 2 (PATH-2) trial: statistical analysis plan for a multicentre, double-blinded, parallel-group, placebo-controlled randomised clinical trial. <i>Trials</i> , 2018, 19, 464.	1.6	4
27	Changing the System - Major Trauma Patients and Their Outcomes in the NHS (England) 2008â€“17. <i>EClinicalMedicine</i> , 2018, 2-3, 13-21.	7.1	208
28	Participation in a trial in the emergency situation: a qualitative study of patient experience in the UK WOLLF trial. <i>Trials</i> , 2018, 19, 328.	1.6	10
29	Negative-pressure wound therapy versus standard dressings for adults with an open lower limb fracture: the WOLLF RCT. <i>Health Technology Assessment</i> , 2018, 22, 1-162.	2.8	42
30	Prognostic models for identifying risk of poor outcome in people with acute ankle sprains: the SPRAINED development and external validation study. <i>Health Technology Assessment</i> , 2018, 22, 1-112.	2.8	124
31	Clinical and cost-effectiveness of progressive exercise compared with best practice advice, with or without corticosteroid injection, for the treatment of rotator cuff disorders: protocol for a 2x2 factorial randomised controlled trial (the GRASP trial). <i>BMJ Open</i> , 2017, 7, e018004.	1.9	20
32	Intrarater reliability and agreement of linear encoder derived heel-rise endurance test outcome measures in healthy adults. <i>Journal of Electromyography and Kinesiology</i> , 2017, 36, 34-39.	1.7	17
33	Measuring physical capacity and performance in older people. <i>Best Practice and Research in Clinical Rheumatology</i> , 2017, 31, 243-254.	3.3	22
34	Platelet rich Plasma in Achilles Tendon Healing 2 (PATH-2) trial: protocol for a multicentre, participant and assessor-blinded, parallel-group randomised clinical trial comparing platelet-rich plasma (PRP) injection versus placebo injection for Achilles tendon rupture. <i>BMJ Open</i> , 2017, 7, e018135.	1.9	16
35	Understanding patient and relative/carer experience of hip fracture in acute care: A qualitative study protocol. <i>International Journal of Orthopaedic and Trauma Nursing</i> , 2017, 25, 36-41.	0.9	3
36	Prognostic factors for recovery following acute lateral ankle ligament sprain: a systematic review. <i>BMC Musculoskeletal Disorders</i> , 2017, 18, 421.	1.9	34

#	ARTICLE	IF	CITATIONS
37	Close Contact Casting vs Surgery for Initial Treatment of Unstable Ankle Fractures in Older Adults. JAMA - Journal of the American Medical Association, 2016, 316, 1455.	7.4	94
38	The application of multilevel modelling to account for the influence of walking speed in gait analysis. Gait and Posture, 2016, 43, 216-219.	1.4	12
39	Ageing, Muscle Power and Physical Function: A Systematic Review and Implications for Pragmatic Training Interventions. Sports Medicine, 2016, 46, 1311-1332.	6.5	157
40	How effective are platelet rich plasma injections in treating musculoskeletal soft tissue injuries?. BMJ, The, 2016, 352, i517.	6.0	11
41	The Immediate Effects of Different Types of Ankle Support Introduced 6 Weeks After Surgical Internal Fixation for Ankle Fracture on Gait and Pain: A Randomized Crossover Trial. Journal of Orthopaedic and Sports Physical Therapy, 2016, 46, 157-167.	3.5	2
42	The Ankle Injury Management (AIM) trial: a pragmatic, multicentre, equivalence randomised controlled trial and economic evaluation comparing close contact casting with open surgical reduction and internal fixation in the treatment of unstable ankle fractures in patients aged over 60 years. Health Technology Assessment, 2016, 20, 1-158.	2.8	73
43	Protocol for a randomised controlled trial of standard wound management versus negative pressure wound therapy in the treatment of adult patients with an open fracture of the lower limb: UK Wound management of Open Lower Limb Fractures (UK WOLFF). BMJ Open, 2015, 5, e009087.	1.9	23
44	Quantitative biomechanical comparison of ankle fracture casting methods. Biomedizinische Technik, 2015, 60, 263-7.	0.8	2
45	Effect of platelet-rich plasma on healing tissues in acute ruptured Achilles tendon: a human immunohistochemistry study. Lancet, The, 2015, 385, S19.	13.7	59
46	The effects of ankle supports on gait in adults: A randomized cross-over study. Journal of Electromyography and Kinesiology, 2015, 25, 973-981.	1.7	2
47	Upper extremity deep vein thrombosis (Paget-Schroetter syndrome) after surfing: A case report. Manual Therapy, 2015, 20, 358-360.	1.6	10
48	Ankle Injury Management (AIM): design of a pragmatic multi-centre equivalence randomised controlled trial comparing Close Contact Casting (CCC) to Open surgical Reduction and Internal Fixation (ORIF) in the treatment of unstable ankle fractures in patients over 60 years. BMC Musculoskeletal Disorders, 2014, 15, 79.	1.9	14
49	Early Ankle Movement Versus Immobilization in the Postoperative Management of Ankle Fracture in Adults: A Systematic Review and Meta-analysis. Journal of Orthopaedic and Sports Physical Therapy, 2014, 44, 690-C7.	3.5	41
50	48€...The Effects Of Different Types Of Ankle Support On Gait: A Randomised Cross-over Study. British Journal of Sports Medicine, 2014, 48, A31-A31.	6.7	0
51	Factors associated with mobility outcomes in older people post-ankle fracture: An observational cohort study focussing on peripheral vessel function. Injury, 2013, 44, 987-993.	1.7	6
52	Risk of Mortality: The Relationship With Associated Injuries and Fracture Treatment Methods in Patients With Unilateral or Bilateral Femoral Shaft Fractures. Journal of Trauma, 2010, 69, 405-410.	2.3	33
53	Is cervical spine rotation, as used in the standard vertebrobasilar insufficiency test, associated with a measureable change in intracranial vertebral artery blood flow?. Manual Therapy, 2004, 9, 220-227.	1.6	54
54	Functional Outcomes Following Plating or Tension Band Wiring of Olecranon Fractures. European Journal of Trauma and Emergency Surgery, 2003, 29, 273-277.	0.3	4

#	ARTICLE	IF	CITATIONS
55	Immobilisation versus early ankle movement for treating acute lateral ankle ligament injuries in adults. The Cochrane Library, 0, , .	2.8	0