

# Phillip L Campbell

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10787113/publications.pdf>

Version: 2024-02-01

34  
papers

1,555  
citations

331259

21  
h-index

414034

32  
g-index

36  
all docs

36  
docs citations

36  
times ranked

2093  
citing authors

#	ARTICLE	IF	CITATIONS
1	Inhibition of bromodomain extraterminal histone readers alleviates skin fibrosis in experimental models of scleroderma. <i>JCI Insight</i> , 2022, 7, .	2.3	11
2	Soluble CD13 induces inflammatory arthritis by activating the bradykinin receptor B1. <i>Journal of Clinical Investigation</i> , 2022, 132, .	3.9	6
3	CD6 is a target for cancer immunotherapy. <i>JCI Insight</i> , 2021, 6, .	2.3	18
4	Attenuation of Murine Collagen-Induced Arthritis by Targeting CD6. <i>Arthritis and Rheumatology</i> , 2020, 72, 1505-1513.	2.9	15
5	Real time visualization of cancer cell death, survival and proliferation using fluorochrome-transfected cells in an IncuCyte® imaging system. <i>Journal of Biological Methods</i> , 2020, 7, e133.	1.0	17
6	Angiogenic and Arthritogenic Properties of the Soluble Form of CD13. <i>Journal of Immunology</i> , 2019, 203, 360-369.	0.4	11
7	Citrullinated Inhibitor of DNA Binding 1 Is a Novel Autoantigen in Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2019, 71, 1241-1251.	2.9	4
8	Identification of Pirin as a Molecular Target of the CCG-1423/CCG-203971 Series of Antifibrotic and Antimetastatic Compounds. <i>ACS Pharmacology and Translational Science</i> , 2019, 2, 92-100.	2.5	28
9	5-Aryl-1,3,4-oxadiazol-2-ylthioalkanoic Acids: A Highly Potent New Class of Inhibitors of Rho/Myocardin-Related Transcription Factor (MRTF)/Serum Response Factor (SRF)-Mediated Gene Transcription as Potential Antifibrotic Agents for Scleroderma. <i>Journal of Medicinal Chemistry</i> , 2019, 62, 4350-4369.	2.9	34
10	A unique role for galectin-9 in angiogenesis and inflammatory arthritis. <i>Arthritis Research and Therapy</i> , 2018, 20, 31.	1.6	39
11	Pharmacokinetic optimization of CCG-203971: Novel inhibitors of the Rho/MRTF/SRF transcriptional pathway as potential antifibrotic therapeutics for systemic scleroderma. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2017, 27, 1744-1749.	1.0	42
12	Takinib, a Selective TAK1 Inhibitor, Broadens the Therapeutic Efficacy of TNF- $\alpha$ Inhibition for Cancer and Autoimmune Disease. <i>Cell Chemical Biology</i> , 2017, 24, 1029-1039.e7.	2.5	104
13	Inflammatory properties of inhibitor of DNA binding 1 secreted by synovial fibroblasts in rheumatoid arthritis. <i>Arthritis Research and Therapy</i> , 2016, 18, 87.	1.6	23
14	Scleroderma dermal microvascular endothelial cells exhibit defective response to pro-angiogenic chemokines. <i>Rheumatology</i> , 2016, 55, 745-754.	0.9	24
15	Activation of the Thromboxane A2 Receptor by 8-Isoprostane Inhibits the Pro-Angiogenic Effect of Vascular Endothelial Growth Factor in Scleroderma. <i>Journal of Investigative Dermatology</i> , 2015, 135, 3153-3162.	0.3	23
16	A key role for Fut1-regulated angiogenesis and ICAM-1 expression in K/BxN arthritis. <i>Annals of the Rheumatic Diseases</i> , 2015, 74, 1459-1466.	0.5	16
17	Dynamic Interactions Between Plasma IL-1 Family Cytokines and Central Endogenous Opioid Neurotransmitter Function in Humans. <i>Neuropsychopharmacology</i> , 2015, 40, 554-565.	2.8	23
18	Fucosyltransferase 1 mediates angiogenesis, cell adhesion and rheumatoid arthritis synovial tissue fibroblast proliferation. <i>Arthritis Research and Therapy</i> , 2014, 16, R28.	1.6	30

#	ARTICLE	IF	CITATIONS
19	Fucosyltransferase 1 Mediates Angiogenesis in Rheumatoid Arthritis. <i>Arthritis and Rheumatology</i> , 2014, 66, 2047-2058.	2.9	9
20	Citrullination of Epithelial Neutrophil-Activating Peptide 78/CXCL5 Results in Conversion From a Non-Monocyte-Recruiting Chemokine to a Monocyte-Recruiting Chemokine. <i>Arthritis and Rheumatology</i> , 2014, 66, 2716-2727.	2.9	51
21	A novel role for inducible Fut2 in angiogenesis. <i>Angiogenesis</i> , 2013, 16, 195-205.	3.7	14
22	Association of Plasma Interleukin-18 Levels with Emotion Regulation and $\mu$ -Opioid Neurotransmitter Function in Major Depression and Healthy Volunteers. <i>Biological Psychiatry</i> , 2011, 69, 808-812.	0.7	71
23	Administration of IL-18BP by gene therapy reduces inflammation and prevents joint destruction by downregulation of MMP9 in rat AIA: role of MMP9 in bone and joint destruction in arthritis. <i>Annals of the Rheumatic Diseases</i> , 2011, 70, A71-A72.	0.5	0
24	Green tea extract inhibits chemokine production, but up-regulates chemokine receptor expression, in rheumatoid arthritis synovial fibroblasts and rat adjuvant-induced arthritis. <i>Rheumatology</i> , 2010, 49, 467-479.	0.9	54
25	Suppression of CC chemokine production and HMVEC chemotaxis by green tea extract. <i>FASEB Journal</i> , 2010, 24, 724.2.	0.2	0
26	Monocyte Chemoattractant Protein-1 and Macrophage Inflammatory Protein-1 $\alpha$ as Possible Biomarkers for the Chronic Pelvic Pain Syndrome. <i>Journal of Urology</i> , 2008, 179, 1857-1862.	0.2	106
27	Epigallocatechin-3-gallate inhibits IL-6 synthesis and suppresses transsignaling by enhancing soluble gp130 production. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008, 105, 14692-14697.	3.3	119
28	Interleukin-18 induces angiogenic factors in rheumatoid arthritis synovial tissue fibroblasts via distinct signaling pathways. <i>Arthritis and Rheumatism</i> , 2007, 56, 1787-1797.	6.7	92
29	CXCL16-mediated cell recruitment to rheumatoid arthritis synovial tissue and murine lymph nodes is dependent upon the MAPK pathway. <i>Arthritis and Rheumatism</i> , 2006, 54, 765-778.	6.7	101
30	The Effect of Sulfasalazine on Rheumatoid Arthritic Synovial Tissue Chemokine Production. <i>Experimental and Molecular Pathology</i> , 2002, 73, 84-92.	0.9	37
31	Ley/H: An Endothelial-Selective, Cytokine-Inducible, Angiogenic Mediator. <i>Journal of Immunology</i> , 2000, 164, 4868-4877.	0.4	60
32	Evaluation of the cytokines interleukin 8 and epithelial neutrophil activating peptide 78 as indicators of inflammation in prostatic secretions. <i>Urology</i> , 2000, 56, 1025-1029.	0.5	123
33	IL-1 $\beta$ AND TNF- $\alpha$ IN PROSTATIC SECRETIONS ARE INDICATORS IN THE EVALUATION OF MEN WITH CHRONIC PROSTATITIS. <i>Journal of Urology</i> , 2000, 164, 214-218.	0.2	198
34	Treatment with sulfasalazine or sulfapyridine, but not 5-aminosalicylic acid, inhibits basic fibroblast growth factor-induced endothelial cell chemotaxis. <i>Arthritis and Rheumatism</i> , 1999, 42, 1927-1935.	6.7	49