

# Cihan DÃ¼ndar

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10803822/publications.pdf>

Version: 2024-02-01

8  
papers

176  
citations

1307594

7  
h-index

1720034

7  
g-index

8  
all docs

8  
docs citations

8  
times ranked

335  
citing authors

#	ARTICLE	IF	CITATIONS
1	Early Detection of Bi-ventricular and Atrial Mechanical Dysfunction Using Two-Dimensional Speckle Tracking Echocardiography in Patients with Sarcoidosis. <i>Lung</i> , 2015, 193, 669-675.	3.3	30
2	Two-Dimensional Strain and Strain Rate Imaging of the Left Atrium and Left Ventricle in Adult Patients with Atrial Septal Defects before and after the Later Stage of Percutaneous Device Closure. <i>Echocardiography</i> , 2015, 32, 470-474.	0.9	11
3	Left Ventricular and Atrial Functions in Hypertrophic Cardiomyopathy Patients with Very High <sc>LVOT</sc> Gradient: A Speckle Tracking Echocardiographic Study. <i>Echocardiography</i> , 2014, 31, 833-841.	0.9	27
4	The value of plasma D-dimer level on admission in predicting no-reflow after primary percutaneous coronary intervention and long-term prognosis in patients with acute ST segment elevation myocardial infarction. <i>Journal of Thrombosis and Thrombolysis</i> , 2014, 38, 339-347.	2.1	41
5	Time Course of Right Ventricular Remodeling after Percutaneous Atrial Septal Defect Closure: Assessment of Regional Deformation Properties with Two-Dimensional Strain and Strain Rate Imaging. <i>Echocardiography</i> , 2013, 30, 324-330.	0.9	25
6	Functional mitral regurgitation and papillary muscle dyssynchrony in patients with left ventricular systolic dysfunction. <i>Anatolian Journal of Cardiology</i> , 2011, 11, 450-5.	0.4	0
7	The importance of papillary muscle dyssynchrony in predicting the severity of functional mitral regurgitation in patients with non-ischaemic dilated cardiomyopathy: a two-dimensional speckle-tracking echocardiography study. <i>European Journal of Echocardiography</i> , 2010, 11, 671-676.	2.3	23
8	Predictors and clinical significance of angiographically detected distal embolization after primary percutaneous coronary interventions. <i>Coronary Artery Disease</i> , 2007, 18, 443-449.	0.7	19