Otilia Vieira

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10818609/publications.pdf

Version: 2024-02-01

| 6 papers | 475 citations | 1478505 6 h-index | 1872680 6 g-index |
|-------------|------------------|-------------------------|-------------------------|
| 6 | 6 | 6 | 597 |
| all docs | docs citations | times ranked | citing authors |

| # | Article | IF | CITATIONS |
|---|---|-----|-----------|
| 1 | Oxidized LDLs alter the activity of the ubiquitinâ€proteasome pathway: potential role in oxidized LDLâ€induced apoptosis. FASEB Journal, 2000, 14, 532-542. | 0.5 | 119 |
| 2 | HDL counterbalance the proinflammatory effect of oxidized LDL by inhibiting intracellular reactive oxygen species rise, proteasome activation, and subsequent NF $\hat{a}\in\mathbb{P}B$ activation in smooth muscle cells. FASEB Journal, 2003, 17, 743-745. | 0.5 | 98 |
| 3 | Inhibition of metmyoglobin/H2o2-dependent low density lipoprotein lipid peroxidation by naturally occurring phenolic acids. Biochemical Pharmacology, 1996, 51, 395-402. | 4.4 | 96 |
| 4 | Effect of dietary phenolic compounds on apoptosis of human cultured endothelial cells induced by oxidized LDL. British Journal of Pharmacology, 1998, 123, 565-573. | 5.4 | 70 |
| 5 | Oxidized LDL and 4-hydroxynonenal modulate tyrosine kinase receptor activity. Molecular Aspects of Medicine, 2003, 24, 251-261. | 6.4 | 62 |
| 6 | Phenolic antioxidants trolox and caffeic acid modulate the oxidized LDL-induced EGF-receptor activation. British Journal of Pharmacology, 2001, 132, 1777-1788. | 5.4 | 30 |