

Feng Zhu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10826308/publications.pdf>

Version: 2024-02-01

14
papers

769
citations

759233
12
h-index

1058476
14
g-index

14
all docs

14
docs citations

14
times ranked

1039
citing authors

#	ARTICLE	IF	CITATIONS
1	Hyperparasitoids Use Herbivore-Induced Plant Volatiles to Locate Their Parasitoid Host. PLoS Biology, 2012, 10, e1001435.	5.6	168
2	Foliar-feeding insects acquire microbiomes from the soil rather than the host plant. Nature Communications, 2019, 10, 1254.	12.8	135
3	Insect herbivore-associated organisms affect plant responses to herbivory. New Phytologist, 2014, 204, 315-321.	7.3	78
4	Plant community composition steers grassland vegetation via soil legacy effects. Ecology Letters, 2020, 23, 973-982.	6.4	76
5	Drought stress affects plant metabolites and herbivore preference but not host location by its parasitoids. Oecologia, 2015, 177, 701-713.	2.0	75
6	Symbiotic polydnavirus and venom reveal parasitoid to its hyperparasitoids. Proceedings of the National Academy of Sciences of the United States of America, 2018, 115, 5205-5210.	7.1	54
7	Beyond Plant Microbiome Composition: Exploiting Microbial Functions and Plant Traits via Integrated Approaches. Frontiers in Bioengineering and Biotechnology, 2020, 8, 896.	4.1	44
8	Parasitism overrides herbivore identity allowing hyperparasitoids to locate their parasitoid host using herbivore-induced plant volatiles. Molecular Ecology, 2015, 24, 2886-2899.	3.9	40
9	Parasitic wasp-associated symbiont affects plant-mediated species interactions between herbivores. Ecology Letters, 2018, 21, 957-967.	6.4	34
10	Body Odors of Parasitized Caterpillars Give Away the Presence of Parasitoid Larvae to Their Primary Hyperparasitoid Enemies. Journal of Chemical Ecology, 2014, 40, 986-995.	1.8	22
11	Taking plant-soil feedbacks to the field in a temperate grassland. Basic and Applied Ecology, 2019, 40, 30-42.	2.7	17
12	Soil Microbial Composition and phoD Gene Abundance Are Sensitive to Phosphorus Level in a Long-Term Wheat-Maize Crop System. Frontiers in Microbiology, 2020, 11, 605955.	3.5	17
13	Development of a solitary koinobiont hyperparasitoid in different instars of its primary and secondary hosts. Journal of Insect Physiology, 2016, 90, 36-42.	2.0	5
14	Intrinsic competition between primary hyperparasitoids of the solitary endoparasitoid <i>Microgaster rubecula</i> . Ecological Entomology, 2016, 41, 292-300.	2.2	4