

Tomas Andersson

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10868071/publications.pdf>

Version: 2024-02-01

47
papers

2,948
citations

304743

22
h-index

265206

42
g-index

47
all docs

47
docs citations

47
times ranked

4909
citing authors

#	ARTICLE	IF	CITATIONS
1	Calculating measures of biological interaction. <i>European Journal of Epidemiology</i> , 2005, 20, 575-579.	5.7	1,111
2	The incidence of SUDEP. <i>Neurology</i> , 2017, 89, 170-177.	1.1	209
3	Clinical risk factors in SUDEP. <i>Neurology</i> , 2020, 94, e419-e429.	1.1	197
4	Plasma Levels of Tissue Plasminogen Activator/Plasminogen Activator Inhibitor-1 Complex and von Willebrand Factor Are Significant Risk Markers for Recurrent Myocardial Infarction in the Stockholm Heart Epidemiology Program (SHEEP) Study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2000, 20, 2019-2023.	2.4	178
5	Hospitalization for psychiatric disorders before and after onset of unprovoked seizures/epilepsy. <i>Neurology</i> , 2012, 78, 396-401.	1.1	168
6	Work Stress and Low Sense of Coherence Is Associated With Type 2 Diabetes in Middle-Aged Swedish Women. <i>Diabetes Care</i> , 2003, 26, 719-724.	8.6	156
7	Explanations of Socioeconomic Differences in Excess Risk of Type 2 Diabetes in Swedish Men and Women. <i>Diabetes Care</i> , 2004, 27, 716-721.	8.6	91
8	Diabetes Prevalence in Sweden at Present and Projections for Year 2050. <i>PLoS ONE</i> , 2015, 10, e0143084.	2.5	73
9	Newly diagnosed single unprovoked seizures and epilepsy in Stockholm, Sweden: First report from the Stockholm Incidence Registry of Epilepsy (SIRE). <i>Epilepsia</i> , 2009, 50, 1094-1101.	5.1	59
10	Unprovoked seizures after traumatic brain injury: A population-based case-control study. <i>Epilepsia</i> , 2015, 56, 1438-1444.	5.1	54
11	Declining incidence trends for hip fractures have not been accompanied by improvements in lifetime risk or post-fracture survival – A nationwide study of the Swedish population 60years and older. <i>Bone</i> , 2015, 78, 55-61.	2.9	52
12	Smoking Is Associated With Reduced Risk of Autoimmune Diabetes in Adults Contrasting With Increased Risk in Overweight Men With Type 2 Diabetes. <i>Diabetes Care</i> , 2013, 36, 604-610.	8.6	51
13	Pharmacologic treatment and SUDEP risk. <i>Neurology</i> , 2020, 95, e2509-e2518.	1.1	48
14	Risk for injuries and accidents in epilepsy. <i>Neurology</i> , 2018, 90, e779-e789.	1.1	44
15	Circumstances of SUDEP: A nationwide population-based case series. <i>Epilepsia</i> , 2018, 59, 1074-1082.	5.1	42
16	Age-Specific Trends in Morbidity, Mortality and Case-Fatality from Cardiovascular Disease, Myocardial Infarction and Stroke in Advanced Age: Evaluation in the Swedish Population. <i>PLoS ONE</i> , 2013, 8, e64928.	2.5	35
17	The incidence of unprovoked seizures and occurrence of neurodevelopmental comorbidities in children at the time of their first epileptic seizure and during the subsequent six months. <i>Epilepsy Research</i> , 2015, 113, 140-150.	1.6	29
18	Incidence and prevalence of type 2 diabetes by occupation: results from all Swedish employees. <i>Diabetologia</i> , 2020, 63, 95-103.	6.3	29

#	ARTICLE	IF	CITATIONS
19	Evolution over time of SUDEP incidence: A nationwide population-based cohort study. <i>Epilepsia</i> , 2018, 59, e120-e124.	5.1	28
20	Smoking and the Risk of LADA: Results From a Swedish Population-Based Case-Control Study. <i>Diabetes Care</i> , 2016, 39, 794-800.	8.6	26
21	Prevalence and Incidence of Diabetes in Stockholm County 1990-2010. <i>PLoS ONE</i> , 2014, 9, e104033.	2.5	26
22	Prior hospitalization for stroke, diabetes, myocardial infarction, and subsequent risk of unprovoked seizures. <i>Epilepsia</i> , 2011, 52, 301-307.	5.1	25
23	Occupational exposure to particles and increased risk of developing chronic obstructive pulmonary disease (COPD): A population-based cohort study in Stockholm, Sweden. <i>Environmental Research</i> , 2021, 200, 111739.	7.5	24
24	Interaction Between Overweight and Genotypes of HLA, TCF7L2, and FTO in Relation to the Risk of Latent Autoimmune Diabetes in Adults and Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 4815-4826.	3.6	22
25	Trends in age at first hospital admission in relation to trends in life expectancy in Swedish men and women above the age of 60. <i>BMJ Open</i> , 2013, 3, e003447.	1.9	20
26	Alcohol and the risk for latent autoimmune diabetes in adults: results based on Swedish ESTRID study. <i>European Journal of Endocrinology</i> , 2014, 171, 535-543.	3.7	17
27	Shift and night work during pregnancy and preterm birth—a cohort study of Swedish health care employees. <i>International Journal of Epidemiology</i> , 2022, 50, 1864-1874.	1.9	17
28	Disability pensions related to heavy physical workload: a cohort study of middle-aged and older workers in Sweden. <i>International Archives of Occupational and Environmental Health</i> , 2021, 94, 1851-1861.	2.3	14
29	Cancer incidence in Stockholm firefighters 1958–2012: an updated cohort study. <i>International Archives of Occupational and Environmental Health</i> , 2018, 91, 285-291.	2.3	13
30	Physical Activity, Genetic Susceptibility, and the Risk of Latent Autoimmune Diabetes in Adults and Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020, 105, e4112-e4123.	3.6	11
31	Does Improved Survival Lead to a More Fragile Population: Time Trends in Second and Third Hospital Admissions among Men and Women above the Age of 60 in Sweden. <i>PLoS ONE</i> , 2014, 9, e99034.	2.5	11
32	Unprovoked seizures in multiple sclerosis and systemic lupus erythematosus: A population-based case-control study. <i>Epilepsy Research</i> , 2012, 101, 284-287.	1.6	10
33	Consumption of red meat, genetic susceptibility, and risk of LADA and type 2 diabetes. <i>European Journal of Nutrition</i> , 2021, 60, 769-779.	3.9	9
34	The magnitude of bias in a cross-sectional study on lifestyle factors in relation to Type 2 diabetes. <i>Scandinavian Journal of Public Health</i> , 2006, 34, 665-668.	2.3	8
35	Tobacco and type 2 diabetes: is the association explained by genetic factors?. <i>International Journal of Epidemiology</i> , 2019, 48, 926-933.	1.9	6
36	Genotypes of HLA, TCF7L2, and FTO as potential modifiers of the association between sweetened beverage consumption and risk of LADA and type 2 diabetes. <i>European Journal of Nutrition</i> , 2020, 59, 127-135.	3.9	6

