

Mohamed El-Sayed

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10872496/publications.pdf>

Version: 2024-02-01

16
papers

301
citations

1039406

9
h-index

1058022

14
g-index

16
all docs

16
docs citations

16
times ranked

352
citing authors

#	ARTICLE	IF	CITATIONS
1	Subacromial injection of autologous platelet-rich plasma versus corticosteroid for the treatment of symptomatic partial rotator cuff tears. <i>European Journal of Orthopaedic Surgery and Traumatology</i> , 2016, 26, 837-842.	0.6	126
2	Solvatochromism, Crystallochromism, and Solid State Structures of Hydrophilically Functionalized Aromatic Amino Ketones Containing Furan and Thiophene Rings. <i>Chemistry of Materials</i> , 2003, 15, 746-754.	3.2	41
3	Linear solvation energy (LSE) correlations of the solvatochromic response and x-ray structure analysis of hydrophilically N-substituted Michler's ketone derivatives. <i>Journal of Physical Organic Chemistry</i> , 2001, 14, 247-255.	0.9	18
4	Ilizarov external fixation for management of severe relapsed clubfeet in older children. <i>Foot and Ankle Surgery</i> , 2013, 19, 177-181.	0.8	18
5	Management of simple (types A and B) closed tibial shaft fractures using percutaneous lag-screw fixation and Ilizarov external fixation in adults. <i>International Orthopaedics</i> , 2012, 36, 2133-2138.	0.9	17
6	Solid-state Structures of N-Substituted Michler's Ketones and Their Relation to Solvatochromism. <i>European Journal of Organic Chemistry</i> , 2002, 2002, 4159-4168.	1.2	16
7	The Solvent-Like Nature of Silica Particles in Organic Solvents. <i>Monatshefte für Chemie</i> , 2001, 132, 1347-1361.	0.9	13
8	UV/Vis Spectroscopic Properties of N-(2-Hydroxy-4-N,N-dimethyl-aminobenzylidene)-4-nitroaniline in Various Solvents and Solid Environments. <i>Monatshefte für Chemie</i> , 2003, 134, 361-370.	0.9	12
9	Linear Solvation Energy Correlation of the Solvatochromic Response of Amino Acid Functionalized Chromophores. <i>Journal of Physical Chemistry B</i> , 2007, 111, 7224-7233.	1.2	10
10	Internal Polarity of Class I and Class II Type Sol-Gel Hybrid Materials Using Aromatic Aminoketones as Solvatochromic Probes for Adsorbed Solvents and the Silicatic Cage. <i>Journal of Sol-Gel Science and Technology</i> , 2005, 34, 77-94.	1.1	8
11	Synthesis, solvatochromism, and photophysical properties of the polymer-tetherable 3-[4-di(2-hydroxyethyl)amino]phenyl-(2-furyl)-2-propene-1-one. <i>Journal of Photochemistry and Photobiology A: Chemistry</i> , 2007, 185, 44-50.	2.0	7
12	Structure and solvatochromism of heteroaromatic aminoketones containing thiophene moieties. <i>Journal of Physical Organic Chemistry</i> , 2005, 18, 1086-1098.	0.9	5
13	Synthesis, properties, and solvatochromism of 1,3-dimethyl-5-[(thien-2-yl)-[4-(1-piperidyl)phenyl]methylidene]-(1H,3H)-pyrimidine-2,4,6-trione. <i>Journal of Physical Organic Chemistry</i> , 2007, 20, 264-270.	0.9	5
14	Is the Denis Browne Splint a Myth? A Long-Term Prospective Cohort Study in Clubfoot Management using Denis Browne Splint Versus Daily Exercise Protocol. <i>Journal of Foot and Ankle Surgery</i> , 2020, 59, 314-322.	0.5	5
15	A modification of the Ponseti method for clubfoot management: a prospective comparative study. <i>Journal of Children's Orthopaedics</i> , 2021, 15, 1-10.	0.4	0
16	The Solvent-Like Nature of Silica Particles in Organic Solvents. , 2002, , 111-125.		0