
## Mohamed El-Sayed

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10872496/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	A modification of the Ponseti method for clubfoot management: a prospective comparative study. Journal of Children's Orthopaedics, 2021, 15, 1-10.	0.4	О
2	ls the Denis Browne Splint a Myth? A Long-Term Prospective Cohort Study in Clubfoot Management using Denis Browne Splint Versus Daily Exercise Protocol. Journal of Foot and Ankle Surgery, 2020, 59, 314-322.	0.5	5
3	Subacromial injection of autologous platelet-rich plasma versus corticosteroid for the treatment of symptomatic partial rotator cuff tears. European Journal of Orthopaedic Surgery and Traumatology, 2016, 26, 837-842.	0.6	126
4	llizarov external fixation for management of severe relapsed clubfeet in older children. Foot and Ankle Surgery, 2013, 19, 177-181.	0.8	18
5	Management of simple (types A and B) closed tibial shaft fractures using percutaneous lag-screw fixation and Ilizarov external fixation in adults. International Orthopaedics, 2012, 36, 2133-2138.	0.9	17
6	Linear Solvation Energy Correlation of the Solvatochromic Response of Amino Acid Functionalized Chromophores. Journal of Physical Chemistry B, 2007, 111, 7224-7233.	1.2	10
7	Synthesis, solvatochromism, and photophysical properties of the polymer-tetherable 3-[4-di(2-hydroxyethyl)amino]phenyl-l-(2-furyl)-2-propene-l-one. Journal of Photochemistry and Photobiology A: Chemistry, 2007, 185, 44-50.	2.0	7
8	Synthesis, properties, and solvatochromism of 1,3-dimethyl-5-{(thien-2-yl)-[4-(1-piperidyl) phenyl]methylidene}-(1H,3H)-pyrimidine-2,4,6-trione. Journal of Physical Organic Chemistry, 2007, 20, 264-270.	0.9	5
9	Internal Polarity of Class I and Class II Type Sol–Gel Hybrid Materials Using Aromatic Aminoketones as Solvatochromic Probes for Adsorbed Solvents and the Silicatic Cage. Journal of Sol-Gel Science and Technology, 2005, 34, 77-94.	1.1	8
10	Structure and solvatochromism of heteroaromatic aminoketones containing thiophene moieties. Journal of Physical Organic Chemistry, 2005, 18, 1086-1098.	0.9	5
11	UV/Vis Spectroscopic Properties of N -(2?-Hydroxy-4?- N , N -dimethyl-aminobenzylidene)-4-nitroaniline in Various Solvents and Solid Environments. Monatshefte Für Chemie, 2003, 134, 361-370.	0.9	12
12	Solvatochromism, Crystallochromism, and Solid State Structures of Hydrophilically Functionalized Aromatic Amino Ketones Containing Furan and Thiophene Rings. Chemistry of Materials, 2003, 15, 746-754.	3.2	41
13	Solid-state Structures of N-Substituted Michler's Ketones and Their Relation to Solvatochromism. European Journal of Organic Chemistry, 2002, 2002, 4159-4168.	1.2	16
14	The Solvent-Like Nature of Silica Particles in Organic Solvents. , 2002, , 111-125.		0
15	The Solvent-Like Nature of Silica Particles in Organic Solvents. Monatshefte Für Chemie, 2001, 132, 1347-1361.	0.9	13
16	Linear solvation energy (LSE) correlations of the solvatochromic response and x-ray structure analysis of hydrophilicallyN-substituted Michler's ketone derivatives. Journal of Physical Organic Chemistry, 2001, 14, 247-255.	0.9	18