

Anders M Abildgaard

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1087344/publications.pdf>

Version: 2024-02-01

17
papers

559
citations

840728

11
h-index

996954

15
g-index

18
all docs

18
docs citations

18
times ranked

1035
citing authors

#	ARTICLE	IF	CITATIONS
1	Probiotic treatment reduces depressive-like behaviour in rats independently of diet. <i>Psychoneuroendocrinology</i> , 2017, 79, 40-48.	2.7	149
2	A high-fat diet exacerbates depressive-like behavior in the Flinders Sensitive Line (FSL) rat, a genetic model of depression. <i>Psychoneuroendocrinology</i> , 2011, 36, 623-633.	2.7	77
3	The microbial metabolite indole-3-propionic acid improves glucose metabolism in rats, but does not affect behaviour. <i>Archives of Physiology and Biochemistry</i> , 2018, 124, 306-312.	2.1	67
4	Severe Acute Respiratory Syndrome Coronavirus 2 Seroprevalence Survey Among 17 971 Healthcare and Administrative Personnel at Hospitals, Prehospital Services, and Specialist Practitioners in the Central Denmark Region. <i>Clinical Infectious Diseases</i> , 2021, 73, e2853-e2860.	5.8	60
5	Altered fecal microbiota composition in the Flinders sensitive line rat model of depression. <i>Psychopharmacology</i> , 2019, 236, 1445-1457.	3.1	44
6	Probiotic treatment protects against the pro-depressant-like effect of high-fat diet in Flinders Sensitive Line rats. <i>Brain, Behavior, and Immunity</i> , 2017, 65, 33-42.	4.1	39
7	GLP-1 receptor agonists have a sustained stimulatory effect on corticosterone release after chronic treatment. <i>Acta Neuropsychiatrica</i> , 2015, 27, 25-32.	2.1	23
8	Chronic high-fat diet increases acute neuroendocrine stress response independently of prenatal dexamethasone treatment in male rats. <i>Acta Neuropsychiatrica</i> , 2014, 26, 8-18.	2.1	22
9	The antidepressant-like effect of probiotics and their faecal abundance may be modulated by the cohabiting gut microbiota in rats. <i>European Neuropsychopharmacology</i> , 2019, 29, 98-110.	0.7	22
10	A diet-induced gut microbiota component and related plasma metabolites are associated with depressive-like behaviour in rats. <i>European Neuropsychopharmacology</i> , 2021, 43, 10-21.	0.7	16
11	Dosage of Anticoagulants in Obesity: Recommendations Based on a Systematic Review. <i>Seminars in Thrombosis and Hemostasis</i> , 2020, 46, 932-969.	2.7	16
12	Reference intervals for plasma vitamin B12 and plasma/serum methylmalonic acid in Danish children, adults and elderly. <i>Clinica Chimica Acta</i> , 2022, 525, 62-68.	1.1	10
13	Lactase persistence genotyping on whole blood by loop-mediated isothermal amplification and melting curve analysis. <i>Clinica Chimica Acta</i> , 2018, 482, 50-56.	1.1	5
14	Lot variation and inter-device differences contribute to poor analytical performance of the DCA Vantage [®] , _c HbA _{1c} POCT instrument in a true clinical setting. <i>Clinical Chemistry and Laboratory Medicine</i> , 2022, 60, 127-134.	2.3	5
15	High-sensitivity troponin T and I in patients suspected of acute myocardial infarction. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2022, 82, 96-103.	1.2	3
16	Circulating trefoil factors in relation to lung cancer, age and lung function: a cross-sectional study in patients referred for suspected lung cancer. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2021, 81, 446-450.	1.2	1
17	The activated partial thromboplastin time may not reveal even severe fibrinogen deficiency. <i>Clinical Chemistry and Laboratory Medicine</i> , 2021, 59, e297-e300.	2.3	0