

Kieran Smith

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1089014/publications.pdf>

Version: 2024-02-01

16
papers

74
citations

1684188

5
h-index

1588992

8
g-index

18
all docs

18
docs citations

18
times ranked

91
citing authors

#	ARTICLE	IF	CITATIONS
1	An acute dose of inorganic dietary nitrate does not improve high-intensity, intermittent exercise performance in temperate or hot and humid conditions. <i>European Journal of Applied Physiology</i> , 2019, 119, 723-733.	2.5	16
2	The Clinical Application of Mealtime Whey Protein for the Treatment of Postprandial Hyperglycaemia for People With Type 2 Diabetes: A Long Whey to Go. <i>Frontiers in Nutrition</i> , 2020, 7, 587843.	3.7	12
3	Postexercise Glycemic Control in Type 1 Diabetes Is Associated With Residual β -Cell Function. <i>Diabetes Care</i> , 2020, 43, 2362-2370.	8.6	11
4	Thrice daily consumption of a novel, premeal shot containing a low dose of whey protein increases time in euglycemia during 7 days of free-living in individuals with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2022, 10, e002820.	2.8	7
5	The Postprandial Glycaemic and Hormonal Responses Following the Ingestion of a Novel, Ready-to-Drink Shot Containing a Low Dose of Whey Protein in Centrally Obese and Lean Adult Males: A Randomised Controlled Trial. <i>Frontiers in Endocrinology</i> , 2021, 12, 696977.	3.5	6
6	Bone turnover and metabolite responses to exercise in people with and without long-duration type 1 diabetes: a case-control study. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001779.	2.8	5
7	Type 1 diabetes patients increase CXCR4+ and CXCR7+ haematopoietic and endothelial progenitor cells with exercise, but the response is attenuated. <i>Scientific Reports</i> , 2021, 11, 14502.	3.3	5
8	Capturing the real-world benefit of residual β -cell function during clinically important time periods in established Type 1 diabetes. <i>Diabetic Medicine</i> , 2022, 39, e14814.	2.3	5
9	294-OR: Type 1 Diabetes Patients with Residual Beta-Cell Function Display Improved Time in Euglycemia and Less Glycaemic Fluctuation after Exercise. <i>Diabetes</i> , 2019, 68, 294-OR.	0.6	2
10	The Effects of a High-Protein Diet on Markers of Muscle Damage Following Exercise in Active Older Adults: A Randomized, Controlled Trial. <i>International Journal of Sport Nutrition and Exercise Metabolism</i> , 2020, 30, 323-329.	2.1	2
11	Type 1 Diabetes Patients With Different Residual Beta-Cell Function but Similar Age, HBA1c, and Cardiorespiratory Fitness Have Differing Exercise-Induced Angiogenic Cell Mobilisation. <i>Frontiers in Endocrinology</i> , 2022, 13, 797438.	3.5	2
12	Identifying Behavioural Determinants to Uptake and Adherence to a Whey Protein Supplement for the Management of Type 2 Diabetes: A Qualitative Interview Study. <i>Nutrients</i> , 2022, 14, 565.	4.1	1
13	2276-PUB: A Palatable, Novel Whey Protein Shot Attenuates Postprandial Glycemia in Lean and Centrally Obese Adult Males. <i>Diabetes</i> , 2019, 68, .	0.6	0
14	680-P: Residual β -Cell Function in Long-Term Type 1 Diabetes Is Associated with a Fivefold Greater Mobilization of Endothelial Progenitor Cells after Exercise. <i>Diabetes</i> , 2020, 69, .	0.6	0
15	679-P: Endothelial and Hematopoietic Progenitor Cells Are Reduced at Rest and Have an Attenuated Mobilization Response to Exercise in People with Type 1 Diabetes. <i>Diabetes</i> , 2020, 69, .	0.6	0
16	2169-PUB: Premeal Consumption of a Small, Ready-to-Drink Whey Protein Shot Suppresses Postprandial Hyperglycemia in Adults with Type 2 Diabetes. <i>Diabetes</i> , 2020, 69, 2169-PUB.	0.6	0