

Peter H Selle

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/10923697/peter-h-selle-publications-by-year.pdf>

Version: 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

64 papers	1,864 citations	19 h-index	42 g-index
66 ext. papers	2,228 ext. citations	3.4 avg, IF	5.22 L-index

#	Paper	IF	Citations
64	Graded methionine dietary inclusions influence growth performance and apparent ileal amino acid digestibility coefficients and disappearance rates in broiler chickens.. <i>Animal Nutrition</i> , 2022 , 8, 160-168	4.8	1
63	Assessment of limiting dietary amino acids in broiler chickens offered reduced crude protein diets. <i>Animal Nutrition</i> , 2022 ,	4.8	1
62	A multivariate Box-Behnken assessment of elevated branched-chain amino acid concentrations in reduced crude protein diets offered to male broiler chickens.. <i>PLoS ONE</i> , 2022 , 17, e0266080	3.7	
61	An evaluation of elevated branched-chain amino acid inclusions on the performance of broiler chickens offered reduced-crude protein, wheat-based diets from 7 to 28 days post-hatch. <i>Animal Feed Science and Technology</i> , 2022 , 286, 115255	3	0
60	Evaluation of dietary crude protein concentrations, fishmeal, and sorghum inclusions in broiler chickens offered wheat-based diet via Box-Behnken response surface design. <i>PLoS ONE</i> , 2021 , 16, e0260285	3.7	25
59	Progress towards reduced-crude protein diets for broiler chickens and sustainable chicken-meat production. <i>Journal of Animal Science and Biotechnology</i> , 2021 , 12, 20	6	8
58	Maize-based diets are more conducive to crude protein reductions than wheat-based diets for broiler chickens. <i>Animal Feed Science and Technology</i> , 2021 , 275, 114867	3	9
57	Starch digestion rates in multiple samples of commonly used feed grains in diets for broiler chickens. <i>Animal Nutrition</i> , 2021 , 7, 450-459	4.8	2
56	The Dynamic Conversion of Dietary Protein and Amino Acids into Chicken-Meat Protein. <i>Animals</i> , 2021 , 11,	3.1	3
55	Implications of excreta uric acid concentrations in broilers offered reduced-crude protein diets and dietary glycine requirements for uric acid synthesis. <i>Animal Nutrition</i> , 2021 , 7, 939-946	4.8	1
54	Addressing the shortfalls of sorghum as a feed grain for chicken-meat production. <i>World's Poultry Science Journal</i> , 2021 , 77, 29-41	3	1
53	Starch digestibility and energy utilisation of maize- and wheat-based diets is superior to sorghum-based diets in broiler chickens offered diets supplemented with phytase and xylanase. <i>Animal Feed Science and Technology</i> , 2020 , 264, 114475	3	8
52	Composition and characterisation of kafirin, the dominant protein fraction in grain sorghum. <i>Animal Production Science</i> , 2020 , 60, 1163	1.4	4
51	Capping dietary starch:protein ratios in moderately reduced crude protein, wheat-based diets showed promise but further reductions generated inferior growth performance in broiler chickens. <i>Animal Nutrition</i> , 2020 , 6, 168-178	4.8	13
50	Synthetic and Crystalline Amino Acids: Alternatives to Soybean Meal in Chicken-Meat Production. <i>Animals</i> , 2020 , 10,	3.1	25
49	Potential of CIELAB colour scores to gauge the quality of sorghum as a feed grain for chicken-meat production. <i>Animal Production Science</i> , 2020 , 60, 930	1.4	1
48	Effects of reducing dietary crude protein and whole grain feeding on performance and amino acid metabolism in broiler chickens offered wheat-based diets. <i>Animal Feed Science and Technology</i> , 2020 , 260, 114386	3	6

47	Glycine equivalent and threonine inclusions in reduced-crude protein, maize-based diets impact on growth performance, fat deposition, starch-protein digestive dynamics and amino acid metabolism in broiler chickens. <i>Animal Feed Science and Technology</i> , 2020 , 261, 114387	3	10
46	Effects of reduced crude protein levels, dietary electrolyte balance, and energy density on the performance of broiler chickens offered maize-based diets with evaluations of starch, protein, and amino acid metabolism. <i>Poultry Science</i> , 2020 , 99, 1421-1431	3.9	23
45	Facilitating the acceptance of tangibly reduced-crude protein diets for chicken-meat production. <i>Animal Nutrition</i> , 2020 , 6, 247-257	4.8	11
44	Protein digestive dynamics of meat and bone meals in broiler chickens. <i>Animal Nutrition</i> , 2020 , 6, 521-528	4.8	1
43	Impacts of reduced-crude protein diets on key parameters in male broiler chickens offered maize-based diets. <i>Poultry Science</i> , 2020 , 99, 505-516	3.9	29
42	Extending daily feed access intervals does not influence lysine HCl utilization but enhances amino acid digestibilities in broiler chickens. <i>Poultry Science</i> , 2019 , 98, 4801-4814	3.9	2
41	Box-Behnken optimisation of growth performance, plasma metabolites and carcass traits as influenced by dietary energy, amino acid and starch to lipid ratios in broiler chickens. <i>PLoS ONE</i> , 2019 , 14, e0213875	3.7	6
40	Influence of starch sources and dietary protein levels on intestinal functionality and intestinal mucosal amino acids catabolism in broiler chickens. <i>Journal of Animal Science and Biotechnology</i> , 2019 , 10, 26	6	7
39	The influence of phytase, pre-pellet cracked maize and dietary crude protein level on broiler performance via response surface methodology. <i>Journal of Animal Science and Biotechnology</i> , 2019 , 10, 80	6	1
38	The influence of exogenous phytase on the post-enteral availability of amino acids in broiler chickens offered wheat-based diets. <i>Animal Feed Science and Technology</i> , 2019 , 258, 114300	3	2
37	Feeding whole grain and phytase to meat chickens: recent Australian experience. <i>Animal Production Science</i> , 2019 , 59, 2010	1.4	1
36	Sorghum as a Feed Grain for Animal Production 2019 , 355-391		4
35	Dietary starch influences growth performance, nutrient utilisation and digestive dynamics of protein and amino acids in broiler chickens offered low-protein diets. <i>Animal Feed Science and Technology</i> , 2018 , 237, 55-67	3	26
34	Responses in digestibilities of macro-minerals, trace minerals and amino acids generated by exogenous phytase and xylanase in canola meal diets offered to broiler chickens. <i>Animal Feed Science and Technology</i> , 2018 , 240, 22-30	3	9
33	Outlook: Sorghum as a feed grain for Australian chicken-meat production. <i>Animal Nutrition</i> , 2018 , 4, 17-30	4.8	19
32	Dietary starch to lipid ratios influence growth performance, nutrient utilisation and carcass traits in broiler chickens offered diets with different energy densities. <i>PLoS ONE</i> , 2018 , 13, e0205272	3.7	8
31	Progress in comprehending the phytate-phytase axis in chicken-meat production. <i>Animal Production Science</i> , 2018 , 58, 1767	1.4	2
30	The interactions of exogenous phytase with whole grain feeding and effects of barley as the whole grain component in broiler diets based on wheat, sorghum and wheat-sorghum blends. <i>Animal Feed Science and Technology</i> , 2017 , 227, 1-12	3	13

29	Comparative performance of broiler chickens offered nutritionally equivalent diets based on six diverse, Bannin-free sorghum varieties with quantified concentrations of phenolic compounds, kafirin, and phytate. <i>Animal Production Science</i> , 2017 , 57, 828	1.4	12
28	The potential of rapid visco-analysis starch pasting profiles to gauge the quality of sorghum as a feed grain for chicken-meat production. <i>Animal Nutrition</i> , 2017 , 3, 11-18	4.8	7
27	Fishmeal and maize starch inclusions in sorghum-soybean meal diets generate different responses in growth performance, nutrient utilisation, starch and protein digestive dynamics of broiler chickens. <i>Animal Feed Science and Technology</i> , 2017 , 227, 32-41	3	13
26	Pre- and post-pellet whole grain inclusions enhance feed conversion efficiency, energy utilisation and gut integrity in broiler chickens offered wheat-based diets. <i>Animal Feed Science and Technology</i> , 2017 , 224, 115-123	3	19
25	Effects of phytase inclusions in diets containing ground wheat or 12.5% whole wheat (pre- and post-pellet) and phytase and protease additions, individually and in combination, to diets containing 12.5% pre-pellet whole wheat on the performance of broiler chickens. <i>Animal Feed Science and Technology</i> , 2017 , 234, 139-150	3	5
24	The influence of the selection of macronutrients coupled with dietary energy density on the performance of broiler chickens. <i>PLoS ONE</i> , 2017 , 12, e0185480	3.7	3
23	Evaluation of ground grain versus pre- and post-pellet whole grain additions to poultry diets via a response surface design. <i>British Poultry Science</i> , 2017 , 58, 718-728	1.9	6
22	Growth performance, nutrient utilisation and carcass composition respond to dietary protein concentrations in broiler chickens but responses are modified by dietary lipid levels. <i>British Journal of Nutrition</i> , 2017 , 118, 250-262	3.6	9
21	Rapid protein disappearance rates along the small intestine advantage poultry performance and influence the post-enteral availability of amino acids. <i>British Journal of Nutrition</i> , 2017 , 118, 1031-1042	3.6	14
20	Phytase inclusions of 500 and 2000 FTU/kg in maize-based broiler diets impact on growth performance, nutrient utilisation, digestive dynamics of starch, protein (N), sodium and IP6 phytate degradation in the gizzard and four small intestinal segments. <i>Animal Feed Science and Technology</i> , 2017 , 223, 13-22	3	14
19	An assessment of the influence of macronutrients on growth performance and nutrient utilisation in broiler chickens by nutritional geometry. <i>British Journal of Nutrition</i> , 2016 , 116, 2129-2138	3.6	12
18	The influence of meat-and-bone meal and exogenous phytase on growth performance, bone mineralisation and digestibility coefficients of protein (N), amino acids and starch in broiler chickens. <i>Animal Nutrition</i> , 2016 , 2, 86-92	4.8	5
17	The multidimensional causal factors of 'wet litter' in chicken-meat production. <i>Science of the Total Environment</i> , 2016 , 562, 766-776	10.2	60
16	Starch utilisation in chicken-meat production: the foremost influential factors. <i>Animal Production Science</i> , 2016 , 56, 797	1.4	23
15	Reducing agent and exogenous protease additions, individually and in combination, to wheat- and sorghum-based diets interactively influence parameters of nutrient utilisation and digestive dynamics in broiler chickens. <i>Animal Nutrition</i> , 2016 , 2, 303-311	4.8	21
14	Sodium metabisulphite enhances energy utilisation in broiler chickens offered sorghum-based diets with five different grain varieties. <i>Animal Feed Science and Technology</i> , 2016 , 219, 159-174	3	10
13	Standard phytase inclusion in maize-based broiler diets enhances digestibility coefficients of starch, amino acids and sodium in four small intestinal segments and digestive dynamics of starch and protein. <i>Animal Feed Science and Technology</i> , 2015 , 209, 240-248	3	38
12	Concentrations of specific phenolic compounds in six red sorghums influence nutrient utilisation in broiler chickens. <i>Animal Feed Science and Technology</i> , 2015 , 210, 190-199	3	39

11	Performance of broiler chickens offered nutritionally-equivalent diets based on two red grain sorghums with quantified kafirin concentrations as intact pellets or re-ground mash following steam-pelleting at 65 or 97°C conditioning temperatures. <i>Animal Nutrition</i> , 2015 , 1, 220-228	4.8	17
10	Grain Sorghum: A Conundrum for Chicken-Meat Production. <i>Agriculture (Switzerland)</i> , 2015 , 5, 1224-1251	5	26
9	Whole-grain feeding for chicken-meat production: possible mechanisms driving enhanced energy utilisation and feed conversion. <i>Animal Production Science</i> , 2015 , 55, 559	1.4	31
8	Dual effects of sodium phytate on the structural stability and solubility of proteins. <i>Journal of Agricultural and Food Chemistry</i> , 2013 , 61, 290-5	5.7	30
7	Strategies to enhance the performance of pigs and poultry on sorghum-based diets. <i>Animal Feed Science and Technology</i> , 2013 , 181, 1-14	3	37
6	Protein-phytate interactions in pig and poultry nutrition: a reappraisal. <i>Nutrition Research Reviews</i> , 2012 , 25, 1-17	7	149
5	Consequences of calcium interactions with phytate and phytase for poultry and pigs. <i>Livestock Science</i> , 2009 , 124, 126-141	1.7	262
4	Phytate-degrading enzymes in pig nutrition. <i>Livestock Science</i> , 2008 , 113, 99-122	1.7	200
3	Microbial phytase in poultry nutrition. <i>Animal Feed Science and Technology</i> , 2007 , 135, 1-41	3	470
2	Phytase Supplementation of Wheat-Based Broiler Diets Reduces Dependence on Meat-and-Bone Meal. <i>Journal of Poultry Science</i> , 2006 , 43, 330-338	1.6	9
1	Influence of Dietary Phytate and Exogenous Phytase on Amino Acid Digestibility in Poultry: A Review. <i>Journal of Poultry Science</i> , 2006 , 43, 89-103	1.6	66