

Sergio Romagnani

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/10942937/sergio-romagnani-publications-by-year.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

188
papers

24,198
citations

76
h-index

155
g-index

192
ext. papers

26,302
ext. citations

7.1
avg, IF

7.03
L-index

#	Paper	IF	Citations
188	Impaired immune cell cytotoxicity in severe COVID-19 is IL-6 dependent. <i>Journal of Clinical Investigation</i> , 2020 , 130, 4694-4703	15.9	261
187	Cell-mediated and humoral adaptive immune responses to SARS-CoV-2 are lower in asymptomatic than symptomatic COVID-19 patients. <i>European Journal of Immunology</i> , 2020 , 50, 2013-2024	6.1	35
186	Eomes controls the development of Th17-derived (non-classic) Th1 cells during chronic inflammation. <i>European Journal of Immunology</i> , 2019 , 49, 79-95	6.1	34
185	The Kinetics of Antidrug Antibodies, Drug Levels, and Clinical Outcomes in Infliximab-Exposed Patients with Immune-Mediated Disorders. <i>Journal of Allergy and Clinical Immunology: in Practice</i> , 2018 , 6, 2065-2072.e2	5.4	11
184	Therapeutic Efficacy of Autologous Non-Mobilized Enriched Circulating Endothelial Progenitors in Patients With Critical Limb Ischemia - The SCELTA Trial. <i>Circulation Journal</i> , 2018 , 82, 1688-1698	2.9	18
183	Musculin inhibits human T-helper 17 cell response to interleukin 2 by controlling STAT5B activity. <i>European Journal of Immunology</i> , 2017 , 47, 1427-1442	6.1	13
182	IL-10-Producing Infliximab-Specific T Cells Regulate the Antidrug T Cell Response in Exposed Patients. <i>Journal of Immunology</i> , 2017 , 199, 1283-1289	5.3	10
181	Human circulating group 2 innate lymphoid cells can express CD154 and promote IgE production. <i>Journal of Allergy and Clinical Immunology</i> , 2017 , 139, 964-976.e4	11.5	61
180	How pregnancy can affect autoimmune diseases progression?. <i>Clinical and Molecular Allergy</i> , 2016 , 14, 11	3.7	56
179	Th1 Cells 2016 , 287-293		1
178	IL411: Key immunoregulator at a crossroads of divergent T-cell functions. <i>European Journal of Immunology</i> , 2016 , 46, 2302-2305	6.1	10
177	Interleukin-17-producing decidual CD4+ T cells are not deleterious for human pregnancy when they also produce interleukin-4. <i>Clinical and Molecular Allergy</i> , 2016 , 14, 1	3.7	22
176	Chitinase 3-like-1 is produced by human Th17 cells and correlates with the level of inflammation in juvenile idiopathic arthritis patients. <i>Clinical and Molecular Allergy</i> , 2016 , 14, 16	3.7	8
175	Dermatophagoides pteronyssinus group 2 allergen bound to 8-OH modified adenine reduces the Th2-mediated airway inflammation without inducing a Th17 response and autoimmunity. <i>Molecular Immunology</i> , 2016 , 77, 60-70	4.3	2
174	Demethylation of the RORC2 and IL17A in human CD4+ T lymphocytes defines Th17 origin of nonclassic Th1 cells. <i>Journal of Immunology</i> , 2015 , 194, 3116-26	5.3	54
173	Treatment with 8-OH-modified adenine (TLR7 ligand)-allergen conjugates decreases T helper type 2-oriented murine airway inflammation. <i>Immunology</i> , 2015 , 145, 570-82	7.8	6
172	The 3 major types of innate and adaptive cell-mediated effector immunity. <i>Journal of Allergy and Clinical Immunology</i> , 2015 , 135, 626-35	11.5	378

171	T helper cell mediated-tolerance towards fetal allograft in successful pregnancy. <i>Clinical and Molecular Allergy</i> , 2015 , 13, 9	3.7	38
170	Human Th1 dichotomy: origin, phenotype and biologic activities. <i>Immunology</i> , 2014 , 144, 343	7.8	30
169	Brief report: etanercept inhibits the tumor necrosis factor α -driven shift of Th17 lymphocytes toward a nonclassic Th1 phenotype in juvenile idiopathic arthritis. <i>Arthritis and Rheumatology</i> , 2014 , 66, 1372-7	9.5	38
168	T cell subpopulations. <i>Chemical Immunology and Allergy</i> , 2014 , 100, 155-64		20
167	IL-4-induced gene 1 maintains high Tob1 expression that contributes to TCR unresponsiveness in human T helper 17 cells. <i>European Journal of Immunology</i> , 2014 , 44, 654-61	6.1	25
166	Th17 and non-classic Th1 cells in chronic inflammatory disorders: two sides of the same coin. <i>International Archives of Allergy and Immunology</i> , 2014 , 164, 171-7	3.7	61
165	Reasons for rarity of Th17 cells in inflammatory sites of human disorders. <i>Seminars in Immunology</i> , 2013 , 25, 299-304	10.7	21
164	A novel allergen-adjuvant conjugate suitable for specific immunotherapy of respiratory allergy. <i>Journal of Allergy and Clinical Immunology</i> , 2013 , 132, 84-92	11.5	12
163	Main features of human T helper 17 cells. <i>Annals of the New York Academy of Sciences</i> , 2013 , 1284, 66-706.5		34
162	HLA-G5 induces IL-4 secretion critical for successful pregnancy through differential expression of ILT2 receptor on decidual CD4+ T cells and macrophages. <i>Journal of Immunology</i> , 2013 , 191, 3651-62	5.3	32
161	Distinctive features of classic and nonclassic (Th17 derived) human Th1 cells. <i>European Journal of Immunology</i> , 2012 , 42, 3180-8	6.1	87
160	Rarity of human T helper 17 cells is due to retinoic acid orphan receptor-dependent mechanisms that limit their expansion. <i>Immunity</i> , 2012 , 36, 201-14	32.3	93
159	Defining the human T helper 17 cell phenotype. <i>Trends in Immunology</i> , 2012 , 33, 505-12	14.4	115
158	Drug-specific Th2 cells and IgE antibodies in a patient with anaphylaxis to rituximab. <i>International Archives of Allergy and Immunology</i> , 2012 , 159, 321-6	3.7	45
157	Human TH17 Cells 2011 , 231-242		
156	Mouse T helper 17 phenotype: not so different than in man after all. <i>Cytokine</i> , 2011 , 56, 112-5	4	13
155	Frequency of regulatory T cells in peripheral blood and in tumour-infiltrating lymphocytes correlates with poor prognosis in renal cell carcinoma. <i>BJU International</i> , 2011 , 107, 1500-6	5.6	81
154	Evidence of the transient nature of the Th17 phenotype of CD4+CD161+ T cells in the synovial fluid of patients with juvenile idiopathic arthritis. <i>Arthritis and Rheumatism</i> , 2011 , 63, 2504-15		160

153	The TLR7 ligand 9-benzyl-2-butoxy-8-hydroxy adenine inhibits IL-17 response by eliciting IL-10 and IL-10-inducing cytokines. <i>Journal of Immunology</i> , 2011 , 186, 4707-15	5.3	31
152	Identification of a novel subset of human circulating memory CD4(+) T cells that produce both IL-17A and IL-4. <i>Journal of Allergy and Clinical Immunology</i> , 2010 , 125, 222-30.e1-4	11.5	228
151	Human and murine Th17. <i>Current Opinion in HIV and AIDS</i> , 2010 , 5, 114-9	4.2	26
150	Evidence for a cross-talk between human neutrophils and Th17 cells. <i>Blood</i> , 2010 , 115, 335-43	2.2	520
149	CD161 is a marker of all human IL-17-producing T-cell subsets and is induced by RORC. <i>European Journal of Immunology</i> , 2010 , 40, 2174-81	6.1	266
148	The transient nature of the Th17 phenotype. <i>European Journal of Immunology</i> , 2010 , 40, 3312-6	6.1	55
147	Modified adenine (9-benzyl-2-butoxy-8-hydroxyadenine) redirects Th2-mediated murine lung inflammation by triggering TLR7. <i>Journal of Immunology</i> , 2009 , 182, 880-9	5.3	22
146	Regeneration of glomerular podocytes by human renal progenitors. <i>Journal of the American Society of Nephrology: JASN</i> , 2009 , 20, 322-32	12.7	407
145	TGF-beta indirectly favors the development of human Th17 cells by inhibiting Th1 cells. <i>European Journal of Immunology</i> , 2009 , 39, 207-15	6.1	118
144	Human Th17 cells: are they different from murine Th17 cells?. <i>European Journal of Immunology</i> , 2009 , 39, 637-40	6.1	47
143	Type 17 T helper cells-origins, features and possible roles in rheumatic disease. <i>Nature Reviews Rheumatology</i> , 2009 , 5, 325-31	8.1	169
142	Properties and origin of human Th17 cells. <i>Molecular Immunology</i> , 2009 , 47, 3-7	4.3	137
141	Heterogeneity of human effector CD4+ T cells. <i>Arthritis Research and Therapy</i> , 2009 , 11, 257	5.7	120
140	Do studies in humans better depict Th17 cells?. <i>Blood</i> , 2009 , 114, 2213-9	2.2	71
139	Toll-like receptors 3 and 4 are expressed by human bone marrow-derived mesenchymal stem cells and can inhibit their T-cell modulatory activity by impairing Notch signaling. <i>Stem Cells</i> , 2008 , 26, 279-89	5.8	380
138	Human immature myeloid dendritic cells trigger a TH2-polarizing program via Jagged-1/Notch interaction. <i>Journal of Allergy and Clinical Immunology</i> , 2008 , 121, 1000-5.e8	11.5	61
137	Human Th17 cells. <i>Arthritis Research and Therapy</i> , 2008 , 10, 206	5.7	125
136	Activation of p38(MAPK) mediates the angiostatic effect of the chemokine receptor CXCR3-B. <i>International Journal of Biochemistry and Cell Biology</i> , 2008 , 40, 1764-74	5.6	53

135	The phenotype of human Th17 cells and their precursors, the cytokines that mediate their differentiation and the role of Th17 cells in inflammation. <i>International Immunology</i> , 2008 , 20, 1361-8	4.9	152
134	Functional deficit of T regulatory cells in Fulani, an ethnic group with low susceptibility to Plasmodium falciparum malaria. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2008 , 105, 646-51	11.5	105
133	Human interleukin 17-producing cells originate from a CD161+CD4+ T cell precursor. <i>Journal of Experimental Medicine</i> , 2008 , 205, 1903-16	16.6	569
132	Essential but differential role for CXCR4 and CXCR7 in the therapeutic homing of human renal progenitor cells. <i>Journal of Experimental Medicine</i> , 2008 , 205, 479-90	16.6	230
131	T cells specific for Candida albicans antigens and producing type 2 cytokines in lesional mucosa of untreated HIV-infected patients with pseudomembranous oropharyngeal candidiasis. <i>Microbes and Infection</i> , 2008 , 10, 166-74	9.3	9
130	Role of chemokines in endocrine autoimmune diseases. <i>Endocrine Reviews</i> , 2007 , 28, 492-520	27.2	190
129	Demonstration of circulating allergen-specific CD4+CD25highFoxp3+ T-regulatory cells in both nonatopic and atopic individuals. <i>Journal of Allergy and Clinical Immunology</i> , 2007 , 120, 429-36	11.5	61
128	Eosinophilia in cholesterol atheroembolic disease. <i>Journal of Allergy and Clinical Immunology</i> , 2007 , 120, 1470-1; author reply 1471	11.5	6
127	Th1/Th2 Cells. <i>Inflammatory Bowel Diseases</i> , 2007 , 5, 285-294	4.5	48
126	Specific recruitment of CD4+CD25++ regulatory T cells into the allograft in heart transplant recipients. <i>American Journal of Physiology - Heart and Circulatory Physiology</i> , 2007 , 292, H2425-31	5.2	12
125	Regenerative potential of embryonic renal multipotent progenitors in acute renal failure. <i>Journal of the American Society of Nephrology: JASN</i> , 2007 , 18, 3128-38	12.7	172
124	PF-4/CXCL4 and CXCL4L1 exhibit distinct subcellular localization and a differentially regulated mechanism of secretion. <i>Blood</i> , 2007 , 109, 4127-34	2.2	54
123	Phenotypic and functional features of human Th17 cells. <i>Journal of Experimental Medicine</i> , 2007 , 204, 1849-61	16.6	1476
122	Immune regulation by mesenchymal stem cells derived from adult spleen and thymus. <i>Stem Cells and Development</i> , 2007 , 16, 797-810	4.4	100
121	Immunological tolerance and autoimmunity. <i>Internal and Emergency Medicine</i> , 2006 , 1, 187-96	3.7	38
120	Peripheral blood as a source of stem cells for regenerative medicine. <i>Expert Opinion on Biological Therapy</i> , 2006 , 6, 193-202	5.4	13
119	Isolation and characterization of multipotent progenitor cells from the Bowman's capsule of adult human kidneys. <i>Journal of the American Society of Nephrology: JASN</i> , 2006 , 17, 2443-56	12.7	556
118	Redirection of allergen-specific TH2 responses by a modified adenine through Toll-like receptor 7 interaction and IL-12/IFN release. <i>Journal of Allergy and Clinical Immunology</i> , 2006 , 118, 511-7	11.5	43

117	Role for interferon-gamma in the immunomodulatory activity of human bone marrow mesenchymal stem cells. <i>Stem Cells</i> , 2006 , 24, 386-98	5.8	1030
116	Chemokine receptors in tissue cells and angiogenesis 2006 , 183-203		1
115	CXCR3-mediated opposite effects of CXCL10 and CXCL4 on TH1 or TH2 cytokine production. <i>Journal of Allergy and Clinical Immunology</i> , 2005 , 116, 1372-9	11.5	86
114	CD14+CD34 ^{low} cells with stem cell phenotypic and functional features are the major source of circulating endothelial progenitors. <i>Circulation Research</i> , 2005 , 97, 314-22	15.7	218
113	Thymic regulatory T cells. <i>Autoimmunity Reviews</i> , 2005 , 4, 579-86	13.6	134
112	Functional features of human CD25 ⁺ regulatory thymocytes. <i>Microbes and Infection</i> , 2005 , 7, 1017-22	9.3	10
111	Human 60-kDa heat shock protein is a target autoantigen of T cells derived from atherosclerotic plaques. <i>Journal of Immunology</i> , 2005 , 174, 6509-17	5.3	106
110	A Case of Chronic Urticaria Due to <i>Dirofilaria</i> Infestation. <i>Allergology International</i> , 2004 , 53, 383-385	4.4	
109	The increased prevalence of allergy and the hygiene hypothesis: missing immune deviation, reduced immune suppression, or both?. <i>Immunology</i> , 2004 , 112, 352-63	7.8	328
108	Immunologic influences on allergy and the TH1/TH2 balance. <i>Journal of Allergy and Clinical Immunology</i> , 2004 , 113, 395-400	11.5	329
107	CXC chemokines: the regulatory link between inflammation and angiogenesis. <i>Trends in Immunology</i> , 2004 , 25, 201-9	14.4	327
106	Th2 cells are less susceptible than Th1 cells to the suppressive activity of CD25 ⁺ regulatory thymocytes because of their responsiveness to different cytokines. <i>Blood</i> , 2004 , 103, 3117-21	2.2	149
105	An alternatively spliced variant of CXCR3 mediates the inhibition of endothelial cell growth induced by IP-10, Mig, and I-TAC, and acts as functional receptor for platelet factor 4. <i>Journal of Experimental Medicine</i> , 2003 , 197, 1537-49	16.6	560
104	Human CD8 ⁺ CD25 ⁺ thymocytes share phenotypic and functional features with CD4 ⁺ CD25 ⁺ regulatory thymocytes. <i>Blood</i> , 2003 , 102, 4107-14	2.2	297
103	The novel synthetic immune response modifier R-848 (Resiquimod) shifts human allergen-specific CD4 ⁺ TH2 lymphocytes into IFN-gamma-producing cells. <i>Journal of Allergy and Clinical Immunology</i> , 2003 , 111, 380-8	11.5	84
102	T helper type 1 lymphocytes drive inflammation in human atherosclerotic lesions. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 6658-63	11.5	125
101	New therapeutic strategies in allergic diseases. <i>Drugs of Today</i> , 2003 , 39, 849-65		6
100	Th2-oriented profile of male offspring T cells present in women with systemic sclerosis and reactive with maternal major histocompatibility complex antigens. <i>Arthritis and Rheumatism</i> , 2002 , 46, 445-50		105

99	Role of low nuclear grading of renal carcinoma cells in the functional profile of tumor-infiltrating T cells. <i>International Journal of Cancer</i> , 2002 , 98, 674-81	7.5	10
98	Phenotype, localization, and mechanism of suppression of CD4(+)CD25(+) human thymocytes. <i>Journal of Experimental Medicine</i> , 2002 , 196, 379-87	16.6	333
97	Type 1 T helper cells specific for <i>Candida albicans</i> antigens in peripheral blood and vaginal mucosa of women with recurrent vaginal candidiasis. <i>Journal of Infectious Diseases</i> , 2002 , 186, 87-93	7	19
96	Interferon-inducible protein 10, monokine induced by interferon gamma, and interferon-inducible T-cell alpha chemoattractant are produced by thymic epithelial cells and attract T-cell receptor (TCR) alphabeta+ CD8+ single-positive T cells, TCRgammadelta+ T cells, and natural killer-type cells. <i>Journal of Experimental Medicine</i> , 2001 , 197, 121-7	2.2	98
95	Production of IL-4 and leukemia inhibitory factor by T cells of the cumulus oophorus: a favorable microenvironment for pre-implantation embryo development. <i>European Journal of Immunology</i> , 2001 , 31, 2431-7	6.1	57
94	Defective production of LIF, M-CSF and Th2-type cytokines by T cells at fetomaternal interface is associated with pregnancy loss. <i>Journal of Reproductive Immunology</i> , 2001 , 52, 35-43	4.2	101
93	Chemoattractant receptors expressed on type 2 T cells and their role in disease. <i>International Archives of Allergy and Immunology</i> , 2001 , 125, 273-9	3.7	31
92	Expression of the chemokine receptor CCR3 on human mast cells. <i>International Archives of Allergy and Immunology</i> , 2001 , 124, 146-50	3.7	64
91	Polarization of PPD-specific T-cell response of patients with tuberculosis from Th0 to Th1 profile after successful antimycobacterial therapy or in vitro conditioning with interferon-alpha or interleukin-12. <i>American Journal of Respiratory Cell and Molecular Biology</i> , 2001 , 24, 187-94	5.7	52
90	Reversal of human allergen-specific CRTH2+ T(H)2 cells by IL-12 or the PS-DSP30 oligodeoxynucleotide. <i>Journal of Allergy and Clinical Immunology</i> , 2001 , 108, 815-21	11.5	38
89	H(+),K(+)-atpase (proton pump) is the target autoantigen of Th1-type cytotoxic T cells in autoimmune gastritis. <i>Gastroenterology</i> , 2001 , 120, 377-86	13.3	115
88	Chemokines and lymphopoiesis in human thymus. <i>Trends in Immunology</i> , 2001 , 22, 277-81	14.4	57
87	Cell cycle-dependent expression of CXC chemokine receptor 3 by endothelial cells mediates angiostatic activity. <i>Journal of Clinical Investigation</i> , 2001 , 107, 53-63	15.9	300
86	T-cell responses in allergy and asthma. <i>Current Opinion in Allergy and Clinical Immunology</i> , 2001 , 1, 73-8	3.3	49
85	CRTH2: marker for the detection of human Th2 and Tc2 cells. <i>Advances in Experimental Medicine and Biology</i> , 2001 , 495, 25-9	3.6	9
84	Environmental factors favoring the allergen-specific Th2 response in allergic subjects. <i>Annals of the New York Academy of Sciences</i> , 2000 , 917, 844-52	6.5	11
83	Macrophage-derived chemokine production by activated human T cells in vitro and in vivo: preferential association with the production of type 2 cytokines. <i>European Journal of Immunology</i> , 2000 , 30, 204-10	6.1	97
82	CRTH2 is the most reliable marker for the detection of circulating human type 2 Th and type 2 T cytotoxic cells in health and disease. <i>European Journal of Immunology</i> , 2000 , 30, 2972-9	6.1	244

81	Cytokines and chemokines in T lymphopoiesis and T-cell effector function. <i>Trends in Immunology</i> , 2000 , 21, 416-8		16
80	Limited expression of R5-tropic HIV-1 in CCR5-positive type 1 β polarized T cells explained by their ability to produce RANTES, MIP-1 β and MIP-1 α . <i>Blood</i> , 2000 , 95, 1167-1174	2.2	46
79	Enhanced expression of the CXCR4 co-receptor in HIV-1-infected individuals correlates with the emergence of syncytia-inducing strains. <i>Cytokines, Cellular & Molecular Therapy</i> , 2000 , 6, 19-24		7
78	Macrophage-derived chemokine and EB1-ligand chemokine attract human thymocytes in different stage of development and are produced by distinct subsets of medullary epithelial cells: possible implications for negative selection. <i>Journal of Immunology</i> , 2000 , 165, 238-46	5.3	60
77	The role of lymphocytes in allergic disease. <i>Journal of Allergy and Clinical Immunology</i> , 2000 , 105, 399-408	1.5	360
76	T-cell subsets (Th1 versus Th2). <i>Annals of Allergy, Asthma and Immunology</i> , 2000 , 85, 9-18; quiz 18, 21	3.2	543
75	Chemokine Receptors on Human Mast Cells 2000 , 579-596		1
74	Macrophage-derived chemokine production by activated human T cells in vitro and in vivo: preferential association with the production of type 2 cytokines 2000 , 30, 204		9
73	Macrophage-derived chemokine production by activated human T cells in vitro and in vivo: preferential association with the production of type 2 cytokines 2000 , 30, 204		1
72	Macrophage-Derived Chemokine Is Localized to Thymic Medullary Epithelial Cells and Is a Chemoattractant for CD3+, CD4+, CD8low Thymocytes. <i>Blood</i> , 1999 , 94, 1890-1898	2.2	94
71	Chemokine receptors and other surface molecules preferentially associated with human Th1 or Th2 cells. <i>Microbes and Infection</i> , 1999 , 1, 103-6	9.3	17
70	Relaxin favors the development of activated human T cells into Th1-like effectors. <i>European Journal of Immunology</i> , 1999 , 29, 2241-7	6.1	56
69	Impaired T-cell regulation of B-cell growth in Helicobacter pylori--related gastric low-grade MALT lymphoma. <i>Gastroenterology</i> , 1999 , 117, 1105-12	13.3	91
68	Tryptase-chymase double-positive human mast cells express the eotaxin receptor CCR3 and are attracted by CCR3-binding chemokines. <i>American Journal of Pathology</i> , 1999 , 155, 1195-204	5.8	206
67	Assessment of chemokine receptor expression by human Th1 and Th2 cells in vitro and in vivo. <i>Journal of Leukocyte Biology</i> , 1999 , 65, 691-9	6.5	145
66	Th1/Th2 cells. <i>Inflammatory Bowel Diseases</i> , 1999 , 5, 285-94	4.5	270
65	Defective production of both leukemia inhibitory factor and type 2 T-helper cytokines by decidual T cells in unexplained recurrent abortions. <i>Nature Medicine</i> , 1998 , 4, 1020-4	50.5	551
64	Influence of both TCR repertoire and severity of the atopic status on the cytokine secretion profile of Parietaria officinalis-specific T cells. <i>European Journal of Immunology</i> , 1998 , 28, 37-46	6.1	13

63	Enhanced HIV expression during Th2-oriented responses explained by the opposite regulatory effect of IL-4 and IFN-gamma of fusin/CXCR4. <i>European Journal of Immunology</i> , 1998 , 28, 3280-90	6.1	65
62	High CD30 Ligand Expression by Epithelial Cells and Hassalß Corpuscles in the Medulla of Human Thymus. <i>Blood</i> , 1998 , 91, 3323-3332	2.2	68
61	High CD30 Ligand Expression by Epithelial Cells and Hassalß Corpuscles in the Medulla of Human Thymus. <i>Blood</i> , 1998 , 91, 3323-3332	2.2	7
60	Functional Characterization and Modulation of Cytokine Production by CD8+ T Cells from Human Immunodeficiency Virus-Infected Individuals. <i>Blood</i> , 1997 , 89, 3672-3681	2.2	41
59	The Th1/Th2 paradigm. <i>Trends in Immunology</i> , 1997 , 18, 263-6		812
58	Different cytokine profile and antigen-specificity repertoire in Helicobacter pylori-specific T cell clones from the antrum of chronic gastritis patients with or without peptic ulcer. <i>European Journal of Immunology</i> , 1997 , 27, 1751-5	6.1	184
57	Opposite role for interleukin-4 and interferon-gamma on CD30 and lymphocyte activation gene-3 (LAG-3) expression by activated naive T cells. <i>European Journal of Immunology</i> , 1997 , 27, 2239-44	6.1	57
56	Functional Characterization and Modulation of Cytokine Production by CD8+ T Cells from Human Immunodeficiency Virus-Infected Individuals. <i>Blood</i> , 1997 , 89, 3672-3681	2.2	10
55	Pathophysiology of Th1 and Th2 Responses in Humans 1997 , 149-166		
54	Expression and release of LAG-3-encoded protein by human CD4+ T cells are associated with IFN-gamma production. <i>FASEB Journal</i> , 1996 , 10, 769-76	0.9	78
53	Editorial The role of T-cell derived cytokines in allergic diseases. <i>Expert Opinion on Therapeutic Patents</i> , 1996 , 6, 223-228	6.8	
52	Th1 and Th2 in human diseases. <i>Clinical Immunology and Immunopathology</i> , 1996 , 80, 225-35		299
51	Regulation of fetal allograft survival by a hormone-controlled Th1- and Th2-type cytokines. <i>Immunologic Research</i> , 1996 , 15, 141-50	4.3	119
50	Genetic variability of the human CD4 V2 domain. <i>Immunogenetics</i> , 1996 , 44, 70-2	3.2	7
49	CD30 cell expression and abnormal soluble CD30 serum accumulation in Omennß syndrome: evidence for a T helper 2-mediated condition. <i>European Journal of Immunology</i> , 1996 , 26, 329-34	6.1	101
48	Effects of interferon-alpha on cytokine profile, T cell receptor repertoire and peptide reactivity of human allergen-specific T cells. <i>European Journal of Immunology</i> , 1996 , 26, 697-703	6.1	103
47	Development in vitro of human CD4+ thymocytes into functionally mature Th2 cells. Exogenous interleukin-12 is required for priming thymocytes to produce both Th1 cytokines and interleukin-10. <i>European Journal of Immunology</i> , 1996 , 26, 1083-7	6.1	29
46	Abnormal production of T helper 2 cytokines interleukin-4 and interleukin-5 by T cells from newborns with atopic parents. <i>European Journal of Immunology</i> , 1996 , 26, 2293-8	6.1	67

45	Role for CD30 in HIV expression. <i>Immunology Letters</i> , 1996 , 51, 83-8	4.1	14
44	The pathology of OmennB syndrome. <i>American Journal of Surgical Pathology</i> , 1996 , 20, 773-4	6.7	8
43	Human T-Cell Responses to Grass Pollen Allergens 1996 , 164-175		1
42	Genetic variability of the human CD4 V2 domain. <i>Immunogenetics</i> , 1996 , 44, 70-72	3.2	
41	Biology of human TH1 and TH2 cells. <i>Journal of Clinical Immunology</i> , 1995 , 15, 121-9	5.7	329
40	CD30, Th2 cytokines and HIV infection: a complex and fascinating link. <i>Trends in Immunology</i> , 1995 , 16, 76-80		133
39	Activation of HIV expression by CD30 triggering in CD4+ T cells from HIV-infected individuals. <i>Immunity</i> , 1995 , 3, 251-5	32.3	50
38	T Helper Cell Subsets. <i>BioDrugs</i> , 1995 , 4, 83-88		1
37	CD30 and type 2 T helper (Th2) responses. <i>Journal of Leukocyte Biology</i> , 1995 , 57, 726-30	6.5	113
36	Polyinosinic acid: polycytidylic acid promotes T helper type 1-specific immune responses by stimulating macrophage production of interferon-alpha and interleukin-12. <i>European Journal of Immunology</i> , 1995 , 25, 2656-60	6.1	120
35	CD30 ligation induces nuclear factor-kappa B activation in human T cell lines. <i>European Journal of Immunology</i> , 1995 , 25, 2870-6	6.1	53
34	Role for CD30 antigen in human T helper 2-type responses. <i>Novartis Foundation Symposium</i> , 1995 , 195, 55-62; discussion 62-7		1
33	Human Th1 and Th2 cells: functional properties, regulation of development and role in autoimmunity. <i>Autoimmunity</i> , 1994 , 18, 301-8	3	68
32	An alternative view of the Th1/Th2 switch hypothesis in HIV infection. <i>AIDS Research and Human Retroviruses</i> , 1994 , 10, iii-ix	1.6	45
31	Regulation of the development of type 2 T-helper cells in allergy. <i>Current Opinion in Immunology</i> , 1994 , 6, 838-46	7.8	216
30	Th1 versus Th2 responses in AIDS. <i>Current Opinion in Immunology</i> , 1994 , 6, 616-22	7.8	49
29	A toxic shock syndrome toxin-1 peptide that shows homology to amino acids 180-193 of mycobacterial heat shock protein 65 is presented as conventional antigen. <i>Immunological Investigations</i> , 1994 , 23, 381-91	2.9	7
28	The role of TH1 and TH2 subsets in human infectious diseases. <i>Trends in Microbiology</i> , 1994 , 2, 4-6	12.4	53

27	Lymphokine production by human T cells in disease states. <i>Annual Review of Immunology</i> , 1994 , 12, 227-57.7	57.7	1155
26	Membrane tumour necrosis factor-alpha is involved in the polyclonal B-cell activation induced by HIV-infected human T cells. <i>Nature</i> , 1993 , 363, 464-6	50.4	131
25	Allergen exposure induces the activation of allergen-specific Th2 cells in the airway mucosa of patients with allergic respiratory disorders. <i>European Journal of Immunology</i> , 1993 , 23, 1445-9	6.1	306
24	Human TH1 and TH2 Subsets. <i>International Archives of Allergy and Immunology</i> , 1992 , 99, 242-245	3.7	14
23	Induction of TH1 and TH2 responses: a key role for the Natural Immune response?. <i>Trends in Immunology</i> , 1992 , 13, 379-81		685
22	Both CD8+ and CD16+ human T cell clones can provide B cell help for immunoglobulin production. <i>International Journal of Clinical and Laboratory Research</i> , 1992 , 22, 36-9		1
21	Aberrant interleukin (IL)-4 and IL-5 production in vitro by CD4+ helper T cells from atopic subjects. <i>European Journal of Immunology</i> , 1992 , 22, 1615-20	6.1	73
20	Human TH1 and TH2 subsets: doubt no more. <i>Trends in Immunology</i> , 1991 , 12, 256-7		996
19	The IgE response in atopy and infections. <i>Clinical and Experimental Allergy</i> , 1991 , 21 Suppl 1, 72-8	4.1	5
18	In vitro infection with HIV of antigen-specific T cell clones derived from HIV-seronegative individuals. Effects on cytokine production and helper function. <i>Research in Clinic and Laboratory</i> , 1991 , 21, 85-90		1
17	T lymphocytes from tonsil and peripheral blood show different cytolytic and helper activities. <i>Acta Oto-Laryngologica</i> , 1991 , 111, 974-80	1.6	1
16	Cytokine production by allergen (Der pI)-specific CD4+ T cell clones derived from a patient with severe atopic disease. <i>International Journal of Clinical and Laboratory Research</i> , 1991 , 21, 186-9		18
15	A novel pathway of human B cell activation initiated by CK226 surface antigen. <i>European Journal of Immunology</i> , 1990 , 20, 1161-5	6.1	2
14	Regulation and deregulation of human IgE synthesis. <i>Trends in Immunology</i> , 1990 , 11, 316-21		228
13	Cytokine action on B cells in disease 1990 , 215-251		2
12	Regulatory mechanisms of in vitro human IgE synthesis*. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 1989 , 44, 9-15	9.3	6
11	Role of interleukins in induction and regulation of human IgE synthesis. <i>Clinical Immunology and Immunopathology</i> , 1989 , 50, S13-23		77
10	The immune derangement and strategies for immunotherapy. <i>Cancer Treatment and Research</i> , 1989 , 41, 53-88	3.5	1

9	Profiles of lymphokine activities and helper function for IgE in human T cell clones. <i>European Journal of Immunology</i> , 1988 , 18, 1045-50	6.1	198
8	Diminished Production of Interleukin 2 and Gamma-Interferon by Cloned T _H Cells from Patients with the Acquired Immunodeficiency Syndrome (AIDS). <i>Tumori</i> , 1987 , 73, 273-278	1.7	1
7	Reduced production of interleukin 2 and interferon-gamma and enhanced helper activity for IgG synthesis by cloned CD4+ T cells from patients with AIDS. <i>European Journal of Immunology</i> , 1987 , 17, 1685-90	6.1	88
6	Analysis of the role of interferon-gamma, interleukin 2 and a third factor distinct from interferon-gamma and interleukin 2 in human B cell proliferation. Evidence that they act at different times after B cell activation. <i>European Journal of Immunology</i> , 1986 , 16, 623-9	6.1	42
5	Human T cell clones can induce in vitro IgE synthesis in normal B cells regardless of alloantigen recognition or specificity for peculiar antigens. <i>European Journal of Immunology</i> , 1986 , 16, 1509-14	6.1	28
4	Heterogeneity of B cell growth factor (BCGF)-producing T cells in humans. Clonal analysis of BCGF-producing cells within T4+ and T8+ subsets and evidence for the involvement of different growth factors in different BCGF assays. <i>Research in Clinic and Laboratory</i> , 1986 , 16, 23-8		
3	Synergy of B cell growth factor and interleukin 2 in the proliferation of activated human B cells. <i>European Journal of Immunology</i> , 1985 , 15, 1158-64	6.1	19
2	Frequent coexpression of cytolytic activity and lymphokine production among human T lymphocytes. Production of B cell growth factor and interleukin 2 by T8+ and T4+ cytolytic clones. <i>European Journal of Immunology</i> , 1984 , 14, 1066-9	6.1	37
1	Short- and long-term effects of radiation on T-cell subsets in peripheral blood of patients with Hodgkin's disease. <i>Cancer</i> , 1980 , 46, 2590-5	6.4	14