Amir Schajnovitz

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/10967091/publications.pdf Version: 2024-02-01



AMID SCHAINOVITZ

#	Article	IF	CITATIONS
1	Induction of a Timed Metabolic Collapse to Overcome Cancer Chemoresistance. Cell Metabolism, 2020, 32, 391-403.e6.	16.2	79
2	Rapid Mobilization Reveals a Highly Engraftable Hematopoietic Stem Cell. Cell, 2018, 172, 191-204.e10.	28.9	92
3	The Wave2 scaffold Hem-1 is required for transition of fetal liver hematopoiesis to bone marrow. Nature Communications, 2018, 9, 2377.	12.8	15
4	Distinct bone marrow blood vessels differentially regulate haematopoiesis. Nature, 2016, 532, 323-328.	27.8	553
5	Inhibition of Dihydroorotate Dehydrogenase Overcomes Differentiation Blockade in Acute Myeloid Leukemia. Cell, 2016, 167, 171-186.e15.	28.9	353
6	Non-genotoxic conditioning for hematopoietic stem cell transplantation using a hematopoietic-cell-specific internalizing immunotoxin. Nature Biotechnology, 2016, 34, 738-745.	17.5	176
7	Inhibition of the Enzyme Dihydroorotate Dehydrogenase Overcomes Differentiation Blockade in Acute Myeloid Leukemia. Blood, 2016, 128, 1656-1656.	1.4	3
8	Rapid Mobilization Reveals a Highly Engraftable Hematopoietic Stem Cell. Blood, 2016, 128, 368-368.	1.4	0
9	Distinct Bone Marrow Blood Vessels Differentially Regulate Normal and Malignant Hematopoietic Stem and Progenitor Cells. Blood, 2015, 126, 664-664.	1.4	1
10	Bone's dark side: mutated osteoblasts implicated in leukemia. Cell Research, 2014, 24, 383-384.	12.0	5
11	Inhibiting stromal cell heparan sulfate synthesis improves stem cell mobilization and enables engraftment without cytotoxic conditioning. Blood, 2014, 124, 2937-2947.	1.4	39
12	Human and Murine β-Defensin-Derived Peptides Induce Rapid Mobilization Of Murine Hematopoietic Stem and Progenitor Cells Via Activation Of CXCR4 Signaling and CXCL12 Release. Blood, 2013, 122, 890-890.	1.4	0
13	S1P promotes murine progenitor cell egress and mobilization via S1P1-mediated ROS signaling and SDF-1 release. Blood, 2012, 119, 2478-2488.	1.4	175
14	FGF-2 expands murine hematopoietic stem and progenitor cells via proliferation of stromal cells, c-Kit activation, and CXCL12 down-regulation. Blood, 2012, 120, 1843-1855.	1.4	99
15	Monocytes-macrophages that express α-smooth muscle actin preserve primitive hematopoietic cells in the bone marrow. Nature Immunology, 2012, 13, 1072-1082.	14.5	196
16	Enhanced c-Met activity promotes G-CSF–induced mobilization of hematopoietic progenitor cells via ROS signaling. Blood, 2011, 117, 419-428.	1.4	114
17	CXCL12 secretion by bone marrow stromal cells is dependent on cell contact and mediated by connexin-43 and connexin-45 gap junctions. Nature Immunology, 2011, 12, 391-398.	14.5	142
18	Rapid mobilization of hematopoietic progenitors by AMD3100 and catecholamines is mediated by CXCR4-dependent SDF-1 release from bone marrow stromal cells. Leukemia, 2011, 25, 1286-1296.	7.2	180

#	Article	IF	CITATIONS
19	The Chemotactic Lipid S1P Regulates Hematopoietic Progenitor Cell Egress and Mobilization Via Its Major Receptor S1P1 and by SDF-1 Inhibition In a p38/Akt/mTOR Dependent Manner. Blood, 2010, 116, 553-553.	1.4	7
20	Expansion of Normal and Leukemic Hematopoietic Progenitor Cells by PTH Requires bFGF Activation of c-Kit and Its Downstream JAK2/STAT5 Signaling Blood, 2009, 114, 2511-2511.	1.4	1
21	Functional SDF-1 Secretion from BM Stromal Cells Is a Cell Contact-Dependent Event Mediated by Cx43 and Cx45 Gap-Junctions. Blood, 2008, 112, 319-319.	1.4	0
22	In Vivo Mobilization of Leukemic Human Precursor-B-ALL Cells by the CXCR4-Antagonist AMD3100 Is Via Secretion of SDF-1 and Synergistically by Catecholamine Action Blood, 2008, 112, 1920-1920.	1.4	0
23	Induction of a Timed Metabolic Collapse to Overcome Cancer Chemoresistance. SSRN Electronic Journal, 0, , .	0.4	0