

Stine Julie Tingskov

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10989428/publications.pdf>

Version: 2024-02-01

13
papers

175
citations

1163117

8
h-index

1199594

12
g-index

13
all docs

13
docs citations

13
times ranked

251
citing authors

#	ARTICLE	IF	CITATIONS
1	EP ₁ receptor antagonism mitigates early and late stage renal fibrosis. <i>Acta Physiologica</i> , 2022, 234, e13780.	3.8	6
2	Tamoxifen attenuates renal fibrosis in human kidney slices and rats subjected to unilateral ureteral obstruction. <i>Biomedicine and Pharmacotherapy</i> , 2021, 133, 111003.	5.6	13
3	Olive Leaf Extract (OLE) impaired vasopressin-induced aquaporin-2 trafficking through the activation of the calcium-sensing receptor. <i>Scientific Reports</i> , 2021, 11, 4537.	3.3	11
4	Local Inhibition of Indoleamine 2,3-Dioxygenase Mitigates Renal Fibrosis. <i>Biomedicines</i> , 2021, 9, 856.	3.2	5
5	Sorting Nexin 27 Regulates the Lysosomal Degradation of Aquaporin-2 Protein in the Kidney Collecting Duct. <i>Cells</i> , 2020, 9, 1208.	4.1	17
6	Estrogen regulates aquaporin-2 expression in the kidney. <i>Vitamins and Hormones</i> , 2020, 112, 243-264.	1.7	5
7	Vasopressin-Independent Regulation of Aquaporin-2 by Tamoxifen in Kidney Collecting Ducts. <i>Frontiers in Physiology</i> , 2019, 10, 948.	2.8	8
8	Activation of the prostaglandin E ₂ EP ₂ receptor attenuates renal fibrosis in unilateral ureteral obstructed mice and human kidney slices. <i>Acta Physiologica</i> , 2019, 227, e13291.	3.8	41
9	Administration of a selective prostaglandin E ₂ receptor agonist attenuates renal fibrosis in vitro and in vivo. <i>FASEB Journal</i> , 2019, 33, 863.9.	0.5	0
10	Tamoxifen Decreases Lithium-Induced Natriuresis in Rats With Nephrogenic Diabetes Insipidus. <i>Frontiers in Physiology</i> , 2018, 9, 903.	2.8	7
11	Tamoxifen attenuates development of lithium-induced nephrogenic diabetes insipidus in rats. <i>American Journal of Physiology - Renal Physiology</i> , 2018, 314, F1020-F1025.	2.7	15
12	Theranostic poly(lactic-co-glycolic acid) nanoparticle for magnetic resonance/infrared fluorescence bimodal imaging and efficient siRNA delivery to macrophages and its evaluation in a kidney injury model. <i>Nanomedicine: Nanotechnology, Biology, and Medicine</i> , 2017, 13, 2451-2462.	3.3	27
13	Can bladder fibrosis in congenital urinary tract obstruction be reversed?. <i>Journal of Pediatric Urology</i> , 2017, 13, 574-580.	1.1	20