

Rohit Singhal

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/10992251/publications.pdf>

Version: 2024-02-01

14
papers

205
citations

1163117

8
h-index

1281871

11
g-index

14
all docs

14
docs citations

14
times ranked

278
citing authors

#	ARTICLE	IF	CITATIONS
1	Soy protein isolate feeding does not result in reproductive toxicity in the pre-pubertal rat testis. <i>Experimental Biology and Medicine</i> , 2018, 243, 695-707.	2.4	4
2	RNA-sequencing data analysis of uterus in ovariectomized rats fed with soy protein isolate, 17 β -estradiol and casein. <i>Data in Brief</i> , 2016, 7, 1491-1496.	1.0	0
3	Uterine responses to feeding soy protein isolate and treatment with 17 β -estradiol differ in ovariectomized female rats. <i>Toxicology and Applied Pharmacology</i> , 2016, 297, 68-80.	2.8	11
4	Mammary Gland Morphology and Gene Expression Differ in Female Rats Treated with 17 β -Estradiol or Fed Soy Protein Isolate. <i>Endocrinology</i> , 2012, 153, 6021-6032.	2.8	17
5	Complement Activation in Acetaminophen-Induced Liver Injury in Mice. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2012, 341, 377-385.	2.5	40
6	Uterine physiological responses and global gene expression in ovariectomized (OVX) rats treated with soy protein isolate (SPI) or 17 β -estradiol. <i>FASEB Journal</i> , 2012, 26, 243.2.	0.5	0
7	Carbohydrate-Responsive Gene Expression in the Adipose Tissue of Rats. <i>Endocrinology</i> , 2010, 151, 153-164.	2.8	37
8	Feeding soy protein isolate (SPI) does not result in an estrogenic gene expression profile in the mammary of ovariectomized (OVX) female rats. <i>FASEB Journal</i> , 2010, 24, 212.2.	0.5	0
9	Hepatic gene expression following consumption of soy protein isolate in female Sprague-Dawley rats differs from that produced by 17 β -estradiol treatment. <i>Journal of Endocrinology</i> , 2009, 202, 141-152.	2.6	26
10	Estrogenic status modulates the effect of soy on hepatic responses to 7,12-dimethylbenz(a)anthracene (DMBA). <i>Toxicology and Applied Pharmacology</i> , 2009, 234, 89-97.	2.8	7
11	Rats fed soy protein isolate (SPI) have impaired hepatic CYP1A1 induction by polycyclic aromatic hydrocarbons as a result of interference with aryl hydrocarbon receptor signaling. <i>Toxicology and Applied Pharmacology</i> , 2008, 227, 275-283.	2.8	11
12	Estrogenic status modulates aryl hydrocarbon receptor-mediated hepatic gene expression and carcinogenicity. <i>Carcinogenesis</i> , 2008, 29, 227-236.	2.8	35
13	Reduction in 7,12-Dimethylbenz[a]anthracene-Induced Hepatic Cytochrome-P450 1A1 Expression Following Soy Consumption in Female Rats Is Mediated by Degradation of the Aryl Hydrocarbon Receptor. <i>Journal of Nutrition</i> , 2007, 137, 19-24.	2.9	16
14	IS SOY ESTROGENIC? HEPATIC GENE EXPRESSION IN THE PRESENCE OR ABSENCE OF ENDOGENOUS ESTROGEN.. <i>FASEB Journal</i> , 2007, 21, A61.	0.5	1