

# Junming Li

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/1099954/junming-li-publications-by-year.pdf>  
**Version:** 2024-04-10

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.  
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

45 papers	785 citations	16 h-index	26 g-index
53 ext. papers	1,115 ext. citations	4 avg, IF	4.33 L-index

#	Paper	IF	Citations
45	PPAR $\delta$ Ameliorates H37Ra-Induced Foamy Macrophage Formation via the ABCG1-Dependent Cholesterol Efflux Pathway in THP-1 Macrophages.. <i>Frontiers in Microbiology</i> , <b>2022</b> , 13, 829870	5.7	
44	Expression profile and diagnostic value of circRNAs in peripheral blood from patients with systemic lupus erythematosus. <i>Molecular Medicine Reports</i> , <b>2021</b> , 23,	2.9	10
43	Molecular Analysis and Antimicrobial Resistance Pattern of Tigecycline-Non-Susceptible Isolated from a Tertiary Care Hospital of East Asia. <i>Infection and Drug Resistance</i> , <b>2021</b> , 14, 4147-4155	4.2	0
42	Tim-4 expressing monocytes as a novel indicator to assess disease activity and severity of ulcerative colitis. <i>Life Sciences</i> , <b>2021</b> , 269, 119077	6.8	1
41	Circular RNAs hsa-circ0000175 and hsa-circ0044235 in plasma are novel biomarkers for new-onset rheumatoid arthritis. <i>Autoimmunity</i> , <b>2021</b> , 54, 234-242	3	2
40	Increased TIM-3PD-1 NK cells are associated with the disease activity and severity of systemic lupus erythematosus. <i>Clinical and Experimental Medicine</i> , <b>2021</b> , 1	4.9	0
39	Characteristics of immune cell infiltration and associated diagnostic biomarkers in ulcerative colitis: results from bioinformatics analysis. <i>Bioengineered</i> , <b>2021</b> , 12, 252-265	5.7	14
38	Elevated frequencies of CD14HLA-DR MDSCs in COVID-19 patients. <i>Aging</i> , <b>2021</b> , 13, 6236-6246	5.6	5
37	T-Cell Subset Counts in Peripheral Blood Can Be Used as Discriminatory Biomarkers for Diagnosis and Severity Prediction of Coronavirus Disease 2019. <i>Journal of Infectious Diseases</i> , <b>2020</b> , 222, 198-202	7	82
36	The study of METTL14, ALKBH5, and YTHDF2 in peripheral blood mononuclear cells from systemic lupus erythematosus. <i>Molecular Genetics &amp; Genomic Medicine</i> , <b>2020</b> , 8, e1298	2.3	17
35	Decreased Peripheral Blood Correlates with Markers of Autoimmune Response in Systemic Lupus Erythematosus. <i>Disease Markers</i> , <b>2020</b> , 2020, 8193895	3.2	15
34	Circular RNAs hsa_circ_0000479 in peripheral blood mononuclear cells as novel biomarkers for systemic lupus erythematosus. <i>Autoimmunity</i> , <b>2020</b> , 53, 167-176	3	12
33	Long non-coding RNA expression profiling of macrophage line RAW264.7 infected by. <i>Biotechnic and Histochemistry</i> , <b>2020</b> , 95, 403-410	1.8	2
32	Up-regulation of circRNA-0003528 promotes mycobacterium tuberculosis associated macrophage polarization via down-regulating miR-224-5p, miR-324-5p and miR-488-5p and up-regulating CTLA4. <i>Aging</i> , <b>2020</b> , 12, 25658-25672	5.6	10
31	Expression and clinical significance of circular RNA hsa_circ_0079787 in the peripheral blood of patients with axial spondyloarthritis. <i>Molecular Medicine Reports</i> , <b>2020</b> , 22, 4197-4206	2.9	2
30	Expression and clinical significance of circular RNAs hsa_circ_0000175 and hsa_circ_0008410 in peripheral blood mononuclear cells from patients with rheumatoid arthritis. <i>International Journal of Molecular Medicine</i> , <b>2020</b> , 45, 1203-1212	4.4	9
29	Immune characteristics distinguish patients with severe disease associated with SARS-CoV-2. <i>Immunologic Research</i> , <b>2020</b> , 68, 398-404	4.3	10

28	Novel serological biomarkers for inflammation in predicting disease severity in patients with COVID-19. <i>International Immunopharmacology</i> , <b>2020</b> , 89, 107065	5.8	32
27	Decreased , , and in Peripheral Blood Are as Risk Factors for Rheumatoid Arthritis. <i>BioMed Research International</i> , <b>2020</b> , 2020, 5735279	3	18
26	Low-Density Granulocytes Affect T-SPOT.TB Assay by Inhibiting the Production of Interferon- $\gamma$ in T Cells via PD-L1/PD-1 Pathway. <i>Frontiers in Microbiology</i> , <b>2020</b> , 11, 622389	5.7	5
25	Peripheral blood circular RNA hsa_circ_0082688-hsa_circ_0008675 can be used as a candidate biomarker of systemic lupus erythematosus with renal involvement. <i>Clinical and Experimental Rheumatology</i> , <b>2020</b> , 38, 822-833	2.2	2
24	Identification of circular RNAs hsa_circ_0044235 and hsa_circ_0068367 as novel biomarkers for systemic lupus erythematosus. <i>International Journal of Molecular Medicine</i> , <b>2019</b> , 44, 1462-1472	4.4	17
23	Serum PGLYRP-1 is a highly discriminatory biomarker for the diagnosis of rheumatoid arthritis. <i>Molecular Medicine Reports</i> , <b>2019</b> , 19, 589-594	2.9	6
22	Circular RNAs Hsa_circ_0002715 and Hsa_circ_0035197 in Peripheral Blood Are Novel Potential Biomarkers for New-Onset Rheumatoid Arthritis. <i>Disease Markers</i> , <b>2019</b> , 2019, 2073139	3.2	18
21	Elevated expression of PD-1 on T cells correlates with disease activity in rheumatoid arthritis. <i>Molecular Medicine Reports</i> , <b>2018</b> , 17, 3297-3305	2.9	7
20	Overexpression of CD64 on CD14CD16 and CD14CD16 monocytes of rheumatoid arthritis patients correlates with disease activity. <i>Experimental and Therapeutic Medicine</i> , <b>2018</b> , 16, 2703-2711	2.1	6
19	Integrative analysis of long non-coding RNAs and messengerRNA expression profiles in systemic lupus erythematosus. <i>Molecular Medicine Reports</i> , <b>2018</b> , 17, 3489-3496	2.9	24
18	Circulating circular RNAs hsa_circ_0001204 and hsa_circ_0001747 act as diagnostic biomarkers for active tuberculosis detection. <i>International Journal of Clinical and Experimental Pathology</i> , <b>2018</b> , 11, 586-594	1.4	6
17	Decreased expression of TIGIT in NK cells correlates negatively with disease activity in systemic lupus erythematosus. <i>International Journal of Clinical and Experimental Pathology</i> , <b>2018</b> , 11, 2408-2418	1.4	5
16	Multicentre laboratory validation of the nitrate reductase assay using liquid medium for the rapid detection of multidrug-resistant and extensively drug-resistant Mycobacterium tuberculosis. <i>Tuberculosis</i> , <b>2018</b> , 113, 242-248	2.6	
15	Plasma Circular RNAs hsa_circ_0001953 and hsa_circ_0009024 as Diagnostic Biomarkers for Active Tuberculosis. <i>Frontiers in Microbiology</i> , <b>2018</b> , 9, 2010	5.7	22
14	A genome-wide association study identifies six novel risk loci for primary biliary cholangitis. <i>Nature Communications</i> , <b>2017</b> , 8, 14828	17.4	66
13	Elevated expression of TIGIT on CD3CD4 T cells correlates with disease activity in systemic lupus erythematosus. <i>Allergy, Asthma and Clinical Immunology</i> , <b>2017</b> , 13, 15	3.2	11
12	Identification of differentially expressed circular RNAs in human monocyte derived macrophages response to Mycobacterium tuberculosis infection. <i>Scientific Reports</i> , <b>2017</b> , 7, 13673	4.9	16
11	Comprehensive analysis of long non-coding RNA and mRNA expression profiles in rheumatoid arthritis. <i>Experimental and Therapeutic Medicine</i> , <b>2017</b> , 14, 5965-5973	2.1	25

10	Elevated Expression of Immunoreceptor Tyrosine-Based Inhibitory Motif (TIGIT) on T Lymphocytes is Correlated with Disease Activity in Rheumatoid Arthritis. <i>Medical Science Monitor</i> , <b>2017</b> , 23, 1232-1241 <sup>3,2</sup>	17
9	Identification of Differentially Expressed Long Non-coding RNAs in Polarized Macrophages. <i>Scientific Reports</i> , <b>2016</b> , 6, 19705	46
8	PD-L1-expressing neutrophils as a novel indicator to assess disease activity and severity of systemic lupus erythematosus. <i>Arthritis Research and Therapy</i> , <b>2016</b> , 18, 47	37
7	Low-Density Granulocytes Are Elevated in Mycobacterial Infection and Associated with the Severity of Tuberculosis. <i>PLoS ONE</i> , <b>2016</b> , 11, e0153567	49
6	EBP50 induces apoptosis in macrophages by upregulating nitric oxide production to eliminate intracellular Mycobacterium tuberculosis. <i>Scientific Reports</i> , <b>2016</b> , 6, 18961	8
5	Evaluation of the microscopic observation drug susceptibility assay for the rapid detection of MDR-TB and XDR-TB in China: a prospective multicentre study. <i>Journal of Antimicrobial Chemotherapy</i> , <b>2015</b> , 70, 456-62	12
4	Mycobacterium tuberculosis-Induced Polarization of Human Macrophage Orchestrates the Formation and Development of Tuberculous Granulomas In Vitro. <i>PLoS ONE</i> , <b>2015</b> , 10, e0129744	95
3	Rapid diagnosis of pleural tuberculosis by Xpert MTB/RIF assay using pleural biopsy and pleural fluid specimens. <i>Journal of Research in Medical Sciences</i> , <b>2015</b> , 20, 26-31	27
2	Evaluation of MODS assay for rapid detection of Mycobacterium tuberculosis resistance to second-line drugs in a tertiary care tuberculosis hospital in China. <i>Tuberculosis</i> , <b>2014</b> , 94, 506-10	5
1	The 10-23 DNA enzyme generated by a novel expression vector mediate inhibition of taco expression in macrophage. <i>Oligonucleotides</i> , <b>2010</b> , 20, 61-8	11