

# Faiz Ali

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11011117/publications.pdf>

Version: 2024-02-01

17  
papers

319  
citations

1162889

8  
h-index

887953

17  
g-index

17  
all docs

17  
docs citations

17  
times ranked

313  
citing authors

#	ARTICLE	IF	CITATIONS
1	Recent applications of molecular imprinted polymers for enantio-selective recognition. <i>Talanta</i> , 2013, 106, 45-59.	2.9	87
2	Comprehensive overview of recent preparation and application trends of various open tubular capillary columns in separation science. <i>Journal of Chromatography A</i> , 2013, 1308, 1-24.	1.8	72
3	Catalyst assisted synthesis of initiator attached silica monolith particles via isocyanate-hydroxyl reaction for production of polystyrene bound chromatographic stationary phase of excellent separation efficiency. <i>Journal of Chromatography A</i> , 2014, 1324, 115-120.	1.8	22
4	Polystyrene bound stationary phase of excellent separation efficiency based on partially sub-2 $\mu$ m silica monolith particles. <i>Journal of Chromatography A</i> , 2013, 1303, 9-17.	1.8	21
5	Sedimentation assisted preparation of ground particles of silica monolith and their C18 modification resulting in a chromatographic phase of improved separation efficiency. <i>Journal of Chromatography A</i> , 2017, 1525, 79-86.	1.8	18
6	Open tubular capillary column for the separation of cytochrome C tryptic digest in capillary electrochromatography. <i>Journal of Separation Science</i> , 2015, 38, 3645-3654.	1.3	16
7	C <sub>18</sub> -bound porous silica monolith particles as a low-cost high-performance liquid chromatography stationary phase with an excellent chromatographic performance. <i>Journal of Separation Science</i> , 2014, 37, 3426-3434.	1.3	15
8	Open tubular capillary electrochromatography with an N-phenylacrylamide-styrene copolymer-based stationary phase for the separation of anomers of glucose and structural isomers of maltotriose. <i>Journal of Separation Science</i> , 2015, 38, 1763-1770.	1.3	15
9	Cheap C <sub>18</sub> -modified Silica Monolith Particles as HPLC Stationary Phase of Good Separation Efficiency. <i>Bulletin of the Korean Chemical Society</i> , 2015, 36, 1733-1736.	1.0	8
10	High Efficiency Robust Open Tubular Capillary Electrochromatography Column for the Separation of Peptides. <i>Bulletin of the Korean Chemical Society</i> , 2016, 37, 1374-1377.	1.0	8
11	Mixed-mode open tubular column for peptide separations by capillary electrochromatography. <i>Journal of Separation Science</i> , 2021, 44, 2602-2611.	1.3	8
12	Immobilization of Styrene-acrylamide Co-polymer on Either Silica Particles or Inner Surface of Silica Capillary for the Separation of D-Glucose Anomers. <i>Bulletin of the Korean Chemical Society</i> , 2014, 35, 539-545.	1.0	8
13	Particle packed mixed-mode chromatographic stationary phase for the separation of peptide in liquid chromatography. <i>Journal of Separation Science</i> , 2021, 44, 1430-1439.	1.3	6
14	Synthesis, column packing and liquid chromatography of molecularly imprinted polymers for the acid black 1, acid black 210, and acid Brown 703 dyes. <i>RSC Advances</i> , 2022, 12, 19611-19623.	1.7	5
15	Ground Organic Monolith Particles Having a Large Volume of Macropores as Chromatographic Separation Media. <i>Bulletin of the Korean Chemical Society</i> , 2014, 35, 2033-2037.	1.0	4
16	Demonstration of high separation efficiency for polystyrene-modified sub-1 $\mu$ m particles originating from silica monolith under isocratic elution mode in liquid chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2019, 42, 662-672.	0.5	3
17	100 Micrometer bore open tubular capillary column modified with linear co-polymer chains for application in low pressure liquid chromatography. <i>Journal of Liquid Chromatography and Related Technologies</i> , 2020, 43, 66-73.	0.5	3