Douglas Londono

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11016843/publications.pdf

Version: 2024-02-01

933447 996975 15 807 10 15 citations g-index h-index papers 15 15 15 1090 docs citations times ranked citing authors all docs

| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Investigation of previously implicated genetic variants in chronic tic disorders: a transmission disequilibrium test approach. European Archives of Psychiatry and Clinical Neuroscience, 2018, 268, 301-316. | 3.2 | 23 |
| 2 | Bone marrow-derived epithelial cells and hair follicle stem cells contribute to development of chronic cutaneous neoplasms. Nature Communications, 2018, 9, 5293. | 12.8 | 9 |
| 3 | An Analytic Solution to the Computation of Power and Sample Size for Genetic Association Studies under a Pleiotropic Mode of Inheritance. Human Heredity, 2016, 81, 194-209. | 0.8 | 3 |
| 4 | A PAX1 enhancer locus is associated with susceptibility to idiopathic scoliosis in females. Nature Communications, 2015, 6, 6452. | 12.8 | 122 |
| 5 | Chromatin Profiling Reveals Regulatory Network Shifts and a Protective Role for Hepatocyte Nuclear Factor 4α during Colitis. Molecular and Cellular Biology, 2014, 34, 3291-3304. | 2.3 | 41 |
| 6 | Mapping genes with longitudinal phenotypes via Bayesian posterior probabilities. BMC Proceedings, 2014, 8, S81. | 1.6 | 6 |
| 7 | A meta-analysis identifies adolescent idiopathic scoliosis association with <i>LBX1 </i> locus in multiple ethnic groups. Journal of Medical Genetics, 2014, 51, 401-406. | 3.2 | 79 |
| 8 | Genetic variants in GPR126 are associated with adolescent idiopathic scoliosis. Nature Genetics, 2013, 45, 676-679. | 21.4 | 240 |
| 9 | A novel method for analyzing genetic association with longitudinal phenotypes. Statistical Applications in Genetics and Molecular Biology, 2013, 12, 241-61. | 0.6 | 16 |
| 10 | TDT-HET: A new transmission disequilibrium test that incorporates locus heterogeneity into the analysis of family-based association data. BMC Bioinformatics, 2012, 13, 13. | 2.6 | 27 |
| 11 | Single-Variant and Multi-Variant Trend Tests for Genetic Association with Next-Generation Sequencing That Are Robust to Sequencing Error. Human Heredity, 2012, 74, 172-183. | 0.8 | 10 |
| 12 | Genome-wide association studies of adolescent idiopathic scoliosis suggest candidate susceptibility genes. Human Molecular Genetics, 2011, 20, 1456-1466. | 2.9 | 172 |
| 13 | <i>EDNRA</i> variants associate with smooth muscle mRNA levels, cell proliferation rates, and cystic fibrosis pulmonary disease severity. Physiological Genomics, 2010, 41, 71-77. | 2.3 | 34 |
| 14 | A Cost-Effective Statistical Method to Correct for Differential Genotype Misclassification When Performing Case-Control Genetic Association. Human Heredity, 2010, 70, 102-108. | 0.8 | 9 |
| 15 | Beta 2 adrenergic receptor polymorphisms in cystic fibrosis. Pediatric Pulmonology, 2005, 39, 544-550. | 2.0 | 16 |