

Gerardo F. Barbero

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/1102044/gerardo-f-barbero-publications-by-year.pdf>

Version: 2024-04-23

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

110
papers

2,801
citations

29
h-index

49
g-index

113
ext. papers

3,420
ext. citations

4.8
avg, IF

5.59
L-index

#	Paper	IF	Citations
110	Optimization of an Ultrasound-Assisted Extraction Method Applied to the Extraction of Flavonoids from Moringa Leaves (<i>Moringa oleifera</i> Lam.). <i>Agronomy</i> , 2022 , 12, 261	3.6	1
109	Toxic elements and trace elements in <i>Macrolepiota procera</i> mushrooms from southern Spain and northern Morocco. <i>Journal of Food Composition and Analysis</i> , 2022 , 108, 104419	4.1	3
108	EscapeWine!. <i>Advances in Game-based Learning Book Series</i> , 2022 , 356-375	0.5	
107	A comparison study between ultrasound-assisted and enzyme-assisted extraction of anthocyanins from blackcurrant (L.). <i>Food Chemistry: X</i> , 2022 , 13, 100192	4.7	2
106	Comparison of different processing approaches by SVM and RF on HS-MS eNose and NIR Spectrometry data for the discrimination of gasoline samples. <i>Microchemical Journal</i> , 2022 , 172, 106893 ^{4.8}	4.8	2
105	Analysis of Compounds with Oenological Interest in Somatic Variants of Grapevines. <i>Horticulturae</i> , 2022 , 8, 22	2.5	0
104	Extraction of Antioxidant Compounds from Onion Bulb (<i>Allium cepa</i> L.) Using Individual and Simultaneous Microwave-Assisted Extraction Methods. <i>Antioxidants</i> , 2022 , 11, 846	7.1	1
103	Co-precipitation of grape residue extract using sub- and supercritical CO ₂ technology. <i>Journal of CO₂ Utilization</i> , 2022 , 61, 102010	7.6	2
102	Exposure to Essential and Toxic Elements via Consumption of Agaricaceae, Amanitaceae, Boletaceae, and Russulaceae Mushrooms from Southern Spain and Northern Morocco. <i>Journal of Fungi (Basel, Switzerland)</i> , 2022 , 8, 545	5.6	0
101	Optimization through a BoxBehnken Experimental Design of the Microwave-Assisted Extraction of the Psychoactive Compounds in Hallucinogenic Fungi (<i>Psilocibe cubensis</i>). <i>Journal of Fungi (Basel, Switzerland)</i> , 2022 , 8, 598	5.6	1
100	Development of Optimized Ultrasound-Assisted Extraction Methods for the Recovery of Total Phenolic Compounds and Anthocyanins from Onion Bulbs. <i>Antioxidants</i> , 2021 , 10,	7.1	2
99	An Evaluation of the Equilibrium Properties in Hexane and Ethanol Extractive Systems for <i>Moringa oleifera</i> Seeds and Fatty Acid Profiles of the Extracts. <i>Separations</i> , 2021 , 8, 217	3.1	2
98	Development of a Rapid UHPLC-PDA Method for the Simultaneous Quantification of Flavonol Contents in Onions (L.). <i>Pharmaceuticals</i> , 2021 , 14,	5.2	1
97	Optimization of an Ultrasound-Assisted Extraction Method for the Analysis of Major Anthocyanin Content in Flowers. <i>Molecules</i> , 2021 , 26,	4.8	1
96	Optimization by Means of Chemometric Tools of an Ultrasound-Assisted Method for the Extraction of Betacyanins from Red Dragon Fruit (<i>Hylocereus polyrhizus</i>). <i>Agronomy</i> , 2021 , 11, 1053	3.6	0
95	A Methodology Based on FT-IR Data Combined with Random Forest Model to Generate for the Characterization of High-Quality Vinegars. <i>Foods</i> , 2021 , 10,	4.9	2
94	Metal concentrations in <i>Lactarius</i> mushroom species collected from Southern Spain and Northern Morocco: Evaluation of health risks and benefits. <i>Journal of Food Composition and Analysis</i> , 2021 , 99, 103859	4.1	2

93	Optimization of the Microwave-Assisted Extraction of Simple Phenolic Compounds from Grape Skins and Seeds. <i>Agronomy</i> , 2021 , 11, 1527	3.6	2
92	Development of a rapid and accurate UHPLC-PDA-FL method for the quantification of phenolic compounds in grapes. <i>Food Chemistry</i> , 2021 , 334, 127569	8.5	14
91	Volatile and Semi-Volatile Organic Compounds May Help Reduce Pollinator-Prey Overlap in the Carnivorous Plant <i>Drosophyllum lusitanicum</i> (Drosophyllaceae). <i>Journal of Chemical Ecology</i> , 2021 , 47, 73-86	2.7	2
90	Optimization and Comparison of Ultrasound and Microwave-Assisted Extraction of Phenolic Compounds from Cotton-Lavender (<i>Santolina chamaecyparissus</i> L.). <i>Agronomy</i> , 2021 , 11, 84	3.6	9
89	Flavonol Composition and Antioxidant Activity of Onions (L.) Based on the Development of New Analytical Ultrasound-Assisted Extraction Methods. <i>Antioxidants</i> , 2021 , 10,	7.1	15
88	Extraction and Analysis of Natural Product in Plant. <i>Agronomy</i> , 2021 , 11, 415	3.6	3
87	A Novel Ultrasound-Assisted Extraction Method for the Analysis of Anthocyanins in Potatoes (L.). <i>Antioxidants</i> , 2021 , 10,	7.1	5
86	How Different Cooking Methods Affect the Phenolic Composition of Sweet Potato for Human Consumption (<i>Ipomea batata</i> (L.) Lam). <i>Agronomy</i> , 2021 , 11, 1636	3.6	1
85	Simultaneous determination by UHPLC-PDA of major capsaicinoids and capsinoids contents in peppers. <i>Food Chemistry</i> , 2021 , 356, 129688	8.5	1
84	Optimization of a Novel Method Based on Ultrasound-Assisted Extraction for the Quantification of Anthocyanins and Total Phenolic Compounds in Blueberry Samples (L.). <i>Foods</i> , 2020 , 9,	4.9	12
83	Characterization of Biodegraded Ignitable Liquids by Headspace-Ion Mobility Spectrometry. <i>Sensors</i> , 2020 , 20,	3.8	1
82	Concentration of bioactive compounds from grape marc using pressurized liquid extraction followed by integrated membrane processes. <i>Separation and Purification Technology</i> , 2020 , 250, 117206	8.3	11
81	Characterization of Arabica and Robusta Coffees by Ion Mobility Sum Spectrum. <i>Sensors</i> , 2020 , 20,	3.8	8
80	Novel method based on ion mobility spectroscopy for the quantification of adulterants in honeys. <i>Food Control</i> , 2020 , 114, 107236	6.2	11
79	Extraction of Anthocyanins and Total Phenolic Compounds from Aġi (<i>Euterpe oleracea</i> Mart.) Using an Experimental Design Methodology. Part 2: Ultrasound-Assisted Extraction. <i>Agronomy</i> , 2020 , 10, 326	3.6	16
78	Optimization of Analytical Ultrasound-Assisted Methods for the Extraction of Total Phenolic Compounds and Anthocyanins from Sloes (<i>Prunus spinosa</i> L.). <i>Agronomy</i> , 2020 , 10, 966	3.6	9
77	Extraction of Anthocyanins and Total Phenolic Compounds from Aġi (<i>Euterpe oleracea</i> Mart.) Using an Experimental Design Methodology. Part 3: Microwave-Assisted Extraction. <i>Agronomy</i> , 2020 , 10, 179	3.6	9
76	Extraction of Anthocyanins and Total Phenolic Compounds from Aġi (<i>Euterpe oleracea</i> Mart.) Using an Experimental Design Methodology. Part 1: Pressurized Liquid Extraction. <i>Agronomy</i> , 2020 , 10, 183	3.6	12

75	Simultaneous extraction and separation of bioactive compounds from apple pomace using pressurized liquids coupled on-line with solid-phase extraction. <i>Food Chemistry</i> , 2020 , 318, 126450	8.5	29
74	Influence of Fruit Ripening on the Total and Individual Capsaicinoids and Capsiate Content in Naga Jolokia Peppers (<i>Capsicum chinense</i> Jacq.). <i>Agronomy</i> , 2020 , 10, 252	3.6	7
73	Changes in Capsiate Content in Four Chili Pepper Genotypes (<i>Capsicum</i> spp.) at Different Ripening Stages. <i>Agronomy</i> , 2020 , 10, 1337	3.6	4
72	Extraction of Flavonoids From Natural Sources Using Modern Techniques. <i>Frontiers in Chemistry</i> , 2020 , 8, 507887	5	67
71	Content of Capsaicinoids and Capsiate in "Filius" Pepper Varieties as Affected by Ripening. <i>Plants</i> , 2020 , 9,	4.5	1
70	Isolation of gallic acid, caffeine and flavonols from black tea by on-line coupling of pressurized liquid extraction with an adsorbent for the production of functional bakery products. <i>LWT - Food Science and Technology</i> , 2020 , 117, 108661	5.4	22
69	Closed-loop spray drying with N ₂ of <i>Moringa oleifera</i> leaf ethanolic extracts: Effects on bioactive compounds and antiradical activity. <i>Drying Technology</i> , 2020 , 1-13	2.6	
68	Discrimination of Myrtle Ecotypes from Different Geographic Areas According to Their Morphological Characteristics and Anthocyanins Composition. <i>Plants</i> , 2019 , 8,	4.5	4
67	A screening method based on Visible-NIR spectroscopy for the identification and quantification of different adulterants in high-quality honey. <i>Talanta</i> , 2019 , 203, 235-241	6.2	28
66	A Screening Method Based on Headspace-Ion Mobility Spectrometry to Identify Adulterated Honey. <i>Sensors</i> , 2019 , 19,	3.8	12
65	Modeling of counter-current multistage extraction of <i>Moringa oleifera</i> leaves using a mechanistic model. <i>Food and Bioproducts Processing</i> , 2019 , 115, 165-174	4.9	7
64	Alternative Ultrasound-Assisted Method for the Extraction of the Bioactive Compounds Present in Myrtle (L.). <i>Molecules</i> , 2019 , 24,	4.8	23
63	Novel method based on ion mobility spectrometry sum spectrum for the characterization of ignitable liquids in fire debris. <i>Talanta</i> , 2019 , 199, 189-194	6.2	10
62	Assessment of Ultrasound Assisted Extraction as an Alternative Method for the Extraction of Anthocyanins and Total Phenolic Compounds from Maqui Berries (<i>Aristotelia chilensis</i> (Mol.) Stuntz). <i>Agronomy</i> , 2019 , 9, 148	3.6	18
61	Progression of the Total and Individual Capsaicinoids Content in the Fruits of Three Different Cultivars of <i>Capsicum chinense</i> Jacq.. <i>Agronomy</i> , 2019 , 9, 141	3.6	9
60	Pressurized liquid extraction of bioactive compounds from grape marc. <i>Journal of Food Engineering</i> , 2019 , 240, 105-113	6	70
59	Quantitation of capsiate and dihydrocapsiate and tentative identification of minor capsinoids in pepper fruits (<i>Capsicum</i> spp.) by HPLC-ESI-MS/MS(QTOF). <i>Food Chemistry</i> , 2019 , 270, 264-272	8.5	19
58	Ultrasound-Assisted Extraction of Two Types of Antioxidant Compounds (TPC and TA) from Black Chokeberry (<i>Aronia melanocarpa</i> L.): Optimization of the Individual and Simultaneous Extraction Methods. <i>Agronomy</i> , 2019 , 9, 456	3.6	12

57	Assessment of Capsaicinoid and Capsinoid Accumulation Patterns during Fruit Development in Three Chili Pepper Genotypes (spp.) Carrying and Alleles Related to Pungency. <i>Journal of Agricultural and Food Chemistry</i> , 2019 , 67, 12219-12227	5.7	11
56	OTP-PRL: an app for occupational risk prevention in policing activities. <i>BMC Public Health</i> , 2019 , 19, 15494.1	4.1	1
55	Optimizing and Comparing Ultrasound- and Microwave-Assisted Extraction Methods Applied to the Extraction of Antioxidant Capsinoids in Peppers. <i>Agronomy</i> , 2019 , 9, 633	3.6	14
54	Extraction of Antioxidants from Blackberry (<i>Rubus ulmifolius</i> L.): Comparison between Ultrasound- and Microwave-Assisted Extraction Techniques. <i>Agronomy</i> , 2019 , 9, 745	3.6	10
53	Escape Classroom: Can You Solve a Crime Using the Analytical Process?. <i>Journal of Chemical Education</i> , 2019 , 96, 267-273	2.4	34
52	FT-IR, Vis spectroscopy, color and multivariate analysis for the control of ageing processes in distinctive Spanish wines. <i>Food Chemistry</i> , 2019 , 277, 6-11	8.5	18
51	Characterization of petroleum-based products in water samples by HS-MS. <i>Fuel</i> , 2018 , 222, 506-512	7.1	5
50	Co-precipitation of anthocyanins of the extract obtained from blackberry residues by pressurized antisolvent process. <i>Journal of Supercritical Fluids</i> , 2018 , 137, 81-92	4.2	21
49	Alternative Extraction Method of Bioactive Compounds from Mulberry (<i>Morus nigra</i> L.) Pulp Using Pressurized-Liquid Extraction. <i>Food Analytical Methods</i> , 2018 , 11, 2384-2395	3.4	16
48	Fast analysis of capsaicinoids in Naga Jolokia extracts (<i>Capsicum chinense</i>) by high-performance liquid chromatography using fused core columns. <i>Food Chemistry</i> , 2018 , 239, 217-224	8.5	29
47	Economic analysis of oleoresin production from malagueta peppers (<i>Capsicum frutescens</i>) by supercritical fluid extraction. <i>Journal of Supercritical Fluids</i> , 2018 , 133, 86-93	4.2	39
46	An Electronic Nose Based Method for the Discrimination of Weathered Petroleum-Derived Products. <i>Sensors</i> , 2018 , 18,	3.8	12
45	Study of the Weathering Process of Gasoline by eNose. <i>Sensors</i> , 2018 , 18,	3.8	6
44	Development of New Analytical Microwave-Assisted Extraction Methods for Bioactive Compounds from Myrtle (L.). <i>Molecules</i> , 2018 , 23,	4.8	21
43	Optimization of Microwave-Assisted Extraction for the Recovery of Bioactive Compounds from the Chilean Superfruit (<i>Aristotelia chilensis</i> (Mol.) Stuntz). <i>Agronomy</i> , 2018 , 8, 240	3.6	23
42	Encapsulation of anthocyanin-rich extract from blackberry residues by spray-drying, freeze-drying and supercritical antisolvent. <i>Powder Technology</i> , 2018 , 340, 553-562	5.2	42
41	Application of Headspace Gas Chromatography-Ion Mobility Spectrometry for the Determination of Ignitable Liquids from Fire Debris. <i>Separations</i> , 2018 , 5, 41	3.1	9
40	Rapid quantification of honey adulteration by visible-near infrared spectroscopy combined with chemometrics. <i>Talanta</i> , 2018 , 188, 288-292	6.2	71

39	Validation of an HS-MS method for direct determination and classification of ignitable liquids. <i>Microchemical Journal</i> , 2017 , 132, 358-364	4.8	10
38	Recovery of anthocyanins from residues of <i>Rubus fruticosus</i> , <i>Vaccinium myrtillus</i> and <i>Eugenia brasiliensis</i> by ultrasound assisted extraction, pressurized liquid extraction and their combination. <i>Food Chemistry</i> , 2017 , 231, 1-10	8.5	85
37	Ontogenetic Variation of Individual and Total Capsaicinoids in Malagueta Peppers (<i>Capsicum frutescens</i>) during Fruit Maturation. <i>Molecules</i> , 2017 , 22,	4.8	18
36	Obtaining anthocyanin-rich extracts from frozen aïi (<i>Euterpe oleracea</i> Mart.) pulp using pressurized liquid extraction. <i>Food Science and Technology</i> , 2017 , 37, 48-54	2	15
35	Extraction of phenolic compounds and anthocyanins from juïra (<i>Euterpe edulis</i> Mart.) residues using pressurized liquids and supercritical fluids. <i>Journal of Supercritical Fluids</i> , 2017 , 119, 9-16	4.2	116
34	Authentication of virgin olive oil by a novel curve resolution approach combined with visible spectroscopy. <i>Food Chemistry</i> , 2017 , 220, 331-336	8.5	30
33	Ultrasound-assisted extraction of bioactive compounds from dedo de moï pepper (<i>Capsicum baccatum</i> L.): Effects on the vegetable matrix and mathematical modeling. <i>Journal of Food Engineering</i> , 2017 , 198, 36-44	6	44
32	Optimization of the ultrasound-assisted extraction of anthocyanins and total phenolic compounds in mulberry (<i>Morus nigra</i>) pulp. <i>Food Chemistry</i> , 2017 , 219, 23-32	8.5	118
31	Tryptophan Levels during Grape Ripening: Effects of Cultural Practices. <i>Molecules</i> , 2017 , 22,	4.8	1
30	Characterization and Differentiation of Petroleum-Derived Products by E-Nose Fingerprints. <i>Sensors</i> , 2017 , 17,	3.8	15
29	Comparative Study of Capsaicinoid Composition in Capsicum Peppers Grown in Brazil. <i>International Journal of Food Properties</i> , 2016 , 19, 1292-1302	3	24
28	Fast Separation of Capsaicinoids from Peppers by Reversed Phase Ultra-Performance Liquid Chromatography: Comparison with Traditional High-Performance Liquid Chromatography Methods. <i>International Journal of Food Properties</i> , 2016 , 19, 984-992	3	13
27	Evolution of Capsaicinoids in Peter Pepper (<i>Capsicum annum</i> var. <i>annuum</i>) During Fruit Ripening. <i>Chemistry and Biodiversity</i> , 2016 , 13, 1068-75	2.5	15
26	Capsaicinoid Contents in Peppers and Pepper-Related Spicy Foods. <i>International Journal of Food Properties</i> , 2016 , 19, 485-493	3	35
25	Effect of ultrasound on the supercritical CO ₂ extraction of bioactive compounds from dedo de moï pepper (<i>Capsicum baccatum</i> L. var. <i>pendulum</i>). <i>Ultrasonics Sonochemistry</i> , 2016 , 31, 284-94	8.9	49
24	Fast analysis of curcuminoids from turmeric (<i>Curcuma longa</i> L.) by high-performance liquid chromatography using a fused-core column. <i>Food Chemistry</i> , 2016 , 200, 167-74	8.5	51
23	Multivariate optimization by statistical methods of ultra high performance liquid chromatography conditions for the separation of 17 capsaicinoids. <i>Analytical Methods</i> , 2016 , 8, 1659-1666	3.2	6
22	Determination of Ignitable Liquids in Fire Debris: Direct Analysis by Electronic Nose. <i>Sensors</i> , 2016 , 16,	3.8	22

21	Encapsulation of pepper oleoresin by supercritical fluid extraction of emulsions. <i>Journal of Supercritical Fluids</i> , 2016 , 112, 37-43	4.2	34
20	Use of multivariate statistical techniques to optimize the separation of 17 capsinoids by ultra performance liquid chromatography using different columns. <i>Talanta</i> , 2015 , 134, 256-263	6.2	13
19	Supercritical carbon dioxide extraction of capsaicinoids from malagueta pepper (<i>Capsicum frutescens</i> L.) assisted by ultrasound. <i>Ultrasonics Sonochemistry</i> , 2015 , 22, 78-88	8.9	106
18	Determination of Melatonin in Rice (<i>Oryza sativa</i>) Grains by Pressurized Liquid Extraction. <i>Journal of Agricultural and Food Chemistry</i> , 2015 , 63, 1107-1115	5.7	20
17	Pressurized liquid extraction of bioactive compounds from blackberry (<i>Rubus fruticosus</i> L.) residues: a comparison with conventional methods. <i>Food Research International</i> , 2015 , 77, 675-683	7	148
16	Evolution of total and individual capsaicinoids in peppers during ripening of the Cayenne pepper plant (<i>Capsicum annum</i> L.). <i>Food Chemistry</i> , 2014 , 153, 200-6	8.5	79
15	Fast analysis of Ecdysone in Brazilian ginseng (<i>Pfaffia glomerata</i>) extracts by high-performance liquid chromatography using a fused-core column. <i>Analytical Methods</i> , 2014 , 6, 2452-2459	3.2	13
14	Extraction of antioxidant compounds from blackberry (<i>Rubus</i> sp.) bagasse using supercritical CO ₂ assisted by ultrasound. <i>Journal of Supercritical Fluids</i> , 2014 , 94, 223-233	4.2	117
13	Extraction of phenolic compounds and anthocyanins from blueberry (<i>Vaccinium myrtillus</i> L.) residues using supercritical CO ₂ and pressurized liquids. <i>Journal of Supercritical Fluids</i> , 2014 , 95, 8-16	4.2	129
12	Supercritical fluid extraction and low pressure extraction of Biquinho pepper (<i>Capsicum chinense</i>). <i>LWT - Food Science and Technology</i> , 2014 , 59, 1239-1246	5.4	34
11	A new solid phase extraction for the determination of anthocyanins in grapes. <i>Molecules</i> , 2014 , 19, 21398-410	4.8	15
10	Stability of anthocyanins from red grape skins under pressurized liquid extraction and ultrasound-assisted extraction conditions. <i>Molecules</i> , 2014 , 19, 21034-43	4.8	18
9	Supercritical carbon dioxide extraction of <i>Capsicum</i> peppers: Global yield and capsaicinoid content. <i>Journal of Supercritical Fluids</i> , 2013 , 81, 210-216	4.2	51
8	Application of Hansch's model to capsaicinoids and capsinoids: a study using the quantitative structure-activity relationship. A novel method for the synthesis of capsinoids. <i>Journal of Agricultural and Food Chemistry</i> , 2010 , 58, 3342-9	5.7	51
7	Rapid Determination of Simple Polyphenols in Grapes by LC Using a Monolithic Column. <i>Chromatographia</i> , 2010 , 72, 417-424	2.1	11
6	Ultrasound-assisted extraction of capsaicinoids from peppers. <i>Talanta</i> , 2008 , 75, 1332-7	6.2	152
5	Fast determination of capsaicinoids from peppers by high-performance liquid chromatography using a reversed phase monolithic column. <i>Food Chemistry</i> , 2008 , 107, 1276-1282	8.5	74
4	Optimization of a New Extraction Technique for Analysis of Verbenone and cis-Verbenol in Pine Seeds. <i>Chromatographia</i> , 2007 , 66, 571-575	2.1	4

3	Pressurized liquid extraction of capsaicinoids from peppers. <i>Journal of Agricultural and Food Chemistry</i> , 2006 , 54, 3231-6	5.7	85
2	Determination of capsaicinoids in peppers by microwave-assisted extraction-high-performance liquid chromatography with fluorescence detection. <i>Analytica Chimica Acta</i> , 2006 , 578, 227-33	6.6	91
1	Composition and antifungal effects of aqueous extracts of <i>Cymbopogon citratus</i> , <i>Laurus nobilis</i> and <i>Santolina chamaecyparissus</i> on the growth of <i>Fusarium oxysporum</i> f. sp. <i>lentis</i> . <i>Archives of Phytopathology and Plant Protection</i> , 1-19	1	0