

# Tatiana Lipina

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11020825/publications.pdf>

Version: 2024-02-01

8  
papers

811  
citations

1307594  
7  
h-index

1720034  
7  
g-index

8  
all docs

8  
docs citations

8  
times ranked

1463  
citing authors

#	ARTICLE	IF	CITATIONS
1	Kainate receptor auxiliary subunit NETO2 is required for normal fear expression and extinction. <i>Neuropsychopharmacology</i> , 2019, 44, 1855-1866.	5.4	13
2	Assessment of Social Interaction Behaviors. <i>Journal of Visualized Experiments</i> , 2011, , .	0.3	306
3	A new model of the disrupted latent inhibition in C57BL/6J mice after bupropion treatment. <i>Psychopharmacology</i> , 2010, 208, 487-498.	3.1	15
4	Neto1 Is a Novel CUB-Domain NMDA Receptor-Interacting Protein Required for Synaptic Plasticity and Learning. <i>PLoS Biology</i> , 2009, 7, e1000041.	5.6	150
5	Mice with reduced NMDA receptor glycine affinity model some of the negative and cognitive symptoms of schizophrenia. <i>Psychopharmacology</i> , 2008, 200, 217-230.	3.1	127
6	The Ampakine CX546 Restores the Prepulse Inhibition and Latent Inhibition Deficits in mGluR5-Deficient Mice. <i>Neuropsychopharmacology</i> , 2007, 32, 745-756.	5.4	52
7	Modulators of the glycine site on NMDA receptors, d-serine and ALX 5407, display similar beneficial effects to clozapine in mouse models of schizophrenia. <i>Psychopharmacology</i> , 2005, 179, 54-67.	3.1	146
8	The genetics of latent inhibition: studies of inbred and mutant mice. , 0, , 225-251.		2