

Bing Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1102435/publications.pdf>

Version: 2024-02-01

21
papers

351
citations

840776

11
h-index

839539

18
g-index

21
all docs

21
docs citations

21
times ranked

331
citing authors

#	ARTICLE	IF	CITATIONS
1	Perennial Flowering Plants Sustain Natural Enemy Populations in Gobi Desert Oases of Southern Xinjiang, China. <i>Insects</i> , 2022, 13, 399.	2.2	1
2	No influence on population dynamics of spider mites in cotton fields of intercropping with walnut, a poor-quality host. <i>Crop Protection</i> , 2021, 148, 105733.	2.1	1
3	The outbreaks of nontarget mirid bugs promote arthropod pest suppression in Bt cotton agroecosystems. <i>Plant Biotechnology Journal</i> , 2020, 18, 322-324.	8.3	18
4	Whorl-stage maize provides a microclimate refuge for predatory ladybeetles. <i>Biological Control</i> , 2020, 142, 104162.	3.0	6
5	Effects of Aphid Density and Plant Taxa on Predatory Ladybeetle Abundance at Field and Landscape Scales. <i>Insects</i> , 2020, 11, 695.	2.2	7
6	A molecular detection approach for a cotton aphid-parasitoid complex in northern China. <i>Scientific Reports</i> , 2019, 9, 15836.	3.3	3
7	Perception of and Behavioral Responses to Host Plant Volatiles for Three <i>Adelphocoris</i> Species. <i>Journal of Chemical Ecology</i> , 2019, 45, 779-788.	1.8	28
8	Floral feeding increases diet breadth in a polyphagous mirid. <i>Journal of Pest Science</i> , 2019, 92, 1089-1100.	3.7	9
9	Non-crop habitats promote the abundance of predatory ladybeetles in maize fields in the agricultural landscape of northern China. <i>Agriculture, Ecosystems and Environment</i> , 2019, 277, 44-52.	5.3	15
10	Mixed effects of landscape complexity and insecticide use on ladybeetle abundance in wheat fields. <i>Pest Management Science</i> , 2019, 75, 1638-1645.	3.4	14
11	Landscape structure alters the abundance and species composition of early-season aphid populations in wheat fields. <i>Agriculture, Ecosystems and Environment</i> , 2019, 269, 167-173.	5.3	24
12	Secondary crops and non-crop habitats within landscapes enhance the abundance and diversity of generalist predators. <i>Agriculture, Ecosystems and Environment</i> , 2018, 258, 30-39.	5.3	23
13	Change in ladybeetle abundance and biological control of wheat aphids over time in agricultural landscape. <i>Agriculture, Ecosystems and Environment</i> , 2018, 255, 102-110.	5.3	20
14	Intercropping With Fruit Trees Increases Population Abundance and Alters Species Composition of Spider Mites on Cotton. <i>Environmental Entomology</i> , 2018, 47, 781-787.	1.4	7
15	Characterization of the natural enemy community attacking cotton aphid in the Bt cotton ecosystem in Northern China. <i>Scientific Reports</i> , 2016, 6, 24273.	3.3	42
16	Landscape diversity enhances parasitism of cotton bollworm (<i>Helicoverpa armigera</i>) eggs by <i>Trichogramma chilonis</i> in cotton. <i>Biological Control</i> , 2016, 93, 15-23.	3.0	31
17	Influence of Landscape Diversity and Composition on the Parasitism of Cotton Bollworm Eggs in Maize. <i>PLoS ONE</i> , 2016, 11, e0149476.	2.5	12
18	Effects of temperature and humidity on immature development of <i>Lygus pratensis</i> (L.) (Hemiptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	8.9	11

#	ARTICLE	IF	CITATIONS
19	Lethal and sublethal effects of cycloxyprid, a novel cis-nitromethylene neonicotinoid insecticide, on the mirid bug <i>Apolygus lucorum</i> . <i>Journal of Pest Science</i> , 2014, 87, 731-738.	3.7	68
20	Performance of Three <i>Adelphocoris</i> spp. (Hemiptera: Miridae) on Flowering and Non-flowering Cotton and Alfalfa. <i>Journal of Integrative Agriculture</i> , 2014, 13, 1727-1735.	3.5	5
21	Life Table Parameters of Three Mirid Bug (<i>Adelphocoris</i>) Species (Hemiptera: Miridae) under Contrasted Relative Humidity Regimes. <i>PLoS ONE</i> , 2014, 9, e115878.	2.5	6