

Hans-Peter Zenner

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11045834/publications.pdf>

Version: 2024-02-01

48
papers

2,086
citations

218677

26
h-index

243625

44
g-index

49
all docs

49
docs citations

49
times ranked

1490
citing authors

#	ARTICLE	IF	CITATIONS
1	Implantability of endaurally insertable active vibratory middle-ear implants – an anatomical study. <i>Acta Oto-Laryngologica</i> , 2019, 139, 561-566.	0.9	0
2	A multidisciplinary systematic review of the treatment for chronic idiopathic tinnitus. <i>European Archives of Oto-Rhino-Laryngology</i> , 2017, 274, 2079-2091.	1.6	117
3	Standardized tinnitus-specific individual cognitive-behavioral therapy: A controlled outcome study with 286 tinnitus patients. <i>Hearing Research</i> , 2013, 298, 117-125.	2.0	20
4	Concept and evaluation of an endaurally insertable middle-ear implant. <i>Medical Engineering and Physics</i> , 2013, 35, 532-536.	1.7	16
5	Conditions for Highly Efficient and Reproducible Round-Window Stimulation in Humans. <i>Audiology and Neuro-Otology</i> , 2012, 17, 133-138.	1.3	29
6	Controlled round-window stimulation in human temporal bones yielding reproducible and functionally relevant stapedial responses. <i>Hearing Research</i> , 2011, 282, 272-282.	2.0	24
7	Townes-Brocks Syndrome. , 2009, , 2092-2094.		0
8	Randomized, double blind, placebo controlled trial on the safety and efficacy of continuous intratympanic dexamethasone delivered via a round window catheter for severe to profound sudden idiopathic sensorineural hearing loss after failure of systemic therapy. <i>Laryngoscope</i> , 2009, 119, 359-369.	2.0	143
9	Response to <i>Randomized, Double Blind, Placebo Controlled Trial on the Safety and Efficacy of Continuous Intratympanic Dexamethasone Delivered Via a Round Window Catheter for Severe to Profound Sudden Idiopathic Sensorineural Hearing Loss After Failure of Systemic Therapy</i> (<i>Laryngoscope</i> 119:359–369, 2009). <i>Laryngoscope</i> , 2009, 119, 2481-2482.	2.0	1
10	Speech-in-noise intelligibility does not correlate with efferent olivocochlear reflex in humans with normal hearing. <i>Acta Oto-Laryngologica</i> , 2008, 128, 53-60.	0.9	53
11	Cochlear Pharmacokinetics with Local Inner Ear Drug Delivery Using a Three-Dimensional Finite-Element Computer Model. <i>Audiology and Neuro-Otology</i> , 2007, 12, 37-48.	1.3	55
12	Hyposmotic stimulation-induced nitric oxide production in outer hair cells of the guinea pig cochlea. <i>Hearing Research</i> , 2007, 227, 59-70.	2.0	8
13	Hyposmotic stimulation-induced nitric oxide production in outer hair cells of the guinea pig cochlea. <i>Hearing Research</i> , 2007, 230, 93-104.	2.0	17
14	Technical Note on Microcatheter Implantation for Local Inner Ear Drug Delivery. <i>Otology and Neurotology</i> , 2006, 27, 912-917.	1.3	47
15	Outcomes research analysis of continuous intratympanic glucocorticoid delivery in patients with acute severe to profound hearing loss: Basis for planning randomized controlled trials. <i>Acta Oto-Laryngologica</i> , 2005, 125, 830-839.	0.9	64
16	Cognitive tinnitus sensitization: Behavioral and neurophysiological aspects of tinnitus centralization. <i>Acta Oto-Laryngologica</i> , 2004, 124, 436-439.	0.9	31
17	Aquaporin-mediated fluid regulation in the inner ear. <i>Cellular and Molecular Neurobiology</i> , 2003, 23, 315-329.	3.3	56
18	Thyroid hormone is a critical determinant for the regulation of the cochlear motor protein prestin. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2002, 99, 2901-2906.	7.1	107

#	ARTICLE	IF	CITATIONS
19	Thyroid Hormone-deficient Period Prior to the Onset of Hearing Is Associated with Reduced Levels of β 2-Tectorin Protein in the Tectorial Membrane. <i>Journal of Biological Chemistry</i> , 2001, 276, 39046-39052.	3.4	63
20	Remarks about the depth resolution of heterodyne interferometers in cochlear investigations. <i>Journal of the Acoustical Society of America</i> , 2001, 110, 1725-1728.	1.1	15
21	Molecular characterization of anion exchangers in the cochlea. <i>Molecular and Cellular Biochemistry</i> , 2000, 205, 25-37.	3.1	9
22	Three-Dimensional Motion of the Organ of Corti. <i>Biophysical Journal</i> , 2000, 78, 2285-2297.	0.5	64
23	Evidence for active, nonlinear, negative feedback in the vibration response of the apical region of the in-vivo guinea-pig cochlea. <i>Hearing Research</i> , 2000, 142, 159-183.	2.0	86
24	Characteristics of the travelling wave in the low-frequency region of a temporal-bone preparation of the guinea-pig cochlea. <i>Hearing Research</i> , 2000, 142, 184-202.	2.0	24
25	MECHANISMS OF COUPLING THE ELECTROMECHANICAL FORCES OF THE OUTER HAIR CELLS INTO THE COCHLEAR PARTITION. , 2000, , .		2
26	A Changing Pattern of Brain-Derived Neurotrophic Factor Expression Correlates with the Rearrangement of Fibers during Cochlear Development of Rats and Mice. <i>Journal of Neuroscience</i> , 1999, 19, 3033-3042.	3.6	89
27	Comparative study of visual, auditory, and olfactory function in Usher syndrome. <i>Graefe's Archive for Clinical and Experimental Ophthalmology</i> , 1999, 237, 301-307.	1.9	33
28	Distinct thyroid hormone-dependent expression of trkB and p75NGFR in nonneuronal cells during the critical TH-dependent period of the cochlea. <i>Journal of Neurobiology</i> , 1999, 38, 338-356.	3.6	57
29	Differential expression of trkB.T1 and trkB.T2, truncated trkC, and p75NGFR in the cochlea prior to hearing function. <i>Journal of Comparative Neurology</i> , 1999, 414, 33-49.	1.6	53
30	A second Middle Eastern kindred with autosomal recessive non-syndromic hearing loss segregates DFNB9. <i>European Journal of Human Genetics</i> , 1998, 6, 341-344.	2.8	14
31	The effect of anti-diuretic hormone on the endolymphatic sac of the inner ear. <i>Pflugers Archiv European Journal of Physiology</i> , 1998, 436, 970-975.	2.8	111
32	Low-coherence fibre heterodyne interferometer for both dc and high-frequency vibration measurements in the inner ear. <i>Journal of Modern Optics</i> , 1998, 45, 765-775.	1.3	11
33	Suppression of distortion product otoacoustic emissions (DPOAE) near $2f_1 \sim f_2$ removes DP-gram fine structure—Evidence for a secondary generator. <i>Journal of the Acoustical Society of America</i> , 1998, 103, 1527-1531.	1.1	117
34	Evidence for Opening of Hair-Cell Transducer Channels after Tip-Link Loss. <i>Journal of Neuroscience</i> , 1998, 18, 6748-6756.	3.6	65
35	<title>Laser interferometric measurement of the micromechanics of the inner ear</title>. , 1995, , .		1
36	Caloric Evoked Motile Responses of Mammalian Vestibular Sensory Cells. <i>Acta Oto-Laryngologica</i> , 1995, 115, 484-487.	0.9	5

#	ARTICLE	IF	CITATIONS
37	Synaptophysin and Gap-43 proteins in efferent fibers of the inner ear during postnatal development. <i>Developmental Brain Research</i> , 1995, 89, 73-86.	1.7	52
38	Sound-induced displacement responses in the plane of the organ of Corti in the isolated guinea-pig cochlea. <i>Hearing Research</i> , 1995, 83, 142-150.	2.0	20
39	Acute gentamicin ototoxicity in cochlear outer hair cells of the guinea pig. <i>Brain Research</i> , 1994, 636, 153-156.	2.2	21
40	Frequency response of mature guinea-pig outer hair cells to stereociliary displacement. <i>Hearing Research</i> , 1994, 77, 116-124.	2.0	15
41	High frequency radial movements of the reticular lamina induced by outer hair cell motility. <i>Hearing Research</i> , 1992, 60, 236-246.	2.0	43
42	Motile Responses of Vestibular Hair Cells Following Caloric, Electrical or Chemical Stimuli. <i>Acta Oto-Laryngologica</i> , 1991, 111, 291-297.	0.9	25
43	Active radial and transverse motile responses of outer hair cells in the organ of corti. <i>Hearing Research</i> , 1990, 43, 219-230.	2.0	69
44	The cell potential of isolated inner hair cells in vitro. <i>Hearing Research</i> , 1990, 45, 87-93.	2.0	18
45	Active movements of the cuticular plate induce sensory hair motion in mammalian outer hair cells. <i>Hearing Research</i> , 1988, 34, 233-239.	2.0	73
46	Immunological Aspects of Tonsils and of Tonsillitis. <i>Acta Oto-Laryngologica</i> , 1988, 105, 70-74.	0.9	1
47	Outer Hair Cells as Fast and Slow Cochlear Amplifiers with a Bidirectional Transduction Cycle. <i>Acta Oto-Laryngologica</i> , 1988, 105, 457-462.	0.9	50
48	Evidence that phosphoinositides mediate motility in cochlear outer hair cells. <i>Hearing Research</i> , 1987, 31, 155-159.	2.0	89