

Jing Liu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1105156/publications.pdf>

Version: 2024-02-01

17
papers

263
citations

933447

10
h-index

940533

16
g-index

17
all docs

17
docs citations

17
times ranked

208
citing authors

#	ARTICLE	IF	CITATIONS
1	Deletion of Toxoplasma Rhopty Protein 38 (Pru ^{rop38}) as a Vaccine Candidate for Toxoplasmosis in a Murine Model. <i>Biomedicines</i> , 2022, 10, 1336.	3.2	2
2	Function of Neospora caninum dense granule protein 7 in innate immunity in mice. <i>Parasitology Research</i> , 2021, 120, 197-207.	1.6	4
3	Microneme Protein 6 Is Involved in Invasion and Egress by Neospora caninum. <i>Pathogens</i> , 2021, 10, 201.	2.8	1
4	NcPuf1 Is a Key Virulence Factor in Neospora caninum. <i>Pathogens</i> , 2020, 9, 1019.	2.8	1
5	Toxoplasma gondii metacaspase 2 is an important factor that influences bradyzoite formation in the Pru strain. <i>Parasitology Research</i> , 2020, 119, 2287-2298.	1.6	4
6	Functional characterization of acyl-CoA binding protein in Neospora caninum. <i>Parasites and Vectors</i> , 2020, 13, 85.	2.5	2
7	Synergistic roles of acyl-CoA binding protein (ACBP1) and sterol carrier protein 2 (SCP2) in <i>Toxoplasma</i> lipid metabolism. <i>Cellular Microbiology</i> , 2019, 21, e12970.	2.1	16
8	NcGRA17 is an important regulator of parasitophorous vacuole morphology and pathogenicity of Neospora caninum. <i>Veterinary Parasitology</i> , 2018, 264, 26-34.	1.8	34
9	A new microneme protein of Neospora caninum , NcMIC8 is involved in host cell invasion. <i>Experimental Parasitology</i> , 2017, 175, 21-27.	1.2	9
10	Neospora caninum ROP16 play an important role in the pathogenicity by phosphorylating host cell STAT3. <i>Veterinary Parasitology</i> , 2017, 243, 135-147.	1.8	26
11	Rhopty protein 5 (ROP5) Is a Key Virulence Factor in Neospora caninum. <i>Frontiers in Microbiology</i> , 2017, 8, 370.	3.5	25
12	MIC3, a novel cross-protective antigen expressed in Toxoplasma gondii and Neospora caninum. <i>Parasitology Research</i> , 2015, 114, 3791-3799.	1.6	19
13	Identification and characterization of a microneme protein (NcMIC6) in Neospora caninum. <i>Parasitology Research</i> , 2015, 114, 2893-2902.	1.6	31
14	The Apoptotic Role of Metacaspase in Toxoplasma gondii. <i>Frontiers in Microbiology</i> , 2015, 6, 1560.	3.5	27
15	A Nuclear Factor of High Mobility Group Box Protein in Toxoplasma gondii. <i>PLoS ONE</i> , 2014, 9, e111993.	2.5	24
16	ROP18 Is a Key Factor Responsible for Virulence Difference between Toxoplasma gondii and Neospora caninum. <i>PLoS ONE</i> , 2014, 9, e99744.	2.5	27
17	GRA 14, a novel dense granule protein from <i>Neospora caninum</i> . <i>Acta Biochimica Et Biophysica Sinica</i> , 2013, 45, 607-609.	2.0	11