Hitesh Vaidya

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11087118/publications.pdf Version: 2024-02-01



Ηίτεςη Νλίσνλ

#	Article	IF	CITATIONS
1	Antiâ€diabetic Activity of Swertiamarin is due to an Active Metabolite, Gentianine, that Upregulates PPARâ€Î³ Gene Expression in 3T3‣1 cells. Phytotherapy Research, 2013, 27, 624-627.	5.8	69
2	Antihyperlipidaemic activity of swertiamarin, a secoiridoid glycoside in poloxamer-407-induced hyperlipidaemic rats. Journal of Natural Medicines, 2009, 63, 437-442.	2.3	65
3	Swertiamarin: A lead from Enicostemma littorale Blume. for anti-hyperlipidaemic effect. European Journal of Pharmacology, 2009, 617, 108-112.	3.5	63
4	Beneficial Effects of Swertiamarin on Dyslipidaemia in Streptozotocinâ€induced Type 2 Diabetic Rats. Phytotherapy Research, 2012, 26, 1259-1261.	5.8	33
5	Sea cucumber and blue mussel: new sources of phospholipid enriched omega-3 fatty acids with a potential role in 3T3-L1 adipocyte metabolism. Food and Function, 2014, 5, 3287-3295.	4.6	27
6	Arachidonic acid has a dominant effect to regulate lipogenic genes in 3T3-L1 adipocytes compared to omega-3 fatty acids. Food and Nutrition Research, 2015, 59, 25866.	2.6	12
7	Recent advances in the treatment of thromboembolic diseases: Venous thromboembolism. Medicinal Research Reviews, 2007, 27, 891-914.	10.5	9
8	Shrimp Oil Extracted from Shrimp Processing By-Product Is a Rich Source of Omega-3 Fatty Acids and Astaxanthin-Esters, and Reveals Potential Anti-Adipogenic Effects in 3T3-L1 Adipocytes. Marine Drugs, 2021, 19, 259.	4.6	5
9	Treatment of preadipocytes with fish oil, mixed oil, or soybean oil-based lipid emulsions have differential effects on the regulation of lipogenic and lipolytic genes in mature 3T3-L1 adipocytes. Prostaglandins Leukotrienes and Essential Fatty Acids, 2022, 177, 102396.	2.2	1
10	Marine Derived Bioactives to Combat Obesity: Potential Mechanisms of Action. , 2021, , 373-388.		0