List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11105426/publications.pdf

Version: 2024-02-01

172207 149479 3,433 64 29 56 citations h-index g-index papers 64 64 64 4683 docs citations citing authors all docs times ranked

#	Article	IF	CITATIONS
1	Acceptability, benefits, and challenges of video consulting: a qualitative study in primary care. British Journal of General Practice, 2019, 69, e586-e594.	0.7	353
2	Cognitive behavioural therapy as an adjunct to pharmacotherapy for primary care based patients with treatment resistant depression: results of the CoBalT randomised controlled trial. Lancet, The, 2013, 381, 375-384.	6.3	278
3	Comparing the content and quality of video, telephone, and face-to-face consultations: a non-randomised, quasi-experimental, exploratory study in UK primary care. British Journal of General Practice, 2019, 69, e595-e604.	0.7	253
4	Facilitated physical activity as a treatment for depressed adults: randomised controlled trial. BMJ, The, 2012, 344, e2758-e2758.	3.0	163
5	Reliability of patient responses in pay for performance schemes: analysis of national General Practitioner Patient Survey data in England. BMJ: British Medical Journal, 2009, 339, b3851-b3851.	2.4	129
6	The GP Patient Survey for use in primary care in the National Health Service in the UK – development and psychometric characteristics. BMC Family Practice, 2009, 10, 57.	2.9	124
7	Levels and predictors of exercise referral scheme uptake and adherence: a systematic review. Journal of Epidemiology and Community Health, 2012, 66, 737-744.	2.0	120
8	Prevalence of treatment-resistant depression in primary care: cross-sectional data. British Journal of General Practice, 2013, 63, e852-e858.	0.7	115
9	Understanding help seeking behaviour among male offenders: qualitative interview study. BMJ: British Medical Journal, 2007, 334, 303.	2.4	114
10	Long-term effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: follow-up of the CoBalT randomised controlled trial. Lancet Psychiatry,the, 2016, 3, 137-144.	3.7	103
11	Antibiotic prescription strategies for acute sore throat: a prospective observational cohort study. Lancet Infectious Diseases, The, 2014, 14, 213-219.	4.6	100
12	Patients' views of physical activity as treatment for depression: a qualitative study. British Journal of General Practice, 2011, 61, e149-e156.	0.7	88
13	Predictors of suppurative complications for acute sore throat in primary care: prospective clinical cohort study. BMJ, The, 2013, 347, f6867-f6867.	3.0	84
14	Setting standards based on patients' views on access and continuity: secondary analysis of data from the general practice assessment survey. BMJ: British Medical Journal, 2003, 326, 258-258.	2.4	79
15	Alternatives to the face-to-face consultation in general practice: focused ethnographic case study. British Journal of General Practice, 2018, 68, e293-e300.	0.7	74
16	Pharmacists' perceptions of their emerging general practice roles in UK primary care: a qualitative interview study. British Journal of General Practice, 2017, 67, e650-e658.	0.7	70
17	Clinical effectiveness and cost-effectiveness of cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: the CoBalT randomised controlled trial. Health Technology Assessment, 2014, 18, 1-167, vii-viii.	1.3	69
18	Investigating the relationship between consultation length and patient experience: a cross-sectional study in primary care. British Journal of General Practice, 2016, 66, e896-e903.	0.7	64

#	Article	IF	CITATIONS
19	The potential of alternatives to face-to-face consultation in general practice, and the impact on different patient groups: a mixed-methods case study. Health Services and Delivery Research, 2018, 6, 1-200.	1.4	59
20	Assessing communication quality of consultations in primary care: initial reliability of the Global Consultation Rating Scale, based on the Calgary-Cambridge Guide to the Medical Interview. BMJ Open, 2014, 4, e004339.	0.8	56
21	How can Health Care Organizations be Reliably Compared?. Medical Care, 2011, 49, 724-733.	1.1	55
22	Do English patients want continuity of care, and do they receive it?. British Journal of General Practice, 2012, 62, e567-e575.	0.7	51
23	Bringing light into darkness: A multiple baseline mixed methods case series evaluation of Augmented Depression Therapy (ADepT). Behaviour Research and Therapy, 2019, 120, 103418.	1.6	47
24	Mirtazapine added to SSRIs or SNRIs for treatment resistant depression in primary care: phase III randomised placebo controlled trial (MIR). BMJ: British Medical Journal, 2018, 363, k4218.	2.4	44
25	The Consolations of Going Back to Prison: What †Revolving Door†Prisoners Think of Their Prospects. Journal of Offender Rehabilitation, 2009, 48, 439-461.	0.5	43
26	Supervised Machine Learning Algorithms Can Classify Open-Text Feedback of Doctor Performance With Human-Level Accuracy. Journal of Medical Internet Research, 2017, 19, e65.	2.1	41
27	Variations in GP–patient communication by ethnicity, age, and gender: evidence from a national primary care patient survey. British Journal of General Practice, 2016, 66, e47-e52.	0.7	39
28	Telephone triage systems in UK general practice: analysis of consultation duration during the index day in a pragmatic randomised controlled trial. British Journal of General Practice, 2016, 66, e214-e218.	0.7	36
29	Understanding quit decisions in primary care: a qualitative study of older GPs: TableÂ1. BMJ Open, 2016, 6, e010592.	0.8	33
30	Cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment resistant depression in primary care: The CoBalT randomised controlled trial protocol. Contemporary Clinical Trials, 2012, 33, 312-319.	0.8	32
31	The Evaluation of Physicians' Communication Skills From Multiple Perspectives. Annals of Family Medicine, 2018, 16, 330-337.	0.9	32
32	Do difficulties in accessing in-hours primary care predict higher use of out-of-hours GP services? Evidence from an English National Patient Survey. Emergency Medicine Journal, 2015, 32, 373-378.	0.4	28
33	Cost-effectiveness of cognitive–behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care: economic evaluation of the CoBalT Trial. British Journal of Psychiatry, 2014, 204, 69-76.	1.7	27
34	Doctors' engagements with patient experience surveys in primary and secondary care: a qualitative study. Health Expectations, 2017, 20, 385-394.	1.1	26
35	Parental Autonomy. Journal of Applied Philosophy, 1988, 5, 183-196.	0.7	24
36	General Practitioners' beliefs about physical activity for managing depression in primary care. Mental Health and Physical Activity, 2012, 5, 13-19.	0.9	24

#	Article	IF	Citations
37	Management of treatment-resistant depression in primary care: a mixed-methods study. British Journal of General Practice, 2018, 68, e673-e681.	0.7	23
38	Improving patient experience in primary care: a multimethod programme of research on the measurement and improvement of patient experience. Programme Grants for Applied Research, 2017, 5, 1-452.	0.4	23
39	Investigating the meaning of â€~good' or â€~very good' patient evaluations of care in English general practice: a mixed methods study. BMJ Open, 2017, 7, e014718.	0.8	21
40	Combining mirtazapine with SSRIs or SNRIs for treatment-resistant depression: the MIR RCT. Health Technology Assessment, 2018, 22, 1-136.	1.3	21
41	Capturing patient experience: a qualitative study of implementing real-time feedback in primary care. British Journal of General Practice, 2016, 66, e786-e793.	0.7	20
42	Users' reports and evaluations of out-of-hours health care and the UK national quality requirements: a cross sectional study. British Journal of General Practice, 2009, 59, e8-e15.	0.7	18
43	Evaluating Augmented Depression Therapy (ADepT): study protocol for a pilot randomised controlled trial. Pilot and Feasibility Studies, 2019, 5, 63.	0.5	18
44	Physical activity as a treatment for depression: the TREAD randomised trial protocol. Trials, $2010,11,105.$	0.7	16
45	Moderators of response to cognitive behavioural therapy as an adjunct to pharmacotherapy for treatment-resistant depression in primary care. Journal of Affective Disorders, 2015, 174, 272-280.	2.0	16
46	Understanding negative feedback from South Asian patients: an experimental vignette study. BMJ Open, 2016, 6, e011256.	0.8	16
47	Why do GPs exclude patients from participating in research? An exploration of adherence to and divergence from trial criteria. Family Practice, 2014, 31, 364-370.	0.8	15
48	Locke on Qualities. Canadian Journal of Philosophy, 1980, 10, 567-585.	0.6	14
49	Patients' use and views of realâ€ŧime feedback technology in general practice. Health Expectations, 2017, 20, 419-433.	1.1	14
50	Rating Communication in GP Consultations: The Association Between Ratings Made by Patients and Trained Clinical Raters. Medical Care Research and Review, 2018, 75, 201-218.	1.0	14
51	Symptom response to antibiotic prescribing strategies in acute sore throat in adults: the DESCARTE prospective cohort study in UK general practice. British Journal of General Practice, 2017, 67, e634-e642.	0.7	13
52	Receptionists' role in new approaches to consultations in primary care: a focused ethnographic study. British Journal of General Practice, 2018, 68, e478-e486.	0.7	13
53	Pathways to suicide attempts among male offenders: the role of agency. Sociology of Health and Illness, 2015, 37, 936-951.	1.1	12
54	Individuals' Long Term Use of Cognitive Behavioural Skills to Manage their Depression: A Qualitative Study. Behavioural and Cognitive Psychotherapy, 2017, 45, 46-57.	0.9	11

#	Article	IF	CITATIONS
55	Patients' experiences of participating in a large-scale trial of cognitive behavioural therapy for depression: a mixed methods study. Family Practice, 2013, 30, 705-711.	0.8	10
56	Mirtazapine added to selective serotonin reuptake inhibitors for treatment-resistant depression in primary care (MIR trial): study protocol for a randomised controlled trial. Trials, 2016, 17, 66.	0.7	10
57	Randomised controlled trial of an augmented exercise referral scheme using web-based behavioural support for inactive adults with chronic health conditions: the e-coachER trial. British Journal of Sports Medicine, 2021, 55, 444-450.	3.1	10
58	Adding web-based behavioural support to exercise referral schemes for inactive adults with chronic health conditions: the e-coachER RCT. Health Technology Assessment, 2020, 24, 1-106.	1.3	10
59	Influence of the duration of penicillin prescriptions on outcomes for acute sore throat in adults: the DESCARTE prospective cohort study in UK general practice. British Journal of General Practice, 2017, 67, e623-e633.	0.7	7
60	Telephone first consultations in primary care. BMJ: British Medical Journal, 2017, 358, j4345.	2.4	5
61	Digital Facilitation to Support Patient Access to Web-Based Primary Care Services: Scoping Literature Review. Journal of Medical Internet Research, 2022, 24, e33911.	2.1	5
62	Establishing the validity of English GP Patient Survey items evaluating out-of-hours care. BMJ Quality and Safety, 2016, 25, 842-850.	1.8	1
63	SPINOZA'S THEORY OF PERFECTION AND GOODNESS. Southern Journal of Philosophy, 1980, 18, 259-274.	0.4	O
64	Reply to Bennett. Canadian Journal of Philosophy, 1982, 12, 757-761.	0.6	O