

Cynthia Behling

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/11166791/cynthia-behling-publications-by-year.pdf>

Version: 2024-04-11

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

50 papers	10,761 citations	28 h-index	58 g-index
58 ext. papers	13,020 ext. citations	8.4 avg, IF	5.3 L-index

#	Paper	IF	Citations
50	Two-step strategy, FIB-4 followed by magnetic resonance elastography, for detecting advanced fibrosis in NAFLD.. <i>Clinical Gastroenterology and Hepatology</i> , 2022 ,	6.9	1
49	How to Interpret EUS-FNA Cytology 2022 , 135-145		
48	Validation of the accuracy of the FAST _{score} for detecting patients with at-risk nonalcoholic steatohepatitis (NASH) in a North American cohort and comparison to other non-invasive algorithms.. <i>PLoS ONE</i> , 2022 , 17, e0266859	3.7	0
47	Clinical Utility of Change in Nonalcoholic Fatty Liver Disease Activity Score and Change in Fibrosis in NAFLD. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , 19, 2673-2674.e3	6.9	2
46	Diagnostic accuracy of two-dimensional shear wave elastography and transient elastography in nonalcoholic fatty liver disease. <i>Therapeutic Advances in Gastroenterology</i> , 2021 , 14, 17562848211050438	4.7	4
45	Prospective Study of Outcomes in Adults with Nonalcoholic Fatty Liver Disease. <i>New England Journal of Medicine</i> , 2021 , 385, 1559-1569	59.2	33
44	MRE combined with FIB-4 (MEFIB) index in detection of candidates for pharmacological treatment of NASH-related fibrosis. <i>Gut</i> , 2021 , 70, 1946-1953	19.2	18
43	Clinical utility of 30% relative decline in MRI-PDFF in predicting fibrosis regression in non-alcoholic fatty liver disease. <i>Gut</i> , 2021 ,	19.2	10
42	Alanine Aminotransferase and Gamma-Glutamyl Transpeptidase Predict Histologic Improvement in Pediatric Nonalcoholic Steatohepatitis. <i>Hepatology</i> , 2021 , 73, 937-951	11.2	9
41	MRE plus FIB-4 (MEFIB) versus FAST in detection of candidates for pharmacological treatment of NASH-related fibrosis. <i>Hepatology</i> , 2021 ,	11.2	6
40	Multicenter Validation of Association Between Decline in MRI-PDFF and Histologic Response in NASH. <i>Hepatology</i> , 2020 , 72, 1219-1229	11.2	39
39	Progression of Fatty Liver Disease in Children Receiving Standard of Care Lifestyle Advice. <i>Gastroenterology</i> , 2020 , 159, 1731-1751.e10	13.3	17
38	Sex Hormone Relations to Histologic Severity of Pediatric Nonalcoholic Fatty Liver Disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2020 , 105,	5.6	10
37	Clinical Utility of an Increase in Magnetic Resonance Elastography in Predicting Fibrosis Progression in Nonalcoholic Fatty Liver Disease. <i>Hepatology</i> , 2020 , 71, 849-860	11.2	39
36	Microbiome Signatures Associated With Steatohepatitis and Moderate to Severe Fibrosis in Children With Nonalcoholic Fatty Liver Disease. <i>Gastroenterology</i> , 2019 , 157, 1109-1122	13.3	92
35	Association of Histologic Disease Activity With Progression of Nonalcoholic Fatty Liver Disease. <i>JAMA Network Open</i> , 2019 , 2, e1912565	10.4	108
34	Diagnostic Accuracy of Noninvasive Fibrosis Models to Detect Change in Fibrosis Stage. <i>Clinical Gastroenterology and Hepatology</i> , 2019 , 17, 1877-1885.e5	6.9	63

33	A Pilot Genome-Wide Analysis Study Identifies Loci Associated With Response to Obeticholic Acid in Patients With NASH. <i>Hepatology Communications</i> , 2019 , 3, 1571-1584	6	8
32	Hepatic R2* is more strongly associated with proton density fat fraction than histologic liver iron scores in patients with nonalcoholic fatty liver disease. <i>Journal of Magnetic Resonance Imaging</i> , 2019 , 49, 1456-1466	5.6	13
31	Magnetic Resonance Imaging Proton Density Fat Fraction Associates With Progression of Fibrosis in Patients With Nonalcoholic Fatty Liver Disease. <i>Gastroenterology</i> , 2018 , 155, 307-310.e2	13.3	61
30	In Children With Nonalcoholic Fatty Liver Disease, Zone 1 Steatosis Is Associated With Advanced Fibrosis. <i>Clinical Gastroenterology and Hepatology</i> , 2018 , 16, 438-446.e1	6.9	37
29	Exome sequencing of an adolescent with nonalcoholic fatty liver disease identifies a clinically actionable case of Wilson disease. <i>Journal of Physical Education and Sports Management</i> , 2018 , 4,	2.8	2
28	Hepatic Nuclear Receptor Expression Associates with Features of Histology in Pediatric Nonalcoholic Fatty Liver Disease. <i>Hepatology Communications</i> , 2018 , 2, 1213-1226	6	5
27	Liver histology and diffusion-weighted MRI in children with nonalcoholic fatty liver disease: A MAGNET study. <i>Journal of Magnetic Resonance Imaging</i> , 2017 , 46, 1149-1158	5.6	21
26	Magnetic resonance elastography measured shear stiffness as a biomarker of fibrosis in pediatric nonalcoholic fatty liver disease. <i>Hepatology</i> , 2017 , 66, 1474-1485	11.2	77
25	Low and High Birth Weights Are Risk Factors for Nonalcoholic Fatty Liver Disease in Children. <i>Journal of Pediatrics</i> , 2017 , 187, 141-146.e1	3.6	64
24	In Children With Nonalcoholic Fatty Liver Disease, Cysteamine Bitartrate Delayed Release Improves Liver Enzymes but Does Not Reduce Disease Activity Scores. <i>Gastroenterology</i> , 2016 , 151, 1141-1154.e9	13.3	73
23	EUS FNA cytology: material preparation and interpretation 2016 , 82-87		
22	Prevalence of Prediabetes and Type 2 Diabetes in Children With Nonalcoholic Fatty Liver Disease. <i>JAMA Pediatrics</i> , 2016 , 170, e161971	8.3	123
21	Reproducibility of histological assessments of disease activity in UC. <i>Gut</i> , 2015 , 64, 1765-73	19.2	49
20	Associations between histologic features of nonalcoholic fatty liver disease (NAFLD) and quantitative diffusion-weighted MRI measurements in adults. <i>Journal of Magnetic Resonance Imaging</i> , 2015 , 41, 1629-38	5.6	48
19	Association Between Quantity of Liver Fat and Cardiovascular Risk in Patients With Nonalcoholic Fatty Liver Disease Independent of Nonalcoholic Steatohepatitis. <i>Clinical Gastroenterology and Hepatology</i> , 2015 , 13, 1513-20.e1	6.9	69
18	Magnetic resonance imaging and liver histology as biomarkers of hepatic steatosis in children with nonalcoholic fatty liver disease. <i>Hepatology</i> , 2015 , 61, 1887-95	11.2	107
17	Evidence and recommendations for imaging liver fat in children, based on systematic review. <i>Clinical Gastroenterology and Hepatology</i> , 2014 , 12, 765-73	6.9	78
16	Longitudinal assessment of high blood pressure in children with nonalcoholic fatty liver disease. <i>PLoS ONE</i> , 2014 , 9, e112569	3.7	59

15	Magnetic resonance elastography predicts advanced fibrosis in patients with nonalcoholic fatty liver disease: a prospective study. <i>Hepatology</i> , 2014 , 60, 1920-8	11.2	304
14	Histologic evaluation of ulcerative colitis: a systematic review of disease activity indices. <i>Inflammatory Bowel Diseases</i> , 2014 , 20, 564-75	4.5	78
13	Clinical and histological determinants of nonalcoholic steatohepatitis and advanced fibrosis in elderly patients. <i>Hepatology</i> , 2013 , 58, 1644-54	11.2	109
12	Misuse of scoring systems. <i>Hepatology</i> , 2011 , 54, 369-70; author reply 370-1	11.2	25
11	EUS-guided FNA diagnostic yield of malignancy in solid pancreatic masses: a benchmark for quality performance measurement. <i>Gastrointestinal Endoscopy</i> , 2007 , 66, 277-82	5.2	137
10	Prevalence of fatty liver in children and adolescents. <i>Pediatrics</i> , 2006 , 118, 1388-93	7.4	1019
9	Pediatric nonalcoholic fatty liver disease: a critical appraisal of current data and implications for future research. <i>Journal of Pediatric Gastroenterology and Nutrition</i> , 2006 , 43, 413-27	2.8	174
8	Design and validation of a histological scoring system for nonalcoholic fatty liver disease. <i>Hepatology</i> , 2005 , 41, 1313-21	11.2	6503
7	Histopathology of pediatric nonalcoholic fatty liver disease. <i>Hepatology</i> , 2005 , 42, 641-9	11.2	549
6	Obesity, insulin resistance, and other clinicopathological correlates of pediatric nonalcoholic fatty liver disease. <i>Journal of Pediatrics</i> , 2003 , 143, 500-5	3.6	371
5	Parathyroid hormone-related protein as a novel tumor marker in pancreatic adenocarcinoma. <i>Pancreas</i> , 2002 , 24, 284-90	2.6	17
4	Morbidity, mortality, and placental pathology in excessively long umbilical cords: retrospective study. <i>Pediatric and Developmental Pathology</i> , 2001 , 4, 144-53	2.2	101
3	Human pancreatic adenocarcinomas express parathyroid hormone-related protein. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2001 , 86, 310-6	5.6	16
2	Comparison of the toxicities of two iron formulations in a swine model. <i>Academic Emergency Medicine</i> , 1999 , 6, 1104-8	3.4	6
1	How to Interpret EUS-FNA Cytology133-143		