
Bernard Renaud

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11200152/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Clinical CNS Microdialysis of Clutamate with a Special Methodological Focus on Human Spinal Cord. Neuromethods, 2018, , 523-558.	0.3	1
2	Outcome of Poor-Grade Subarachnoid Hemorrhage as Determined by Biomarkers of Glucose Cerebral Metabolism. Neurocritical Care, 2013, 18, 234-244.	2.4	13
3	In-capillary derivatization and capillary electrophoresis separation of amino acid neurotransmitters from brain microdialysis samples. Journal of Chromatography A, 2008, 1205, 144-149.	3.7	44
4	Highly sensitive assay for the measurement of serotonin in microdialysates using capillary high-performance liquid chromatography with electrochemical detection. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2007, 850, 303-309.	2.3	23
5	Dopaminergic transmission in STOP null mice. Journal of Neurochemistry, 2005, 94, 63-73.	3.9	65
6	Analysis of serotonin in brain microdialysates using capillary electrophoresis and native laser-induced fluorescence detection. Electrophoresis, 2005, 26, 1071-1079.	2.4	32
7	High temporal resolution for in vivo monitoring of neurotransmitters in awake epileptic rats using brain microdialysis and capillary electrophoresis with laser-induced fluorescence detection. Journal of Neuroscience Methods, 2004, 140, 29-38.	2.5	83
8	Capillary electrophoresis combined with microdialysis in the human spinal cord: A new tool for monitoring rapid peroperative changes in amino acid neurotransmitters within the dorsal horn. Electrophoresis, 2004, 25, 1511-1517.	2.4	30
9	Simultaneous determination of vigabatrin and amino acid neurotransmitters in brain microdialysates by capillary electrophoresis with laser-induced fluorescence detection. Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences, 2004, 806, 237-244.	2.3	50
10	Assessment of Jugular Blood Oxygen and Lactate Indices for Detection of Cerebral Ischemia and Prognosis. Journal of Neurosurgical Anesthesiology, 2004, 16, 226-231.	1.2	36
11	Microdialysis monitoring of catecholamines and excitatory amino acids in the rat and mouse brain: recent developments based on capillary electrophoresis with laser-induced fluorescence detectiona mini-review. Cellular and Molecular Neurobiology, 2003, 23, 793-804.	3.3	54
12	In vivo comparison of two 5-HT1A receptors agonists alnespirone (S-20499) and buspirone on locus coeruleus neuronal activity. European Journal of Pharmacology, 2003, 459, 17-26.	3.5	23
13	In vivo simultaneous monitoring of ^{ĵ3} -aminobutyric acid, glutamate, andL-aspartate using brain microdialysis and capillary electrophoresis with laser-induced fluorescence detection: Analytical developments andin vitro/in vivo validations. Electrophoresis, 2003, 24, 3187-3196.	2.4	81
14	Glutamate and aspartate do not exhibit the same changes in their extracellular concentrations in the rat striatum after N-methyl-D-aspartate local administration. Journal of Neuroscience Research, 2003, 71, 445-454.	2.9	15
15	The Switch of Subthalamic Neurons From an Irregular to a Bursting Pattern Does Not Solely Depend on Their GABAergic Inputs in the Anesthetic-Free Rat. Journal of Neuroscience, 2002, 22, 8665-8675.	3.6	58
16	Determination of Subnanomolar Concentrations of Dopamine and Norepinephrine in Nanovolume Samples Using an Automated Capillary Zone Electrophoresis with Laser Induced Fluorescence Detection. Advances in Behavioral Biology, 2002, , 309-312.	0.2	0
17	Microdialysis study of amino acid neurotransmitters in the spinal dorsal horn of patients undergoing microsurgical dorsal root entry zone lesioning. Journal of Neurosurgery: Spine, 2001, 94, 165-173.	1.7	10
18	Amino acids in spinal dorsal horn of patients during surgery for neuropathic pain or spasticity. NeuroReport, 2000, 11, 1795-1798.	1.2	19

Bernard Renaud

#	Article	IF	CITATIONS
19	Single-unit and polygraphic recordings associated with systemic or local pharmacology: A multi-purpose stereotaxic approach for the awake, anaesthetic-free, and head-restrained rat. Journal of Neuroscience Research, 2000, 61, 88-100.	2.9	24
20	Unrelated course of subthalamic nucleus and globus pallidus neuronal activities across vigilance states in the rat. European Journal of Neuroscience, 2000, 12, 3361-3374.	2.6	94
21	Evidence that the neuronal nitric oxide synthase inhibitor 7-nitroindazole inhibits monoamine oxidase in the rat: in vivo effects on extracellular striatal dopamine and 3,4-dihydroxyphenylacetic acid. Neuroscience Letters, 1999, 261, 175-178.	2.1	50
22	Evidence that the neuronal nitric oxide synthase inhibitor 7-nitroindazole inhibits monoamine oxidase in the rat: in vivo effects on extracellular striatal dopamine and 3,4-dihydroxyphenylacetic acid. Neuroscience Letters, 1999, 264, 5-8.	2.1	29
23	Assessment of pharmacodynamic and pharmacokinetic characteristics of drugs using microdialysis sampling and capillary electrophoresis. Electrophoresis, 1998, 19, 2841-2847.	2.4	29
24	Monitoring nitric oxide (NO) in rat locus coeruleus. NeuroReport, 1997, 8, 1321-1325.	1.2	32
25	Microdialysis monitoring of variations in extracellular levels of serotonin, GABA and excitatory amino acids in the frontal cortex of awake rats in response to a single peripheral or central administration of dexfenfluramine. Brain Research, 1996, 737, 221-230.	2.2	32
26	High-speed separation of subnanomolar concentrations of noradrenaline and dopamine using capillary zone electrophoresis with laser-induced fluorescence detection. Electrophoresis, 1996, 17, 523-525.	2.4	32
27	In vivo monitoring of extracellular noradrenaline and glutamate from rat brain cortex with 2-min microdialysis sampling using capillary electrophoresis with laser-induced fluorescence detection. Journal of Neuroscience Methods, 1996, 70, 153-162.	2.5	51
28	Capillary zone electrophoresis with laser-induced fluorescence detection for the determination of nanomolar concentrations of noradrenaline and dopamine: application to brain microdialyzate analysis. Analytical Chemistry, 1995, 67, 1838-1844.	6.5	112
29	Biochemical evidence for an interaction between adrenaline and noradrenaline neurons in the rat brainstem. Brain Research, 1986, 397, 333-340.	2.2	19
30	Changes in tyrosine hydroxylase and dopamine-beta-hydroxylase activities but not in phenylethanolamine-N-methyltransferase activity within central adrenaline neurons after 6-hydroxydopamine administration. Biochemical Pharmacology, 1984, 33, 1887-1891.	4.4	9
31	Effects of chronic β-blocker treatment on catecholamine levels in spontaneously hypertensive rats. Biochemical Pharmacology, 1983, 32, 2739-2743.	4.4	6
32	Increased tyrosine hydroxylase activity in central adrenaline neurons after reserpine treatment. European Journal of Pharmacology, 1983, 92, 243-248.	3.5	7
33	Effects of chronic β-blockers treatment on catecholamine synthesizing enzymes in spontaneously hypertensive rats. Biochemical Pharmacology, 1981, 30, 2673-2678.	4.4	15
34	Early increase in phenylethanolamine-N-methyltransferase activity in a new strain of spontaneously hypertensive rats. Brain Research, 1978, 159, 149-159.	2.2	49
35	Time-course variations in tyrosine hydroxylase activity in the rat locus coeruleus after electrolytic destruction of the nuclei raphe dorsalis or raphe centralis. Brain Research, 1976, 108, 339-349.	2.2	69
36	Increase in tyrosine hydroxylase activity in the locus coeruleus of the rat brain after contralateral lesioning. Brain Research, 1975, 93, 564-569.	2.2	77

#	Article	IF	CITATIONS
37	Effects of 5,6-dihydroxytryptamine on tyrosine-hydroxylase activity in central catecholaminergic neurons of the rat. Biochemical Pharmacology, 1975, 24, 1739-1742.	4.4	76