

# Jane A English

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11217206/publications.pdf>

Version: 2024-02-01

14  
papers

610  
citations

933447

10  
h-index

1058476

14  
g-index

15  
all docs

15  
docs citations

15  
times ranked

3328  
citing authors

#	ARTICLE	IF	CITATIONS
1	2D DIGE analysis implicates cytoskeletal abnormalities in psychiatric disease. <i>Proteomics</i> , 2009, 9, 3368-3382.	2.2	134
2	The Neuroproteomics of Schizophrenia. <i>Biological Psychiatry</i> , 2011, 69, 163-172.	1.3	122
3	Common Proteomic Changes in the Hippocampus in Schizophrenia and Bipolar Disorder and Particular Evidence for Involvement of Cornu Ammonis Regions 2 and 3. <i>Archives of General Psychiatry</i> , 2011, 68, 477.	12.3	90
4	Comparative analysis of OFFGel, strong cation exchange with pH gradient, and RP at high pH for first-dimensional separation of peptides from a membrane-enriched protein fraction. <i>Proteomics</i> , 2009, 9, 5194-5198.	2.2	63
5	Blood-Based Protein Changes in Childhood Are Associated With Increased Risk for Later Psychotic Disorder: Evidence From a Nested Case-Control Study of the ALSPAC Longitudinal Birth Cohort. <i>Schizophrenia Bulletin</i> , 2018, 44, 297-306.	4.3	53
6	Image analysis tools and emerging algorithms for expression proteomics. <i>Proteomics</i> , 2010, 10, 4226-4257.	2.2	46
7	Blood biomarker discovery in drug-free schizophrenia: the contribution of proteomics and multiplex immunoassays. <i>Expert Review of Proteomics</i> , 2016, 13, 1141-1155.	3.0	38
8	Integrated Lipidomics and Proteomics Point to Early Blood-Based Changes in Childhood Preceding Later Development of Psychotic Experiences: Evidence From the Avon Longitudinal Study of Parents and Children. <i>Biological Psychiatry</i> , 2019, 86, 25-34.	1.3	26
9	Omega-3 fatty acid deficiency disrupts endocytosis, neuritogenesis, and mitochondrial protein pathways in the mouse hippocampus. <i>Frontiers in Genetics</i> , 2013, 4, 208.	2.3	17
10	Partitioning the Proteome: Phase Separation for Targeted Analysis of Membrane Proteins in Human Post-Mortem Brain. <i>PLoS ONE</i> , 2012, 7, e39509.	2.5	10
11	Targeted Proteomics for Validation of Biomarkers in Early Psychosis. <i>Biological Psychiatry</i> , 2014, 76, e7-e9.	1.3	5
12	Dataset of mouse hippocampus profiled by LC-MS/MS for label-free quantitation. <i>Data in Brief</i> , 2016, 7, 341-343.	1.0	3
13	ApoE elevation is associated with the persistence of psychotic experiences from age 12 to age 18: Evidence from the ALSPAC birth cohort. <i>Schizophrenia Research</i> , 2019, 209, 141-147.	2.0	2
14	Poster #179 PROTEOMIC ANALYSIS OF THE CORPUS CALLOSUM REVEALS DOWN-REGULATION OF THE GLYCOLYSIS PROTEIN PATHWAY AND FURTHER IMPLICATES CME IN SCHIZOPHRENIA. <i>Schizophrenia Research</i> , 2012, 136, S155.	2.0	1