

Olivier Huet

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/1122543/publications.pdf>

Version: 2024-02-01

45
papers

2,260
citations

257450

24
h-index

233421

45
g-index

45
all docs

45
docs citations

45
times ranked

3372
citing authors

#	ARTICLE	IF	CITATIONS
1	A Post Hoc Analysis of Osmotherapy Use in the Erythropoietin in Traumatic Brain Injury Study—Associations With Acute Kidney Injury and Mortality. <i>Critical Care Medicine</i> , 2021, 49, e394-e403.	0.9	14
2	Effect of Continuous Infusion of Hypertonic Saline vs Standard Care on 6-Month Neurological Outcomes in Patients With Traumatic Brain Injury. <i>JAMA - Journal of the American Medical Association</i> , 2021, 325, 2056.	7.4	64
3	Management and prevention of anemia (acute bleeding excluded) in adult critical care patients. <i>Annals of Intensive Care</i> , 2020, 10, 97.	4.6	24
4	Management and prevention of anemia (acute bleeding excluded) in adult critical care patients. <i>Anaesthesia, Critical Care & Pain Medicine</i> , 2020, 39, 655-664.	1.4	11
5	Erythropoietin in traumatic brain injury associated acute kidney injury: A randomized controlled trial. <i>Acta Anaesthesiologica Scandinavica</i> , 2019, 63, 200-207.	1.6	24
6	Cost-Effectiveness of Erythropoietin in Traumatic Brain Injury: A Multinational Trial-Based Economic Analysis. <i>Journal of Neurotrauma</i> , 2019, 36, 2541-2548.	3.4	12
7	Cause and Timing of Death and Subgroup Differential Effects of Erythropoietin in the EPO-TBI Study. <i>Journal of Neurotrauma</i> , 2018, 35, 333-340.	3.4	13
8	Effect of Early Sustained Prophylactic Hypothermia on Neurologic Outcomes Among Patients With Severe Traumatic Brain Injury. <i>JAMA - Journal of the American Medical Association</i> , 2018, 320, 2211.	7.4	226
9	Statistical analysis plan for the POLAR-RCT: The Prophylactic hypOthermia trial to Lessen trAumatic bRain injury-Randomised Controlled Trial. <i>Trials</i> , 2018, 19, 259.	1.6	9
10	Transactivation of RAGE mediates angiotensin-induced inflammation and atherogenesis. <i>Journal of Clinical Investigation</i> , 2018, 129, 406-421.	8.2	59
11	Protective Effect of Inflammasome Activation by Hydrogen Peroxide in a Mouse Model of Septic Shock. <i>Critical Care Medicine</i> , 2017, 45, e184-e194.	0.9	9
12	Erythropoietin in patients with traumatic brain injury and extracranial injury—A post hoc analysis of the erythropoietin traumatic brain injury trial. <i>Journal of Trauma and Acute Care Surgery</i> , 2017, 83, 449-456.	2.1	14
13	Venous thromboembolic events in critically ill traumatic brain injury patients. <i>Intensive Care Medicine</i> , 2017, 43, 419-428.	8.2	86
14	COBI (COntinuous hyperosmolar therapy for traumatic Brain-Injured patients) trial protocol: a multicentre randomised open-label trial with blinded adjudication of primary outcome. <i>BMJ Open</i> , 2017, 7, e018035.	1.9	19
15	Nitroxyl (HNO) reduces endothelial and monocyte activation and promotes M2 macrophage polarization. <i>Clinical Science</i> , 2016, 130, 1629-1640.	4.3	18
16	Compound 21, a selective agonist of angiotensin AT ₂ receptors, prevents endothelial inflammation and leukocyte adhesion <i>in vitro</i> and <i>in vivo</i> . <i>British Journal of Pharmacology</i> , 2016, 173, 729-740.	5.4	51
17	Lack of glutathione peroxidase-1 facilitates a pro-inflammatory and activated vascular endothelium. <i>Vascular Pharmacology</i> , 2016, 79, 32-42.	2.1	37
18	Norepinephrine Decreases Fluid Requirements and Blood Loss While Preserving Intestinal Villi Microcirculation during Fluid Resuscitation of Uncontrolled Hemorrhagic Shock in Mice. <i>Anesthesiology</i> , 2015, 122, 1093-1102.	2.5	38

#	ARTICLE	IF	CITATIONS
19	Stored red blood cell susceptibility to in vitro transfusion-associated stress conditions is higher after longer storage and increased by storage in saline-glucose-mannitol compared to AS ¹ . <i>Transfusion</i> , 2015, 55, 2197-2206.	1.6	21
20	Direct Endothelial Nitric Oxide Synthase Activation Provides Atheroprotection in Diabetes-Accelerated Atherosclerosis. <i>Diabetes</i> , 2015, 64, 3937-3950.	0.6	60
21	Erythropoietin in traumatic brain injury: study protocol for a randomised controlled trial. <i>Trials</i> , 2015, 16, 39.	1.6	27
22	Erythropoietin in traumatic brain injury (EPO-TBI): a double-blind randomised controlled trial. <i>Lancet, The</i> , 2015, 386, 2499-2506.	13.7	217
23	Dicarbonyl Stress in the Absence of Hyperglycemia Increases Endothelial Inflammation and Atherogenesis Similar to That Observed in Diabetes. <i>Diabetes</i> , 2014, 63, 3915-3925.	0.6	74
24	Microcirculatory Alterations in Traumatic Hemorrhagic Shock*. <i>Critical Care Medicine</i> , 2014, 42, 1433-1441.	0.9	152
25	Septic shock: desperately seeking treatment. <i>Clinical Science</i> , 2014, 126, 31-39.	4.3	36
26	The ethical dimension in published animal research in critical care: the dark side of our moon. <i>Critical Care</i> , 2014, 18, 120.	5.8	6
27	Statistical analysis plan for the Erythropoietin in Traumatic Brain Injury trial: a randomised controlled trial of erythropoietin versus placebo in moderate and severe traumatic brain injury. <i>Trials</i> , 2014, 15, 501.	1.6	16
28	Ensuring Animal Welfare While Meeting Scientific Aims Using a Murine Pneumonia Model of Septic Shock. <i>Shock</i> , 2013, 39, 488-494.	2.1	60
29	Synergistic Deleterious Effect of Hypoxemia and Hypovolemia on Microcirculation in Intestinal Villi*. <i>Critical Care Medicine</i> , 2013, 41, e376-e384.	0.9	23
30	Interleukin 10 Antioxidant Effect Decreases Leukocytes/Endothelial Interaction Induced by Tumor Necrosis Factor α . <i>Shock</i> , 2013, 39, 83-88.	2.1	34
31	The Limits of Succinylcholine for Critically Ill Patients. <i>Anesthesia and Analgesia</i> , 2012, 115, 873-879.	2.2	37
32	Activation of the Renin-Angiotensin System Mediates the Effects of Dietary Salt Intake on Atherogenesis in the Apolipoprotein E Knockout Mouse. <i>Hypertension</i> , 2012, 60, 98-105.	2.7	48
33	Variation in endogenous oxidative stress in <i>Escherichia coli</i> natural isolates during growth in urine. <i>BMC Microbiology</i> , 2012, 12, 120.	3.3	25
34	Changes in urine composition after trauma facilitate bacterial growth. <i>BMC Infectious Diseases</i> , 2012, 12, 330.	2.9	17
35	Coenzyme Q10 deficiency in septic shock patients. <i>Critical Care</i> , 2011, 15, 194.	5.8	7
36	Oxidative stress and endothelial dysfunction during sepsis. <i>Frontiers in Bioscience - Landmark</i> , 2011, 16, 1986.	3.0	100

#	ARTICLE	IF	CITATIONS
37	Postresuscitation syndrome: Potential role of hydroxyl radical-induced endothelial cell damage*. Critical Care Medicine, 2011, 39, 1712-1720.	0.9	57
38	In Vitro Plasma-Induced Endothelial Oxidative Stress and Circulating Markers of Endothelial Dysfunction in Preeclampsia: An Observational Study. Hypertension in Pregnancy, 2009, 28, 212-223.	1.1	13
39	Alterations of mitochondrial function in sepsis and critical illness. Current Opinion in Anaesthesiology, 2009, 22, 143-149.	2.0	96
40	Changes in cerebral blood flow and oxygen extraction during post-resuscitation syndrome. Resuscitation, 2008, 76, 17-24.	3.0	115
41	Pivotal role of glutathione depletion in plasma-induced endothelial oxidative stress during sepsis. Critical Care Medicine, 2008, 36, 2328-2334.	0.9	42
42	Plasma-induced endothelial oxidative stress is related to the severity of septic shock*. Critical Care Medicine, 2007, 35, 821-826.	0.9	90
43	Î2-adrenergic agonist protects human endothelial cells from hypoxia/reoxygenation injury in vitro. Critical Care Medicine, 2006, 34, 165-172.	0.9	8
44	Endothelial oxidative stress induced by serum from patients with severe trauma hemorrhage. Intensive Care Medicine, 2005, 31, 1174-1180.	8.2	31
45	Cardiac Arrest after Injection of Ropivacaine for Posterior Lumbar Plexus Blockade. Anesthesiology, 2003, 99, 1451-1453.	2.5	120