

Sheila M Fleming

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11244867/publications.pdf>

Version: 2024-02-01

53
papers

5,330
citations

159585

30
h-index

197818

49
g-index

55
all docs

55
docs citations

55
times ranked

6910
citing authors

#	ARTICLE	IF	CITATIONS
1	Targeting alpha-synuclein via the immune system in Parkinson's disease: Current vaccine therapies. <i>Neuropharmacology</i> , 2022, 202, 108870.	4.1	14
2	Farey Trees Explain Sequential Effects in Choice Response Time. <i>Frontiers in Physiology</i> , 2021, 12, 611145.	2.8	2
3	Irisin treatment lowers levels of phosphorylated tau in the hippocampus of pre-symptomatic female but not male htau mice. <i>Neuropathology and Applied Neurobiology</i> , 2021, 47, 967-978.	3.2	10
4	Pharmacological inhibition of CSF1R by GW2580 reduces microglial proliferation and is protective against neuroinflammation and dopaminergic neurodegeneration. <i>FASEB Journal</i> , 2020, 34, 1679-1694.	0.5	66
5	Developmental exposure to the organochlorine pesticide dieldrin causes male-specific exacerbation of α -synuclein-preformed fibril-induced toxicity and motor deficits. <i>Neurobiology of Disease</i> , 2020, 141, 104947.	4.4	24
6	Time course and magnitude of alpha-synuclein inclusion formation and nigrostriatal degeneration in the rat model of synucleinopathy triggered by intrastriatal α -synuclein preformed fibrils. <i>Neurobiology of Disease</i> , 2019, 130, 104525.	4.4	67
7	Exacerbation of sensorimotor dysfunction in mice deficient in <i>Atp13a2</i> and overexpressing human wildtype alpha-synuclein. <i>Behavioural Brain Research</i> , 2018, 343, 41-49.	2.2	17
8	<i>Ahr</i> and <i>Cyp1a2</i> genotypes both affect susceptibility to motor deficits following gestational and lactational exposure to polychlorinated biphenyls. <i>NeuroToxicology</i> , 2018, 65, 125-134.	3.0	11
9	The effect of manganese exposure in <i>Atp13a2</i> -deficient mice. <i>NeuroToxicology</i> , 2018, 64, 256-266.	3.0	21
10	Mechanisms of Gene-Environment Interactions in Parkinson's Disease. <i>Current Environmental Health Reports</i> , 2017, 4, 192-199.	6.7	74
11	Intranasal <i>NAP</i> (davunetide) decreases tau hyperphosphorylation and moderately improves behavioral deficits in mice overexpressing α -synuclein. <i>Pharmacology Research and Perspectives</i> , 2014, 2, e00065.	2.4	40
12	Ribosomal <i>s15</i> : A novel therapeutic target for Parkinson's disease. <i>Movement Disorders</i> , 2014, 29, 990-990.	3.9	0
13	Cinnamon in a mouse model of PD. <i>Movement Disorders</i> , 2014, 29, 1466-1466.	3.9	5
14	A GCCase Chaperone Improves Motor Function in a Mouse Model of Synucleinopathy. <i>Neurotherapeutics</i> , 2014, 11, 840-856.	4.4	88
15	Chronic administration of cholesterol oximes in mice increases transcription of cytoprotective genes and improves transcriptome alterations induced by alpha-synuclein overexpression in nigrostriatal dopaminergic neurons. <i>Neurobiology of Disease</i> , 2014, 69, 263-275.	4.4	28
16	Olfactory Assays for Mouse Models of Neurodegenerative Disease. <i>Journal of Visualized Experiments</i> , 2014, , e51804.	0.3	64
17	Impaired Baroreflex Function in Mice Overexpressing Alpha-Synuclein. <i>Frontiers in Neurology</i> , 2013, 4, 103.	2.4	20
18	<i>Atp13a2</i> -deficient mice exhibit neuronal ceroid lipofuscinosis, limited α -synuclein accumulation and age-dependent sensorimotor deficits. <i>Human Molecular Genetics</i> , 2013, 22, 2067-2082.	2.9	124

#	ARTICLE	IF	CITATIONS
19	Assessment of Sensorimotor Function in Mouse Models of Parkinson's Disease. <i>Journal of Visualized Experiments</i> , 2013, , .	0.3	115
20	Morphological and Behavioral Impact of AAV2/5-Mediated Overexpression of Human Wildtype Alpha-Synuclein in the Rat Nigrostriatal System. <i>PLoS ONE</i> , 2013, 8, e81426.	2.5	70
21	Sensorimotor assessment of the unilateral 6-hydroxydopamine mouse model of Parkinson's disease. <i>Behavioural Brain Research</i> , 2012, 230, 309-316.	2.2	108
22	Cranial and related sensorimotor impairments in rodent models of Parkinson's disease. <i>Behavioural Brain Research</i> , 2012, 231, 317-322.	2.2	21
23	Cognitive deficits in a mouse model of pre-manifest Parkinson's disease. <i>European Journal of Neuroscience</i> , 2012, 35, 870-882.	2.6	87
24	Cardiovascular Autonomic Dysfunction in Animal Models of Parkinson's Disease. <i>Journal of Parkinson's Disease</i> , 2011, 1, 321-327.	2.8	11
25	A pilot trial of the microtubule-interacting peptide (NAP) in mice overexpressing alpha-synuclein shows improvement in motor function and reduction of alpha-synuclein inclusions. <i>Molecular and Cellular Neurosciences</i> , 2011, 46, 597-606.	2.2	68
26	HCN channelopathy in external globus pallidus neurons in models of Parkinson's disease. <i>Nature Neuroscience</i> , 2011, 14, 85-92.	14.8	160
27	Comparing Behavioral Assessment of Sensorimotor Function in Rat and Mouse Models of Parkinson's Disease and Stroke. <i>Neuromethods</i> , 2011, , 325-335.	0.3	0
28	Analysis of striatal transcriptome in mice overexpressing human wild-type alpha-synuclein supports synaptic dysfunction and suggests mechanisms of neuroprotection for striatal neurons. <i>Molecular Neurodegeneration</i> , 2011, 6, 83.	10.8	30
29	Cognitive Dysfunction in Genetic Mouse Models of Parkinsonism. <i>Neuromethods</i> , 2011, , 485-492.	0.3	2
30	Mice overexpressing corticotropin-releasing factor show brain atrophy and motor dysfunctions. <i>Neuroscience Letters</i> , 2010, 473, 11-15.	2.1	17
31	Behavioral Outcome Measures for the Assessment of Sensorimotor Function in Animal Models of Movement Disorders. <i>International Review of Neurobiology</i> , 2009, 89, 57-65.	2.0	17
32	Secretion of amyloidogenic gelsolin progressively compromises protein homeostasis leading to the intracellular aggregation of proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009, 106, 11125-11130.	7.1	49
33	Bacterial Artificial Chromosome Transgenic Mice Expressing a Truncated Mutant Parkin Exhibit Age-Dependent Hypokinetic Motor Deficits, Dopaminergic Neuron Degeneration, and Accumulation of Proteinase K-Resistant α -Synuclein. <i>Journal of Neuroscience</i> , 2009, 29, 1962-1976.	3.6	168
34	Modeling Nonmotor Symptoms of Parkinson's Disease in Genetic Mouse Models. <i>Advances in Behavioral Biology</i> , 2009, , 483-491.	0.2	3
35	Olfactory deficits in mice overexpressing human wildtype α -synuclein. <i>European Journal of Neuroscience</i> , 2008, 28, 247-256.	2.6	182
36	Effects of antioxidants on cancer prevention and neuromotor performance in Atm deficient mice. <i>Food and Chemical Toxicology</i> , 2008, 46, 1371-1377.	3.6	30

#	ARTICLE	IF	CITATIONS
37	Abnormal colonic motility in mice overexpressing human wild-type α -synuclein. <i>NeuroReport</i> , 2008, 19, 873-876.	1.2	110
38	Behavioral phenotypes and pharmacology in genetic mouse models of Parkinsonism. <i>Behavioural Pharmacology</i> , 2006, 17, 383-391.	1.7	59
39	Phenotypical Characterization of Genetic Mouse Models of Parkinson Disease. , 2005, , 183-192.		4
40	3,4-Dihydroxyphenylalanine Reverses the Motor Deficits in Pitx3-Deficient Aphakia Mice: Behavioral Characterization of a Novel Genetic Model of Parkinson's Disease. <i>Journal of Neuroscience</i> , 2005, 25, 2132-2137.	3.6	162
41	An intermittent, controlled-rate, slow progressive degeneration model of Parkinson's disease: antiparkinson effects of Sinemet and protective effects of methylphenidate. <i>Behavioural Brain Research</i> , 2005, 156, 201-213.	2.2	57
42	Genetic mouse models of parkinsonism: Strengths and limitations. <i>NeuroRx</i> , 2005, 2, 495-503.	6.0	150
43	Early Behavioral Phenotypes in Mouse Models of Huntington's and Parkinson's Diseases. , 2005, , 349-359.		0
44	Variable effects of chronic subcutaneous administration of rotenone on striatal histology. <i>Journal of Comparative Neurology</i> , 2004, 478, 418-426.	1.6	86
45	Early and Progressive Sensorimotor Anomalies in Mice Overexpressing Wild-Type Human α -Synuclein. <i>Journal of Neuroscience</i> , 2004, 24, 9434-9440.	3.6	428
46	Behavioral and immunohistochemical effects of chronic intravenous and subcutaneous infusions of varying doses of rotenone. <i>Experimental Neurology</i> , 2004, 187, 418-429.	4.1	179
47	Genetic mouse models of Huntington's and Parkinson's diseases: illuminating but imperfect. <i>Trends in Neurosciences</i> , 2004, 27, 691-697.	8.6	170
48	Should the injured and intact hemispheres be treated differently during the early phases of physical restorative therapy in experimental stroke or parkinsonism?. <i>Physical Medicine and Rehabilitation Clinics of North America</i> , 2003, 14, S27-S46.	1.3	36
49	Experimental Focal Ischemic Injury: Behavior-Brain Interactions and Issues of Animal Handling and Housing. <i>ILAR Journal</i> , 2003, 44, 130-143.	1.8	34
50	Parkin-deficient Mice Exhibit Nigrostriatal Deficits but Not Loss of Dopaminergic Neurons. <i>Journal of Biological Chemistry</i> , 2003, 278, 43628-43635.	3.4	784
51	CNS plasticity and assessment of forelimb sensorimotor outcome in unilateral rat models of stroke, cortical ablation, parkinsonism and spinal cord injury. <i>Neuropharmacology</i> , 2000, 39, 777-787.	4.1	1,217
52	Prefrontal cortex infusions of SCH 23390 cause immediate and delayed effects on ventral tegmental area stimulation reward. <i>Brain Research</i> , 1998, 811, 57-62.	2.2	14
53	Involvement of δ - and μ -opioid receptors in the potentiation of brain-stimulation reward. <i>European Journal of Pharmacology</i> , 1996, 316, 137-143.	3.5	26