

Francisco Puerta Martinez

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11261307/publications.pdf>

Version: 2024-02-01

8
papers

145
citations

1478505

6
h-index

1588992

8
g-index

8
all docs

8
docs citations

8
times ranked

252
citing authors

#	ARTICLE	IF	CITATIONS
1	CTCF Binding to the First Intron of the Major Immediate Early (MIE) Gene of Human Cytomegalovirus (HCMV) Negatively Regulates MIE Gene Expression and HCMV Replication. <i>Journal of Virology</i> , 2014, 88, 7389-7401.	3.4	45
2	Murine cytomegalovirus major immediate-early protein 3 interacts with cellular and viral proteins in viral DNA replication compartments and is important for early gene activation. <i>Journal of General Virology</i> , 2010, 91, 2664-2676.	2.9	28
3	Nuclear domain 10 of the viral aspect. <i>World Journal of Virology</i> , 2013, 2, 110.	2.9	21
4	Leucine Zipper Domain Is Required for Kaposi Sarcoma-associated Herpesvirus (KSHV) K-bZIP Protein to Interact with Histone Deacetylase and Is Important for KSHV Replication. <i>Journal of Biological Chemistry</i> , 2012, 287, 15622-15634.	3.4	19
5	Functional Interaction of Nuclear Domain 10 and Its Components with Cytomegalovirus after Infections: Cross-Species Host Cells versus Native Cells. <i>PLoS ONE</i> , 2011, 6, e19187.	2.5	14
6	A Short cis-Acting Motif in the M112-113 Promoter Region Is Essential for IE3 To Activate M112-113 Gene Expression and Is Important for Murine Cytomegalovirus Replication. <i>Journal of Virology</i> , 2013, 87, 2639-2647.	3.4	9
7	Evidence of inability of human cytomegalovirus to reactivate Kaposi's sarcoma-associated herpesvirus from latency in body cavity-based lymphocytes. <i>Journal of Clinical Virology</i> , 2009, 46, 244-248.	3.1	5
8	H2B homology region of major immediate-early protein 1 is essential for murine cytomegalovirus to disrupt nuclear domain 10, but is not important for viral replication in cell culture. <i>Journal of General Virology</i> , 2011, 92, 2006-2019.	2.9	4