

Subramaniam Sotheeswaran

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11269018/publications.pdf>

Version: 2024-02-01

37
papers

1,360
citations

361296
20
h-index

330025
37
g-index

40
all docs

40
docs citations

40
times ranked

1281
citing authors

#	ARTICLE	IF	CITATIONS
1	Anti-HIV coumarins from calophyllum seed oil. Bioorganic and Medicinal Chemistry Letters, 1998, 8, 3475-3478.	1.0	260
2	Distribution of resveratrol oligomers in plants. Phytochemistry, 1993, 32, 1083-1092.	1.4	217
3	Identification of character-impact odorants in coriander and wild coriander leaves using gas chromatography-olfactometry (GCO) and comprehensive two-dimensional gas chromatography-time-of-flight mass spectrometry (GC-TOFMS). Journal of Separation Science, 2005, 28, 1061-1074.	1.3	107
4	Ecdysterone from stem of Diploclisia glaucescens. Phytochemistry, 1989, 28, 1073-1075.	1.4	65
5	¹³ C NMR studies of some lupane and taraxerane triterpenes. Magnetic Resonance in Chemistry, 1980, 14, 462-465.	0.7	59
6	Canaliculatol, an antifungal resveratrol trimer from Stemonoporus canaliculatus. Phytochemistry, 1988, 27, 377-380.	1.4	54
7	Balanocarpol, a new polyphenol from Balanocarpus zeylanicus (trimen) and Hopea jucunda (Thw.) (Dipterocarpaceae). Journal of the Chemical Society Perkin Transactions 1, 1985, , 1807.	0.9	43
8	Triterpenoids and coumarins from the leaves of calophyllum cordato-oblongum. Phytochemistry, 1985, 24, 1553-1556.	1.4	36
9	Another antibacterial polyphenol, copalliferol B, from Vateria copallifera (dipterocarpaceae). Phytochemistry, 1986, 25, 1498-1500.	1.4	36
10	Terpenoid and biflavonoid constituents of Calophyllum calaba and Garcinia spicata from Sri Lanka. Phytochemistry, 1984, 23, 323-328.	1.4	35
11	Polyphenols from Stemonoporus species. Phytochemistry, 1982, 21, 2585-2587.	1.4	32
12	Polyphenols from dipterocarp species. Vaticaffinol and μ -viniferin. Journal of the Chemical Society Perkin Transactions 1, 1985, , 159-162.	0.9	31
13	Horsfieldin, a lignan and other constituents from Horsfieldia iryagedhi. Phytochemistry, 1982, 21, 2719-2723.	1.4	29
14	3,5,7,3',5'-Pentahydroxyflavan and 3'-methoxyfriedelan from Humboldtia laurifolia. Phytochemistry, 1983, 22, 565-567.	1.4	29
15	Comparison of the Composition of the Essential Oils of Alpinia and Hedychium Species "Essential Oils of Fijian Plants, Part 1. Journal of Essential Oil Research, 2002, 14, 409-411.	1.3	28
16	A hemolytic saponin, randianin, from Randia dumetorum. Phytochemistry, 1989, 28, 1544-1546.	1.4	26
17	Diploclisin, a bidesmosidic triterpenoid saponin from Diploclisia glaucescens. Phytochemistry, 1989, 28, 2783-2785.	1.4	23
18	Polyphenols from dipterocarp species. Copalliferol A and stemonoporol. Journal of the Chemical Society Perkin Transactions 1, 1983, , 699.	0.9	22

#	ARTICLE	IF	CITATIONS
19	Two chemically distinct groups of calophyllum species from Sri Lanka. <i>Phytochemistry</i> , 1986, 25, 425-428.	1.4	22
20	Xanthones from roots of three calophyllum species. <i>Phytochemistry</i> , 1986, 25, 1957-1959.	1.4	20
21	Further evidence for the structure of vaticaffinol and a revision of its stereochemistry. <i>Phytochemistry</i> , 1987, 26, 1505-1507.	1.4	18
22	Benzophenones of <i>Garcinia pseudoguttifera</i> (Clusiaceae). <i>Phytochemistry</i> , 2000, 53, 281-284.	1.4	18
23	Spongisoritin A: A New Polyketide from a Fijian Marine Sponge, <i>Spongisorites</i> sp.. <i>Australian Journal of Chemistry</i> , 2005, 58, 18.	0.5	18
24	Two new xanthones, calozeyloxanthonone and zeyloxanthonone, from <i>Calophyllum zeylanicum</i> (Guttiferae). <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1981, , 1831.	0.9	15
25	Triterpenes and neoflavonoids of <i>Calophyllum lankaensis</i> and <i>Calophyllum thwaitesii</i> . <i>Phytochemistry</i> , 1984, 23, 2601-2603.	1.4	15
26	Bark acids of seven <i>Calophyllum</i> species (Guttiferae). <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1983, , 703.	0.9	13
27	Three prenylated phenolic benzophenones from <i>Garcinia myrtifolia</i> . <i>Phytochemistry</i> , 1995, 38, 233-236.	1.4	13
28	Calocalabaxanthonone, the putative isoprenyl precursor of calabaxanthonone from <i>Calophyllum calaba</i> . <i>Phytochemistry</i> , 1982, 21, 807-809.	1.4	12
29	Triterpenoid taxonomic markers for <i>Stemonoporus</i> and other genera of the dipterocarpaceae. <i>Phytochemistry</i> , 1977, 16, 699-701.	1.4	11
30	Convenient preparation of esters of testosterone and 19-norethisterone with hindered acids. <i>Journal of the Chemical Society Chemical Communications</i> , 1978, , 980.	2.0	11
31	Pachyovatamine, a bisbenzylisoquinoline alkaloid, and other alkaloids from <i>Pachygone ovata</i> . <i>Phytochemistry</i> , 1985, 24, 589-592.	1.4	10
32	A revised structure for hermonionic acid. <i>Tetrahedron Letters</i> , 1982, 23, 2987-2990.	0.7	7
33	Pachygonamine and N-Methylpachygonamine, New Bisbenzylisoquinoline Alkaloids from <i>Pachygone ovata</i> . <i>Heterocycles</i> , 1983, 20, 1927.	0.4	7
34	Screening for saponins using the blood hemolysis test: An undergraduate laboratory experiment. <i>Journal of Chemical Education</i> , 1988, 65, 161.	1.1	6
35	Calozeylanic acid, a new bark acid from three species (guttiferae). <i>Tetrahedron Letters</i> , 1981, 22, 5083-5086.	0.7	5
36	Quaesitol, a phenol from <i>Garcinia quaesita</i> . <i>Phytochemistry</i> , 1984, 23, 2679-2681.	1.4	4

#	ARTICLE	IF	CITATIONS
37	Identification of character impact odorants in coriander and wild coriander leaves using GC-olfactometry and GC x GC-TOFMS. Developments in Food Science, 2006, 43, 197-200.	0.0	3