

# Beverley Shields

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/11270442/publications.pdf>

Version: 2024-02-01

19  
papers

12,136  
citations

394286

19  
h-index

752573

20  
g-index

20  
all docs

20  
docs citations

20  
times ranked

16031  
citing authors

#	ARTICLE	IF	CITATIONS
1	A Type 1 Diabetes Genetic Risk Score Can Aid Discrimination Between Type 1 and Type 2 Diabetes in Young Adults. <i>Diabetes Care</i> , 2016, 39, 337-344.	4.3	231
2	Mendelian Randomization Studies Do Not Support a Role for Raised Circulating Triglyceride Levels Influencing Type 2 Diabetes, Glucose Levels, or Insulin Resistance. <i>Diabetes</i> , 2011, 60, 1008-1018.	0.3	77
3	Genetic evidence that raised sex hormone binding globulin (SHBG) levels reduce the risk of type 2 diabetes. <i>Human Molecular Genetics</i> , 2010, 19, 535-544.	1.4	176
4	Polygenic Risk Variants for Type 2 Diabetes Susceptibility Modify Age at Diagnosis in Monogenic <i>HNFA</i> Diabetes. <i>Diabetes</i> , 2010, 59, 266-271.	0.3	37
5	New genetic loci implicated in fasting glucose homeostasis and their impact on type 2 diabetes risk. <i>Nature Genetics</i> , 2010, 42, 105-116.	9.4	1,982
6	A common genetic variant in the 15q24 nicotinic acetylcholine receptor gene cluster (CHRNA5-CHRNA3-CHRNA4) is associated with a reduced ability of women to quit smoking in pregnancy. <i>Human Molecular Genetics</i> , 2009, 18, 2922-2927.	1.4	132
7	Type 2 Diabetes Risk Alleles Are Associated With Reduced Size at Birth. <i>Diabetes</i> , 2009, 58, 1428-1433.	0.3	135
8	Cigarette Smoking during Pregnancy Is Associated with Alterations in Maternal and Fetal Thyroid Function. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2009, 94, 570-574.	1.8	64
9	Meta-analysis of genome-wide association data and large-scale replication identifies additional susceptibility loci for type 2 diabetes. <i>Nature Genetics</i> , 2008, 40, 638-645.	9.4	1,683
10	Common Variation in the <i>FTO</i> Gene Alters Diabetes-Related Metabolic Traits to the Extent Expected Given Its Effect on BMI. <i>Diabetes</i> , 2008, 57, 1419-1426.	0.3	277
11	A Common Variation in Deiodinase 1 Gene <i>DIO1</i> Is Associated with the Relative Levels of Free Thyroxine and Triiodothyronine. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2008, 93, 3075-3081.	1.8	133
12	Detection of Thyroid Dysfunction in Early Pregnancy: Universal Screening or Targeted High-Risk Case Finding?. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007, 92, 203-207.	1.8	362
13	Type 2 Diabetes <i>TCF7L2</i> Risk Genotypes Alter Birth Weight: A Study of 24,053 Individuals. <i>American Journal of Human Genetics</i> , 2007, 80, 1150-1161.	2.6	112
14	A Common Variant in the <i>FTO</i> Gene Is Associated with Body Mass Index and Predisposes to Childhood and Adult Obesity. <i>Science</i> , 2007, 316, 889-894.	6.0	3,884
15	Replication of Genome-Wide Association Signals in UK Samples Reveals Risk Loci for Type 2 Diabetes. <i>Science</i> , 2007, 316, 1336-1341.	6.0	2,040
16	A common variant of <i>HMGA2</i> is associated with adult and childhood height in the general population. <i>Nature Genetics</i> , 2007, 39, 1245-1250.	9.4	373
17	Combining Information from Common Type 2 Diabetes Risk Polymorphisms Improves Disease Prediction. <i>PLoS Medicine</i> , 2006, 3, e374.	3.9	242
18	No Evidence of Association of <i>ENPP1</i> Variants With Type 2 Diabetes or Obesity in a Study of 8,089 U.K. Caucasians. <i>Diabetes</i> , 2006, 55, 3175-3179.	0.3	86

#	ARTICLE	IF	CITATIONS
19	Common Variants of the Hepatocyte Nuclear Factor-4 P2 Promoter Are Associated With Type 2 Diabetes in the U.K. Population. <i>Diabetes</i> , 2004, 53, 3002-3006.	0.3	88