

Shin-ichi Nakano

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/112867/publications.pdf>

Version: 2024-02-01

108
papers

1,913
citations

304743

22
h-index

361022

35
g-index

115
all docs

115
docs citations

115
times ranked

2090
citing authors

#	ARTICLE	IF	CITATIONS
1	Biogeochemical characteristics of the HÃ¶vsgÃ¶lÃ¶“Ustilinsk water system in Mongolia and Russia: the effect of environmental factors on dissolved chemical components. <i>Limnology</i> , 2022, 23, 385-402.	1.5	2
2	Ecogenomics sheds light on diverse lifestyle strategies in freshwater CPR. <i>Microbiome</i> , 2022, 10, .	11.1	22
3	The crucial influence of trophic status on the relative requirement of nitrogen to phosphorus for phytoplankton growth. <i>Water Research</i> , 2022, 222, 118868.	11.3	10
4	Seasonal changes in the cell size and density of the diatom <i>Fragilaria crotonensis</i> Kitton in Lake Biwa. , 2022, 77, 3469-3476.		1
5	Microdiversity and phylogeographic diversification of bacterioplankton in pelagic freshwater systems revealed through long-read amplicon sequencing. <i>Microbiome</i> , 2021, 9, 24.	11.1	17
6	Phylogenetic diversity of the picocyanobacterial community from a novel winter bloom in Lake Biwa. <i>Limnology</i> , 2021, 22, 161-167.	1.5	4
7	The <scp>Asiaâ€Pacifi</scp> Biodiversity Observation Network: 10â€year achievements and new strategies to 2030. <i>Ecological Research</i> , 2021, 36, 232-257.	1.5	11
8	Long-term variation in abundance of the non-native phytoplankton <i>Micrasterias hardyi</i> (Zygnematophyceae, Streptophyta) in Lake Biwa, Japan. <i>Limnology</i> , 2020, 21, 67-72.	1.5	4
9	A freshwater radiation of diplomonads. <i>Environmental Microbiology</i> , 2020, 22, 4658-4668.	3.8	17
10	Influence of potential grazers on picocyanobacterial abundance in Lake Biwa revealed with empirical dynamic modeling. <i>Inland Waters</i> , 2020, 10, 386-396.	2.2	4
11	Distribution of the Harmful Bloom-Forming Cyanobacterium, <i>Microcystis aeruginosa</i>, in 88 Freshwater Environments across Japan. <i>Microbes and Environments</i> , 2020, 35, n/a.	1.6	6
12	Widespread Dominance of Kinetoplastids and Unexpected Presence of Diplomonads in Deep Freshwater Lakes. <i>Frontiers in Microbiology</i> , 2019, 10, 2375.	3.5	16
13	Genomeâ€resolved viral and cellular metagenomes revealed potential key virusâ€host interactions in a deep freshwater lake. <i>Environmental Microbiology</i> , 2019, 21, 4740-4754.	3.8	49
14	Rapid development and characterization of EST-SSR markers for the honey locust seed beetle, <i>Megabruchidius dorsalis</i> (Coleoptera: Bruchidae), using de novo transcriptome analysis based on next-generation sequencing. <i>Applied Entomology and Zoology</i> , 2019, 54, 141-145.	1.2	5
15	Trophic niche breadth of pond zooplankton species using stable isotope analysis and the relationship with the abiotic and biotic factors. <i>Royal Society Open Science</i> , 2019, 6, 180917.	2.4	3
16	Metaepigenomic analysis reveals the unexplored diversity of DNA methylation in an environmental prokaryotic community. <i>Nature Communications</i> , 2019, 10, 159.	12.8	48
17	Distributions and geochemical behaviors of oxyanion-forming trace elements and uranium in the HÃ¶vsgÃ¶lÃ¶“Baikalâ€Yenisei water system of Mongolia and Russia. <i>Journal of Geochemical Exploration</i> , 2018, 188, 123-136.	3.2	8
18	The effect of human activities on benthic macroinvertebrate diversity in tributary lagoons surrounding Lake Biwa. <i>Limnology</i> , 2018, 19, 199-207.	1.5	7

#	ARTICLE	IF	CITATIONS
19	The Broad Habitat Spectrum of the CL500-11 Lineage (Phylum Chloroflexi), a Dominant Bacterioplankton in Oxygenated Hypolimnia of Deep Freshwater Lakes. <i>Frontiers in Microbiology</i> , 2018, 9, 2891.	3.5	23
20	Hidden in plain sight—highly abundant and diverse planktonic freshwater Chloroflexi. <i>Microbiome</i> , 2018, 6, 176.	11.1	130
21	Ubiquity and quantitative significance of bacterioplankton lineages inhabiting the oxygenated hypolimnion of deep freshwater lakes. <i>ISME Journal</i> , 2017, 11, 2279-2293.	9.8	75
22	Differential Responses of Two Ecologically Similar Case-Bearing Caddisfly Species to a Fish Chemical Cue: Implications for a Coexistence Mechanism. <i>Zoological Science</i> , 2017, 34, 461-467.	0.7	3
23	Growth and mortality rates of prokaryotes in the hypolimnion of a deep freshwater lake (Lake Biwa, Japan). <i>Journal of Plankton Research</i> , 2017, 39, 1078-1083.	2.2	2
24	Seasonal dynamics of heterotrophic and plastidic protists in the water column of Lake Biwa, Japan. <i>Aquatic Microbial Ecology</i> , 2017, 80, 123-137.	1.8	18
25	Vertical partitioning of freshwater bacterioplankton community in a deep mesotrophic lake with a fully oxygenated hypolimnion (Lake Biwa, Japan). <i>Environmental Microbiology Reports</i> , 2016, 8, 780-788.	2.4	52
26	Changes in bacterial community structure associated with phytoplankton succession in outdoor experimental ponds. <i>Plankton and Benthos Research</i> , 2015, 10, 34-44.	0.6	2
27	A novel alphaproteobacterial ectosymbiont promotes the growth of the hydrocarbon-rich green alga <i>Botryococcus braunii</i> . <i>Scientific Reports</i> , 2015, 5, 10467.	3.3	55
28	Identification of species and genotypic compositions of <i>Cryptomonas</i> (Cryptophyceae) populations in the eutrophic Lake Hira, Japan, using single-cell PCR. <i>Aquatic Ecology</i> , 2015, 49, 263-272.	1.5	2
29	Kinetoplastid flagellates overlooked by universal primers dominate in the oxygenated hypolimnion of Lake Biwa, Japan. <i>FEMS Microbiology Ecology</i> , 2015, 91, fiv083.	2.7	29
30	High contribution of <i>Synechococcus</i> to phytoplankton biomass in the aphotic hypolimnion in a deep freshwater lake (Lake Biwa, Japan). <i>Aquatic Microbial Ecology</i> , 2015, 75, 69-79.	1.8	13
31	Developing an understanding of dissolved organic matter dynamics in the giant Lake Baikal by ultrahigh resolution mass spectrometry. <i>Limnology</i> , 2014, 15, 127-139.	1.5	7
32	Biodiversity in Aquatic Systems and Environments. <i>SpringerBriefs in Biology</i> , 2014, , .	0.5	6
33	Protistan grazing and viral lysis losses of bacterial carbon production in a large mesotrophic lake (Lake Biwa). <i>Limnology</i> , 2014, 15, 257-270.	1.5	4
34	Progress in the 21st century: a Roadmap for the Ecological Society of Japan. <i>Ecological Research</i> , 2014, 29, 357-368.	1.5	6
35	High-throughput sequencing shows inconsistent results with a microscope-based analysis of the soil prokaryotic community. <i>Soil Biology and Biochemistry</i> , 2014, 76, 53-56.	8.8	13
36	Biodiversity Researches on Microbial Loop in Aquatic Systems. <i>SpringerBriefs in Biology</i> , 2014, , 51-67.	0.5	5

#	ARTICLE	IF	CITATIONS
37	Developing a Regional Network of Biodiversity Observation in the Asia-Pacific Region: Achievements and Challenges of AP BON. Structure and Function of Mountain Ecosystems in Japan, 2014, , 3-28.	0.5	4
38	Long-Term and Spatial Variation in the Diversity of Littoral Benthic Macroinvertebrate Fauna in Lake Biwa, Japan. Structure and Function of Mountain Ecosystems in Japan, 2014, , 151-166.	0.5	2
39	CARD-FISH analysis of prokaryotic community composition and abundance along small-scale vegetation gradients in a dry arctic tundra ecosystem. Soil Biology and Biochemistry, 2013, 64, 147-154.	8.8	6
40	Estimation of carbon biomass and community structure of planktonic bacteria in Lake Biwa using respiratory quinone analysis. Limnology, 2013, 14, 247-256.	1.5	5
41	Anatoxin-a-producing <i>Raphidiopsis mediterranea</i> Skuja var. <i>grandis</i> Hill is one ecotype of non-heterocytous <i>Cuspidothrix issatschenkoi</i> (UsaÅev) Rajaniemi et al. in Japanese lakes. Harmful Algae, 2013, 21-22, 44-53.	4.8	21
42	Seasonal dominance of CL500-11 bacterioplankton (phylum<i>Chloroflexi</i>) in the oxygenated hypolimnion of Lake Biwa, Japan. FEMS Microbiology Ecology, 2013, 83, 82-92.	2.7	69
43	Grazing impact on the cyanobacterium <i>Microcystis aeruginosa</i> by the heterotrophic flagellate <i>Collodictyon triciliatum</i> in an experimental pond. Limnology, 2013, 14, 43-49.	1.5	6
44	Genotypic composition and the relationship between genotypic composition and geographical proximity of the cyanobacterium <i>Microcystis aeruginosa</i> in western Japan. Canadian Journal of Microbiology, 2013, 59, 266-272.	1.7	2
45	PCR primers for selective detection of intra-species variations in the bloom-forming cyanobacterium, <i>Microcystis</i> . Harmful Algae, 2013, 23, 46-54.	4.8	14
46	Biogeochemical control on fluorescent dissolved organic matter dynamics in a large freshwater lake (Lake Biwa, Japan). Limnology and Oceanography, 2013, 58, 2262-2278.	3.1	23
47	Lack of Congruence in Species Diversity Indices and Community Structures of Planktonic Groups Based on Local Environmental Factors. PLoS ONE, 2013, 8, e69594.	2.5	9
48	Detection and identification of potentially toxic cyanobacteria: Ubiquitous distribution of <i>Microcystis aeruginosa</i> and <i>Cuspidothrix issatschenkoi</i> in Japanese lakes. Harmful Algae, 2012, 16, 49-57.	4.8	15
49	Abundance and bacterivory of heterotrophic nanoflagellates in the meromictic Lake Suigetsu, Japan. Aquatic Microbial Ecology, 2012, 66, 149-158.	1.8	12
50	Temporal variation in cyanobacteria species composition and photosynthetic activity in experimentally induced blooms. Journal of Plankton Research, 2011, 33, 1410-1416.	1.8	14
51	Response of the plankton community to herbicide application (triazine herbicide, simetryn) in a eutrophicated system: short-term exposure experiment using microcosms. Limnology, 2011, 12, 11-16.	1.5	10
52	Grazing on <i>Microcystis</i> (Cyanophyceae) by testate amoebae with special reference to cyanobacterial abundance and physiological state. Limnology, 2011, 12, 205-211.	1.5	4
53	Nitrogen and carbon isotope fractionations of zooplankton consumers in ponds: potential effects of seston C:N stoichiometry. Marine and Freshwater Research, 2011, 62, 66.	1.3	4
54	HISTORY OF JAPANESE LIMNOLOGY. Limnology and Oceanography Bulletin, 2010, 19, 78-82.	0.4	0

#	ARTICLE	IF	CITATIONS
55	Shoreline bank construction modifies benthic-pelagic coupling of food webs. <i>Ecological Engineering</i> , 2010, 36, 601-604.	3.6	12
56	Feeding habits of omnivorous <i>Asplanchna</i> : comparison of diet composition among <i>Asplanchna herricki</i> , <i>A. priodonta</i> and <i>A. girodi</i> in pond ecosystems. <i>Journal of Limnology</i> , 2010, 69, 209.	1.1	21
57	Trophic niche breadth variability differs among three <i>Neocalanus</i> species in the subarctic Pacific Ocean. <i>Journal of Plankton Research</i> , 2010, 32, 1733-1737.	1.8	9
58	Effects of nutrient supplies on the growth rates of planktonic bacteria in Uchiumi Bay, Japan. <i>Aquatic Biology</i> , 2010, 9, 123-130.	1.4	4
59	Dispersal, connectivity, and local conditions determine zooplankton community composition in artificially connected ponds. <i>Aquatic Biology</i> , 2010, 10, 47-55.	1.4	4
60	Resource availability and ecosystem size predict food-chain length in pond ecosystems. <i>Oikos</i> , 2009, 118, 138-144.	2.7	51
61	Grazing Effects on Toxic and Non-Toxic <i>Microcystis aeruginosa</i> by the Mixotrophic Flagellate <i>Ochromonas</i> . <i>Journal of Freshwater Ecology</i> , 2009, 24, 367-373.	1.2	10
62	Abundance and pigment type composition of picocyanobacteria in Barguzin Bay, Lake Baikal. <i>Limnology</i> , 2008, 9, 105-114.	1.5	11
63	Longitudinal changes in zooplankton distribution below a reservoir outfall with reference to river planktivory. <i>Limnology</i> , 2008, 9, 125-133.	1.5	53
64	Abundance and composition of the summer phytoplankton community along a transect from the Barguzin River to the central basin of Lake Baikal. <i>Limnology</i> , 2008, 9, 243-250.	1.5	6
65	Evaluation of three phytoplankton species as food for the pearl oyster <i>Pinctada fucata</i> . <i>Aquaculture International</i> , 2008, 16, 309-318.	2.2	7
66	Drifting plankton from a reservoir subsidize downstream food webs and alter community structure. <i>Oecologia</i> , 2008, 156, 363-371.	2.0	67
67	Changes in the abundance and composition of picophytoplankton in relation to the occurrence of a Kyucho and a bottom intrusion in the Bungo Channel, Japan. <i>Estuarine, Coastal and Shelf Science</i> , 2008, 76, 293-303.	2.1	9
68	Role of allochthonous organic matter in Lake Baikal investigated using a 3-dimensional fluorescence excitation-emission matrix spectroscopy and high performance liquid chromatography-mass spectrometry. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 2008, 30, 469-476.	0.1	0
69	Horizontal distribution and nutritional status of picophytoplankton in Lake Baikal in summer. <i>Verhandlungen Der Internationalen Vereinigung Fur Theoretische Und Angewandte Limnologie International Association of Theoretical and Applied Limnology</i> , 2008, 30, 598-602.	0.1	1
70	Attached microalgae contribute to planktonic food webs in bays with fish and pearl oyster farms. <i>Marine Ecology - Progress Series</i> , 2008, 353, 107-113.	1.9	14
71	Growth and grazing mortality rates of <i>Prochlorococcus</i> , <i>Synechococcus</i> and eukaryotic picophytoplankton in a bay of the Uwa Sea, Japan. <i>Journal of Plankton Research</i> , 2007, 30, 241-250.	1.8	27
72	Growth rates of <i>Synechococcus</i> types with different phycoerythrin composition estimated by dual-laser flow cytometry in relationship to the light environment in the Uwa Sea. <i>Journal of Sea Research</i> , 2006, 55, 182-190.	1.6	8

#	ARTICLE	IF	CITATIONS
73	Vertical profiles of current velocity and dissolved oxygen saturation in biofilms on artificial and natural substrates. <i>Limnology</i> , 2006, 7, 213-218.	1.5	6
74	Nutrient limitation of the primary production of phytoplankton in Lake Baikal. <i>Limnology</i> , 2006, 7, 225-229.	1.5	13
75	Assessing Primary and Bacterial Production Rates in Biofilms on Pebbles in Ishite Stream, Japan. <i>Microbial Ecology</i> , 2006, 52, 1-9.	2.8	18
76	Contribution of Chemoautotrophic Production to Freshwater Macroinvertebrates in a Headwater Stream Using Multiple Stable Isotopes. <i>International Review of Hydrobiology</i> , 2006, 91, 501-508.	0.9	15
77	Relative importance of nanoflagellates and ciliates as consumers of bacteria in a coastal sea area dominated by oligotrichous <i>Strombidium</i> and <i>Strobilidium</i> . <i>Aquatic Microbial Ecology</i> , 2006, 42, 139-147.	1.8	33
78	Grazing and growth of the heterotrophic flagellate <i>Diphyllia rotans</i> on the cyanobacterium <i>Microcystis aeruginosa</i> . <i>Aquatic Microbial Ecology</i> , 2006, 45, 163-170.	1.8	22
79	The dynamics of microbial and herbivorous food webs in a coastal sea with special reference to intermittent nutrient supply from bottom intrusion. <i>Aquatic Ecology</i> , 2005, 38, 485-493.	1.5	2
80	Abundance and Community Structure of Picoplankton and Protists in the Microbial Food Web of Barguzin Bay, Lake Baikal. <i>Aquatic Ecology</i> , 2005, 39, 263-270.	1.5	12
81	Abundance, growth and grazing loss rates of picophytoplankton in Barguzin Bay, Lake Baikal. <i>Aquatic Ecology</i> , 2005, 39, 431-438.	1.5	20
82	Seasonal changes in the abundance and composition of picophytoplankton in relation to the occurrence of 'Kyucho' and bottom intrusion in Uchiumi Bay, Japan. <i>Marine Ecology - Progress Series</i> , 2005, 298, 59-67.	1.9	24
83	An improved method for collecting heterotrophic microorganisms living on pebbles in streams. <i>Limnology</i> , 2004, 5, 41-46.	1.5	9
84	Trophic coupling of a testate amoeba and <i>Microcystis</i> species in a hypertrophic pond. <i>Limnology</i> , 2004, 5, 71.	1.5	21
85	The dynamics of microbial and herbivorous food webs in a coastal sea with special reference to intermittent nutrient supply from bottom intrusion. <i>Aquatic Ecology</i> , 2004, 38, 485-493.	1.5	9
86	Discrimination of Two Phycoerythrin-Pigment Types of <i>Synechococcus</i> and Their Seasonal Succession in the Uwa Sea. <i>Microbes and Environments</i> , 2004, 19, 7-12.	1.6	7
87	Title is missing!. <i>Aquatic Ecology</i> , 2003, 37, 37-43.	1.5	13
88	Vertical planktonic structure in the central basin of Lake Baikal in summer 1999, with special reference to the microbial food web. <i>Limnology</i> , 2003, 4, 155-160.	1.5	15
89	Dominance of <i>Microcystis</i> with Special Reference to Carbon Availability in Lake Water.. <i>Microbes and Environments</i> , 2003, 18, 38-42.	1.6	11
90	Effect of nutrient limitation on abundance and growth of phytoplankton in a Japanese pearl farm. <i>Marine Ecology - Progress Series</i> , 2003, 258, 43-50.	1.9	14

#	ARTICLE	IF	CITATIONS
91	Respiration rates of the Japanese pearl oyster, <i>Pinctada fucata martensii</i> , feeding on <i>Pavlova lutheri</i> and <i>Chaetoceros gracilis</i> . <i>Aquaculture Research</i> , 2002, 33, 33-36.	1.8	9
92	Effect of water temperature and chlorophyll abundance on shell growth of the Japanese pearl oyster, <i>Pinctada fucata martensii</i> , in suspended culture at different depths and sites. <i>Aquaculture Research</i> , 2002, 33, 109-116.	1.8	28
93	Effect of heterotrophic nanoflagellates on the loss of virus-like particles in pond water. <i>Ecological Research</i> , 2002, 17, 473-479.	1.5	21
94	Title is missing!. <i>Hydrobiologia</i> , 2002, 481, 181-185.	2.0	25
95	Cyanobacterial blooms in a shallow lake: a largescale enclosure assay to test the importance of diurnal stratification. <i>Fundamental and Applied Limnology</i> , 2001, 150, 491-509.	0.7	22
96	Trophic linkage among heterotrophic nanoflagellates, ciliates and metazoan zooplankton in a hypereutrophic pond. <i>Aquatic Microbial Ecology</i> , 2001, 25, 259-270.	1.8	27
97	Mass mortality of the Japanese pearl oyster <i>Pinctada fucata martensii</i> in relation to water temperature, chlorophyll a and phytoplankton composition. <i>Diseases of Aquatic Organisms</i> , 2001, 44, 61-68.	1.0	46
98	Changes in cell volume of bacteria and heterotrophic nanoflagellates in a hypereutrophic pond. <i>Hydrobiologia</i> , 2000, 428, 197-203.	2.0	10
99	The Vertical Distribution of Pearl Oyster <i>Pinctada fucata martensii</i> Spat in Uchiumi Bay. <i>Fisheries Science</i> , 1999, 65, 358-361.	1.6	8
100	Title is missing!. <i>Hydrobiologia</i> , 1999, 411, 211-216.	2.0	40
101	Seasonal changes in abundance of heterotrophic nanoflagellates and their consumption of bacteria in Lake Biwa with special reference to trophic interactions with <i>Daphnia galeata</i> . <i>Fundamental and Applied Limnology</i> , 1998, 142, 21-34.	0.7	24
102	Trophic roles of heterotrophic nanoflagellates and ciliates among planktonic organisms in a hypereutrophic pond. <i>Aquatic Microbial Ecology</i> , 1998, 16, 153-161.	1.8	57
103	Bacterial response to extracellular dissolved organic carbon released from healthy and senescent <i>Fragilaria crotonensis</i> (Bacillariophyceae) in experimental systems. <i>Hydrobiologia</i> , 1996, 339, 47-55.	2.0	18
104	Seasonal Changes in Horizontal Distribution of Algal Picoplankton in Lake Biwa with Special Reference to Water Temperature, Nutrient Levels and Heterotrophic Flagellates.. <i>Japanese Journal of Limnology</i> , 1996, 57, 49-55.	0.1	11
105	Carbon: nitrogen: phosphorus ratios and nutrient regeneration of a heterotrophic flagellate fed on bacteria with different elemental ratios. <i>Archiv für Hydrobiologie</i> , 1994, 129, 257-271.	1.1	49
106	Rates and Ratios of Nitrogen and Phosphorus Released by a Bacterivorous Flagellate.. <i>Japanese Journal of Limnology</i> , 1994, 55, 115-123.	0.1	10
107	Estimation of Phosphorus Release Rate by Bacterivorous Flagellates in Lake Biwa.. <i>Japanese Journal of Limnology</i> , 1994, 55, 201-211.	0.1	15
108	Changes in Bacterioplankton Production and Dominant Algal Species in the North Basin of Lake Biwa.. <i>Japanese Journal of Limnology</i> , 1992, 53, 145-149.	0.1	9