Harlan F Weisman

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/11296608/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Pharmacodynamic Profile of Short-term Abciximab Treatment Demonstrates Prolonged Platelet Inhibition With Gradual Recovery From GP IIb/IIIa Receptor Blockade. Circulation, 1998, 97, 1680-1688.	1.6	282
2	Bleeding Complications With the Chimeric Antibody to Platelet Glycoprotein IIb/IIIa Integrin in Patients Undergoing Percutaneous Coronary Intervention. Circulation, 1995, 91, 2882-2890.	1.6	216
3	Effects of platelet glycoprotein IIb/IIIa receptor blockade by a chimeric monoclonal antibody (Abciximab) on acute and six-month outcomes after percutaneous transluminal coronary angioplasty for acute myocardial infarction. American Journal of Cardiology, 1996, 77, 1045-1051.	0.7	205
4	Evidence for Prevention of Death and Myocardial Infarction With Platelet Membrane Glycoprotein IIb/IIIa Receptor Blockade by Abciximab (c7E3 Fab) Among Patients With Unstable Angina Undergoing Percutaneous Coronary Revascularization fn1fn1This study was supported by Centocor, Inc., Malvern, Pennsylvania Journal of the American College of Cardiology, 1997, 30, 149-156.	1.2	204
5	Differential inhibition of platelet aggregation induced by adenosine diphosphate or a thrombin receptor-activating peptide in patients treated with bolus chimeric 7E3 Fab: Implications for inhibition of the internal pool of GPIIb/IIIa receptors. Journal of the American College of Cardiology, 1995, 26, 1665-1671.	1.2	158
6	Sustained Suppression of Ischemic Complications of Coronary Intervention by Platelet GP IIb/IIIa Blockade With Abciximab. Circulation, 1999, 99, 1951-1958.	1.6	154
7	Effect of platelet glycoprotein IIb/IIIa integrin blockade on activated clotting time during percutaneous transluminal coronary angioplasty or directional atherectomy (the EPIC trial). American Journal of Cardiology, 1995, 75, 559-562.	0.7	147
8	Monoclonal Antibodies to Platelet Glycoprotein IIb/IIIa as Antithrombotic Agentsa. Annals of the New York Academy of Sciences, 1991, 614, 193-213.	1.8	145
9	Standard Versus Low-Dose Weight-Adjusted Heparin in Patients Treated With the Platelet Glycoprotein IIb/IIIa Receptor Antibody Fragment Abciximab (c7E3 Fab) During Percutaneous Coronary Revascularization. American Journal of Cardiology, 1997, 79, 286-291.	0.7	137
10	Long-term mortality benefit with abciximab in patients undergoing percutaneous coronary intervention. Journal of the American College of Cardiology, 2001, 37, 2059-2065.	1.2	136
11	The immunogenicity of the 7E3 murine monoclonal Fab antibody fragment variable region is dramatically reduced in humans by substitution of human for murine constant regions. Molecular Immunology, 1995, 32, 1271-1281.	1.0	118
12	Therapeutic heparin concentrations augment platelet reactivity: Implications for the pharmacologic assessment of the glycoprotein IIb/IIIa antagonist abciximab. American Heart Journal, 2000, 139, 696-703.	1.2	42
13	Concept and Clinical Application of Platelet Glycoprotein IIb/IIIa Inhibition with Abciximab (c7E3 Fab;) Tj ETQq1	1 0.78431 0.7	4 rgBT /Overlo 7
14	An industry perspective on health economics studies. American Heart Journal, 1999, 137, S129-S132.	1.2	3
15	Abciximab: The First Platelet Glycoprotein IIb/IIIa Receptor Antagonist. , 1999, , 35-71.		2