

J Fernando Bazan

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/11320752/j-fernando-bazan-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

15
papers

9,516
citations

15
h-index

15
g-index

15
ext. papers

10,492
ext. citations

24.3
avg, IF

4.91
L-index

#	Paper	IF	Citations
15	Hedgehog pathway modulation by multiple lipid binding sites on the smoothed effector of signal response. <i>Developmental Cell</i> , 2013 , 26, 346-57	10.2	137
14	Structure of the activating IL-1 receptor signaling complex. <i>Nature Structural and Molecular Biology</i> , 2012 , 19, 455-7	17.6	80
13	Mapping the NPHP-JBTS-MKS protein network reveals ciliopathy disease genes and pathways. <i>Cell</i> , 2011 , 145, 513-28	56.2	435
12	Functional characterization of putative cilia genes by high-content analysis. <i>Molecular Biology of the Cell</i> , 2011 , 22, 1104-19	3.5	56
11	The conserved Bardet-Biedl syndrome proteins assemble a coat that traffics membrane proteins to cilia. <i>Cell</i> , 2010 , 141, 1208-19	56.2	456
10	Smoothed mutation confers resistance to a Hedgehog pathway inhibitor in medulloblastoma. <i>Science</i> , 2009 , 326, 572-4	33.3	676
9	Structure of IL-33 and its interaction with the ST2 and IL-1RAcP receptors--insight into heterotrimeric IL-1 signaling complexes. <i>Structure</i> , 2009 , 17, 1398-410	5.2	132
8	A BBSome subunit links ciliogenesis, microtubule stability, and acetylation. <i>Developmental Cell</i> , 2008 , 15, 854-65	10.2	225
7	A core complex of BBS proteins cooperates with the GTPase Rab8 to promote ciliary membrane biogenesis. <i>Cell</i> , 2007 , 129, 1201-13	56.2	1037
6	IL-33, an interleukin-1-like cytokine that signals via the IL-1 receptor-related protein ST2 and induces T helper type 2-associated cytokines. <i>Immunity</i> , 2005 , 23, 479-90	32.3	2643
5	WSX-1 and glycoprotein 130 constitute a signal-transducing receptor for IL-27. <i>Journal of Immunology</i> , 2004 , 172, 2225-31	5.3	546
4	IL-27, a heterodimeric cytokine composed of EBI3 and p28 protein, induces proliferation of naive CD4+ T cells. <i>Immunity</i> , 2002 , 16, 779-90	32.3	1100
3	Two novel IL-1 family members, IL-1 delta and IL-1 epsilon, function as an antagonist and agonist of NF-kappa B activation through the orphan IL-1 receptor-related protein 2. <i>Journal of Immunology</i> , 2001 , 167, 1440-6	5.3	211
2	Solution structure and dynamics of the CX3C chemokine domain of fractalkine and its interaction with an N-terminal fragment of CX3CR1. <i>Biochemistry</i> , 1999 , 38, 1402-14	3.2	128
1	A new class of membrane-bound chemokine with a CX3C motif. <i>Nature</i> , 1997 , 385, 640-4	50.4	1654